

Solutionbank

Edexcel AS and A Level Modular Mathematics



Exercise A, Question 1
Question:

Draw mapping diagrams and graphs for the following operations:



- 'subtract 5' on the set $\{ 10, 5, 0, -5, x \}$
- 'double and add 3' on the set $\{ -2, 2, 4, 6, x \}$
- 'square and then subtract 1' on the set $\{ -3, -1, 0, 1, 3, x \}$
- 'the positive square root' on the set $\{ -4, 0, 1, 4, 9, x \}$.

Solution:



(a)





(b)



(c)



(d)



Note: You cannot take the square root of a negative number.




Solutionbank


Edexcel AS and A Level Modular Mathematics

Exercise A, Question 2

Question:

Find the missing numbers a to h in the following mapping diagrams:



**Solution:**

$$x \rightarrow 2x \quad \text{is 'doubling'}$$

$$-3 \rightarrow a \quad \text{so } a = -6$$

$$b \rightarrow 9 \quad \text{so } b \times 2 = 9 \quad \Rightarrow \quad b = 4 \frac{1}{2}$$

$$x \rightarrow x^2 - 9 \quad \text{is 'squaring then subtracting 9'}$$

$$3 \rightarrow c \quad \text{so } c = 3^2 - 9 = 0$$

$$d \rightarrow 0 \quad \text{so } d^2 - 9 = 0 \quad \Rightarrow \quad d^2 = 9 \quad \Rightarrow \quad d = \pm 3$$

$$x \rightarrow \frac{x-3}{4} \quad \text{is 'subtract 3, then divide by 4'}$$

$$10 \rightarrow e \quad \text{so } e = (10 - 3) \div 4 = 1.75$$

$$f \rightarrow 5 \quad \text{so } \frac{f-3}{4} = 5 \quad \Rightarrow \quad f = 23$$

$$x \rightarrow +\sqrt{x-3} \quad \text{is 'subtract 3, then take the positive square root'}$$

$$27 \rightarrow g \quad \text{so } g = +\sqrt{27-3} = +\sqrt{24} = +2\sqrt{6}$$

$$h \rightarrow +\sqrt{3} \quad \text{so } \sqrt{h-3} = \sqrt{3} \quad \Rightarrow \quad h-3=3 \quad \Rightarrow \quad h=6$$

$$\text{So } a = -6, b = 4 \frac{1}{2}, c = 0, d = \pm 3, e = 1.75, f = 23, g = 2\sqrt{6}, h = 6$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 1**Question:**

Find:

- (a) $f(3)$ where $f(x) = 5x + 1$
- (b) $g(-2)$ where $g(x) = 3x^2 - 2$
- (c) $h(0)$ where $h : x \rightarrow 3^x$
- (d) $j(-2)$ where $j : x \rightarrow 2^{-x}$

Solution:

$$(a) f(x) = 5x + 1$$

$$\text{Substitute } x = 3 \Rightarrow f(3) = 5 \times 3 + 1 = 16$$

$$(b) g(x) = 3x^2 - 2$$

$$\text{Substitute } x = -2 \Rightarrow g(-2) = 3 \times (-2)^2 - 2 = 3 \times 4 - 2 = 10$$

$$(c) h(x) = 3^x$$

$$\text{Substitute } x = 0 \Rightarrow h(0) = 3^0 = 1$$

$$(d) j(x) = 2^{-x}$$

$$\text{Substitute } x = -2 \Rightarrow j(-2) = 2^{-(-2)} = 2^2 = 4$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 2

Question:

Calculate the value(s) of a , b , c and d given that:

$$(a) p(a) = 16 \text{ where } p(x) = 3x - 2$$

$$(b) q(b) = 17 \text{ where } q(x) = x^2 - 3$$

$$(c) r(c) = 34 \text{ where } r(x) = 2(2^x) + 2$$

$$(d) s(d) = 0 \text{ where } s(x) = x^2 + x - 6$$

Solution:

$$(a) p(x) = 3x - 2$$

Substitute $x = a$ and $p(a) = 16$ then

$$16 = 3a - 2$$

$$18 = 3a$$

$$a = 6$$

$$(b) q(x) = x^2 - 3$$

Substitute $x = b$ and $q(b) = 17$ then

$$17 = b^2 - 3$$

$$20 = b^2$$

$$b = \pm\sqrt{20}$$

$$b = \pm 2\sqrt{5}$$

$$(c) r(x) = 2 \times 2^x + 2$$

Substitute $x = c$ and $r(c) = 34$ then

$$34 = 2 \times 2^c + 2$$

$$32 = 2 \times 2^c$$

$$16 = 2^c$$

$$c = 4$$

$$(d) s(x) = x^2 + x - 6$$

Substitute $x = d$ and $s(d) = 0$ then

$$0 = d^2 + d - 6$$

$$0 = (d + 3)(d - 2)$$

$$d = 2, -3$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 3

Question:

For the following functions

- (i) sketch the graph of the function
- (ii) state the range
- (iii) describe if the function is one-to-one or many-to-one.

(a) $m(x) = 3x + 2$

(b) $n(x) = x^2 + 5$


(c) $p(x) = \sin(x)$

(d) $q(x) = x^3$

Solution:

(a) $m(x) = 3x + 2$

(i)




(ii) Range of $m(x)$ is $-\infty < m(x) < \infty$

or $m(x) \in \mathbb{R}$ (all of the real numbers)

(iii) Function is one-to-one

(b) $n(x) = x^2 + 5$


(i)



- (ii) Range of $n(x)$ is $n(x) \geq 5$
 (iii) Function is many-to-one

(c) $p(x) = \sin(x)$


(i)



- (ii) Range of $p(x)$ is $-1 \leq p(x) \leq 1$
 (iii) Function is many-to-one

(d) $q(x) = x^3$

(i)



- (ii) Range of $q(x)$ is $-\infty < q(x) < \infty$ or $q(x) \in \mathbb{R}$
(iii) Function is one-to-one

Solutionbank


Edexcel AS and A Level Modular Mathematics

Exercise B, Question 4


Question:

State whether or not the following graphs represent functions. Give reasons for your answers and describe the type of function.


(a)




(b)




(c)




(d)




(e)




(f)

**Solution:**


(a)



One-to-one function




One-to-one function



Not a function.

The values left of $x = a$ do not get mapped anywhere.


The values right of $x = a$ get mapped to two values of y .




Not a function. Similar to part (c).

Values of x between $-r$ and $+r$ get mapped to two values of y .

Values outside this don't get mapped anywhere.



Not a function. The value $x = b$ doesn't get mapped anywhere.



Many-to-one function. Two values of x get mapped to the same value of y .

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 1

Question:

The functions below are defined for the discrete domains.

- (i) Represent each function on a mapping diagram, writing down the elements in the range.
- (ii) State if the function is one-to-one or many-to-one.

(a) $f(x) = 2x + 1$ for the domain $\{x = 1, 2, 3, 4, 5\}$.


(b) $g(x) = +\sqrt{x}$ for the domain $\{x = 1, 4, 9, 16, 25, 36\}$.

(c) $h(x) = x^2$ for the domain $\{x = -2, -1, 0, 1, 2\}$.

(d) $j(x) = \frac{2}{x}$ for the domain $\{x = 1, 2, 3, 4, 5\}$.


Solution:

(a) $f(x) = 2x + 1$ ‘Double and add 1’




(ii) One-to-one function

(b) $g(x) = +\sqrt{x}$ ‘The positive square root’




(ii) One-to-one function

(c) $h(x) = x^2$ ‘Square the numbers in the domain’



(ii) Many-to-one function

(d) $j(x) = \frac{2}{x}$ ‘2 divided by numbers in the domain’



(ii) One-to-one function

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 2

Question:

The functions below are defined for continuous domains.

- (i) Represent each function on a graph.
- (ii) State the range of the function.
- (iii) State if the function is one-to-one or many-to-one.

(a) $m(x) = 3x + 2$ for the domain $\{x > 0\}$.


(b) $n(x) = x^2 + 5$ for the domain $\{x \geq -2\}$.

(c) $p(x) = 2 \sin x$ for the domain $\{0 \leq x \leq 180\}$.

(d) $q(x) = +\sqrt{x+2}$ for the domain $\{x \geq -2\}$.

Solution:

(a) $m(x) = 3x + 2$ for $x > 0$




$3x + 2$ is a linear function of gradient 3 passing through 2 on the y axis.

(ii) $x = 0$ does not exist in the domain

So range is $m(x) > 3 \times 0 + 2 \Rightarrow m(x) > 2$

(iii) $m(x)$ is a one-to-one function

(b) $n(x) = x^2 + 5$ for $x \geq -2$



$x^2 + 5$ is a parabola with minimum point at (0, 5).


The domain however is only values bigger than or equal to 2.

(ii) $x = 2$ exists in the domain

$$\text{So range is } n(x) \geq 2^2 + 5 \Rightarrow n(x) \geq 9$$

(iii) $n(x)$ is a one-to-one function

(c) $p(x) = 2 \sin x$ for $0 \leq x \leq 180$




$2 \sin x$ has the same shape as $\sin x$ except that it has been stretched by a factor of 2 parallel to the y axis.

(ii) Range of $p(x)$ is $0 \leq p(x) \leq 2$

(iii) The function is many-to-one

(d) $q(x) = +\sqrt{x+2}$ for $x \geq -2$



$\sqrt{x+2}$ is the \sqrt{x} graph translated 2 units to the left.

(ii) The range of $q(x)$ is $q(x) \geq 0$

(iii) The function is one-to-one

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 3

Question:

The mappings $f(x)$ and $g(x)$ are defined by

$$f(x) = \begin{cases} 4 - x & x < 4 \\ x^2 + 9x & \geq 4 \end{cases}$$

$$g(x) = \begin{cases} 4 - x & x < 4 \\ x^2 + 9x & x > 4 \end{cases}$$

Explain why $f(x)$ is a function and $g(x)$ is not.

Sketch the function $f(x)$ and find


- (a) $f(3)$
- (b) $f(10)$
- (c) the value(s) of a such that $f(a) = 90$.

Solution:

$4 - x$ is a linear function of gradient -1 passing through 4 on the y axis.

$x^2 + 9$ is a \cup -shaped quadratic

At $x = 4$ $4 - x = 0$ and $x^2 + 9 = 25$




$g(x)$ is not a function because the element 4 of the domain does not get mapped anywhere.

In $f(x)$ it gets mapped to 25 .

(a) $f(3) = 4 - 3 = 1$ (Use $4 - x$ as $3 < 4$)

(b) $f(10) = 10^2 + 9 = 109$ (Use $x^2 + 9$ as $10 > 4$)

(c)



The negative value of a is where $4 - a = 90 \Rightarrow a = -86$

The positive value of a is where

$$a^2 + 9 = 90$$

$$a^2 = 81$$

$$a = \pm 9$$

$$a = 9$$

The values of a are -86 and 9 .

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 4
Question:


The function $s(x)$ is defined by

$$s(x) = \begin{cases} x^2 - 6 & x < 0 \\ 10 - x & x \geq 0 \end{cases}$$


- (a) Sketch $s(x)$.
- (b) Find the value(s) of a such that $s(a) = 43$.
- (c) Find the values of the domain that get mapped to themselves in the range.

Solution:

(a) $x^2 - 6$ is a \cup -shaped quadratic with a minimum value of $(0, -6)$.
 $10 - x$ is a linear function with gradient -1 passing through 10 on the y axis.



- (b) There is only one value of a such that $s(a) = 43$ (see graph).



$$s(a) = 43$$

$$a^2 - 6 = 43$$

$$a^2 = 49$$

$$a = \pm 7$$

Value is negative so $a = -7$

(c) If value gets mapped to itself then $s(b) = b$

For $10 - x$ part

$$10 - b = b$$

$$\Rightarrow 10 = 2b$$

$$\Rightarrow b = 5$$

Check. $s(5) = 10 - 5 = 5 \checkmark$

For $x^2 - 6$ part

$$b^2 - 6 = b$$

$$\Rightarrow b^2 - b - 6 = 0$$

$$\Rightarrow (b - 3)(b + 2) = 0$$

$$\Rightarrow b = 3, -2$$

b must be negative

$$\Rightarrow b = -2$$

Check. $s(-2) = (-2)^2 - 6 = 4 - 6 = -2 \checkmark$

Values that get mapped to themselves are -2 and 5 .

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 5

Question:

The function $g(x)$ is defined by $g(x) = cx + d$ where c and d are constants to be found. Given $g(3) = 10$ and $g(8) = 12$ find the values of c and d .

Solution:

$$g(x) = cx + d$$

$$g(3) = 10 \Rightarrow c \times 3 + d = 10$$

$$g(8) = 12 \Rightarrow c \times 8 + d = 12$$

$$3c + d = 10 \quad \textcircled{1}$$

$$8c + d = 12 \quad \textcircled{2}$$

$$\textcircled{2} - \textcircled{1}: \quad 5c = 2 \quad (\div 5)$$

$$\Rightarrow c = 0.4$$

Substitute $c = 0.4$ into $\textcircled{1}$:

$$3 \times 0.4 + d = 10$$

$$1.2 + d = 10$$

$$d = 8.8$$

$$\text{Hence } g(x) = 0.4x + 8.8$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 6

Question:

The function $f(x)$ is defined by $f(x) = ax^3 + bx - 5$ where a and b are constants to be found. Given that $f(1) = -4$ and $f(2) = 9$, find the values of the constants a and b .

Solution:

$$f(x) = ax^3 + bx - 5$$

$$f(1) = -4 \Rightarrow a \times 1^3 + b \times 1 - 5 = -4$$

$$\Rightarrow a + b - 5 = -4$$

$$\Rightarrow a + b = 1 \quad \textcircled{1}$$

$$f(2) = 9 \Rightarrow a \times 2^3 + b \times 2 - 5 = 9$$

$$\Rightarrow 8a + 2b - 5 = 9$$

$$\Rightarrow 8a + 2b = 14$$

$$\Rightarrow 4a + b = 7 \quad \textcircled{2}$$

$$\textcircled{2} - \textcircled{1}: \quad 3a = 6$$

$$\Rightarrow a = 2$$

Substitute $a = 2$ in $\textcircled{1}$:

$$2 + b = 1$$

$$b = -1$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 7


Question:

The function $h(x)$ is defined by $h(x) = x^2 - 6x + 20 \{ x \geq a \}$. Given that $h(x)$ is a one-to-one function find the smallest possible value of the constant a .


Solution:

$$h(x) = x^2 - 6x + 20 = (x - 3)^2 - 9 + 20 = (x - 3)^2 + 11$$

This is a \cup -shaped quadratic with minimum point at $(3, 11)$.



This is a many-to-one function.
For $h(x)$ to be one-to-one, $x \geq 3$



Hence smallest value of a is 3.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 1

Question:

Given the functions $f(x) = 4x + 1$, $g(x) = x^2 - 4$ and $h(x) = \frac{1}{x}$, find expressions for the functions:

(a) $fg(x)$

(b) $gf(x)$

(c) $gh(x)$

(d) $fh(x)$

(e) $f^2(x)$

Solution:

(a) $fg(x) = f(x^2 - 4) = 4(x^2 - 4) + 1 = 4x^2 - 15$

(b) $gf(x) = g(4x + 1) = (4x + 1)^2 - 4 = 16x^2 + 8x - 3$

(c) $gh(x) = g\left(\frac{1}{x}\right) = \left(\frac{1}{x}\right)^2 - 4 = \frac{1}{x^2} - 4$

(d) $fh(x) = f\left(\frac{1}{x}\right) = 4 \times \left(\frac{1}{x}\right) + 1 = \frac{4}{x} + 1$

(e) $f^2(x) = ff(x) = f(4x + 1) = 4(4x + 1) + 1 = 16x + 5$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 2

Question:

For the following functions $f(x)$ and $g(x)$, find the composite functions $fg(x)$ and $gf(x)$. In each case find a suitable domain and the corresponding range when

(a) $f(x) = x - 1$, $g(x) = x^2$

(b) $f(x) = x - 3$, $g(x) = +\sqrt{x}$

(c) $f(x) = 2^x$, $g(x) = x + 3$


Solution:

(a) $f(x) = x - 1$, $g(x) = x^2$

$$fg(x) = f(x^2) = x^2 - 1$$

Domain $x \in \mathbb{R}$


Range $fg(x) \geq -1$



$$gf(x) = g(x - 1) = (x - 1)^2$$

Domain $x \in \mathbb{R}$

Range $gf(x) \geq 0$




$$(b) f(x) = x - 3, g(x) = +\sqrt{x}$$

$$fg(x) = f(+\sqrt{x}) = \sqrt{x} - 3$$

$$\text{Domain } x \geq 0$$

(It will not be defined for negative numbers)


$$\text{Range } fg(x) \geq -3$$



$$gf(x) = g(x - 3) = \sqrt{x - 3}$$

$$\text{Domain } x \geq 3$$

$$\text{Range } gf(x) \geq 0$$




$$(c) f(x) = 2^x, g(x) = x + 3$$

$$fg(x) = f(x + 3) = 2^{x+3}$$

$$\text{Domain } x \in \mathbb{R}$$


$$\text{Range } fg(x) > 0$$



$$gf(x) = g(2^x) = 2^x + 3$$

Domain $x \in \mathbb{R}$

Range $gf(x) > 3$



Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 3

Question:

If $f(x) = 3x - 2$ and $g(x) = x^2$, find the number(s) a such that $fg(a) = gf(a)$.

Solution:

$$f(x) = 3x - 2, g(x) = x^2$$

$$fg(x) = f(x^2) = 3x^2 - 2$$

$$gf(x) = g(3x - 2) = (3x - 2)^2$$

$$\text{If } fg(a) = gf(a)$$

$$3a^2 - 2 = (3a - 2)^2$$

$$3a^2 - 2 = 9a^2 - 12a + 4$$

$$0 = 6a^2 - 12a + 6$$

$$0 = a^2 - 2a + 1$$

$$0 = (a - 1)^2$$

Hence $a = 1$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 4

Question:

Given that $s(x) = \frac{1}{x-2}$ and $t(x) = 3x + 4$ find the number m such that $ts(m) = 16$.

Solution:

$$s(x) = \frac{1}{x-2}, t(x) = 3x + 4$$

$$ts(x) = t\left(\frac{1}{x-2}\right) = 3 \times \left(\frac{1}{x-2}\right) + 4 = \frac{3}{x-2} + 4$$

If $ts(m) = 16$

$$\frac{3}{m-2} + 4 = 16 \quad (-4)$$

$$\frac{3}{m-2} = 12 \quad [\times (m-2)]$$

$$3 = 12(m-2) \quad (\div 12)$$

$$\frac{3}{12} = m-2$$

$$0.25 = m-2$$

$$m = 2.25$$

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 5

Question:

The functions $l(x)$, $m(x)$, $n(x)$ and $p(x)$ are defined by $l(x) = 2x + 1$, $m(x) = x^2 - 1$, $n(x) = \frac{1}{x+5}$ and $p(x) = x^3$. Find in terms of l , m , n and p the functions:

(a) $4x + 3$

(b) $4x^2 + 4x$

(c) $\frac{1}{x^2 + 4}$

(d) $\frac{2}{x+5} + 1$

(e) $(x^2 - 1)^3$

(f) $2x^2 - 1$

(g) x^{27}

Solution:

(a) $4x + 3 = 2(2x + 1) + 1 = 2l(x) + 1 = ll(x)$ [or $l^2(x)$]

(b) $4x^2 + 4x = (2x + 1)^2 - 1 = [l(x)]^2 - 1 = ml(x)$

(c) $\frac{1}{x^2 + 4} = \frac{1}{(x^2 - 1) + 5} = \frac{1}{m(x) + 5} = nm(x)$

(d) $\frac{2}{x+5} + 1 = 2 \times \frac{1}{x+5} + 1 = 2n(x) + 1 = ln(x)$

(e) $(x^2 - 1)^3 = [m(x)]^3 = pm(x)$

(f) $2x^2 - 1 = 2(x^2 - 1) + 1 = 2m(x) + 1 = lm(x)$

$$\begin{aligned}(g) \quad x^{27} &= [(x^3)^3]^3 = \{ [p(x)]^3 \}^3 = [pp(x)]^3 = ppp(x) \\ &= p^3(x)\end{aligned}$$

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 6

Question:

If $m(x) = 2x + 3$ and $n(x) = \frac{x-3}{2}$, prove that $mn(x) = x$.

Solution:

$$m(x) = 2x + 3, n(x) = \frac{x-3}{2}$$

$$mn(x) = m\left(\frac{x-3}{2}\right) = 2\left(\frac{x-3}{2}\right) + 3 = x - 3 + 3 = x$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 7

Question:

If $s(x) = \frac{3}{x+1}$ and $t(x) = \frac{3-x}{x}$, prove that $st(x) = x$.

Solution:

$$s(x) = \frac{3}{x+1}, t(x) = \frac{3-x}{x}$$

$$\begin{aligned} st(x) &= s\left(\frac{3-x}{x}\right) \\ &= \frac{3}{\frac{3-x}{x}+1} \times x \\ &= \frac{3x}{3-x+x} \\ &= \frac{3x}{3} \\ &= x \end{aligned}$$

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 8

Question:

If $f(x) = \frac{1}{x+1}$, prove that $f^2(x) = \frac{x+1}{x+2}$. Hence find an expression for $f^3(x)$.

Solution:

$$f(x) = \frac{1}{x+1}$$

$$\begin{aligned} ff(x) &= f\left(\frac{1}{x+1}\right) \\ &= \frac{1}{\frac{1}{x+1}+1} \times (x+1) \\ &= \frac{x+1}{1+x+1} \\ &= \frac{x+1}{x+2} \end{aligned}$$

$$\begin{aligned} f^3(x) = f[f^2(x)] &= f\left(\frac{x+1}{x+2}\right) \\ &= \frac{1}{\frac{x+1}{x+2}+1} \times (x+2) \\ &= \frac{x+2}{x+1+x+2} \\ &= \frac{x+2}{2x+3} \end{aligned}$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 1

Question:

For the following functions $f(x)$, sketch the graphs of $f(x)$ and $f^{-1}(x)$ on the same set of axes. Determine also the equation of $f^{-1}(x)$.

$$(a) f(x) = 2x + 3 \quad \{ x \in \mathbb{R} \}$$

$$(b) f(x) = \frac{x}{2} \quad \left\{ \begin{array}{l} x \in \mathbb{R} \end{array} \right\}$$

$$(c) f(x) = \frac{1}{x} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, x \neq 0 \end{array} \right\}$$

$$(d) f(x) = 4 - x \quad \{ x \in \mathbb{R} \}$$

$$(e) f(x) = x^2 + 2 \quad \{ x \in \mathbb{R}, x \geq 0 \}$$

$$(f) f(x) = x^3 \quad \{ x \in \mathbb{R} \}$$


Solution:

$$(a) \text{If } y = 2x + 3$$

$$y - 3 = 2x$$

$$\frac{y - 3}{2} = x$$


$$\text{Hence } f^{-1}(x) = \frac{x - 3}{2}$$



(b) If $y = \frac{x}{2}$

$$2y = x$$

$$\text{Hence } f^{-1}(x) = 2x$$




(c) If $y = \frac{1}{x}$

$$yx = 1$$

$$x = \frac{1}{y}$$

$$\text{Hence } f^{-1}(x) = \frac{1}{x}$$

Note that the inverse to the function is identical to the function.




(d) If $y = 4 - x$

$$x + y = 4$$

$$x = 4 - y$$

$$\text{Hence } f^{-1}(x) = 4 - x$$

Note that the inverse to the function is identical to the function.




(e) If $y = x^2 + 2$

$$y - 2 = x^2$$

$$\sqrt{y - 2} = x$$


$$\text{Hence } f^{-1}(x) = \sqrt{x - 2}$$



(f) If $y = x^3$

$$\sqrt[3]{y} = x$$

$$\text{Hence } f^{-1}(x) = \sqrt[3]{x}$$



Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 2

Question:

Determine which of the functions in Question 1 are self inverses. (That is to say the function and its inverse are identical.)

Solution:

Look back at Question 1.

$$1(c) f(x) = \frac{1}{x} \text{ and}$$

1(d) $f(x) = 4 - x$
are both identical to their inverses.

© Pearson Education Ltd 2008

Solutionbank


Edexcel AS and A Level Modular Mathematics

Exercise E, Question 3

Question:

Explain why the function $g(x) = 4 - x \{ x \in \mathbb{R}, x > 0 \}$ is not identical to its inverse.

Solution:



$$g(x) = 4 - x$$

has domain $x > 0$

and range $g(x) < 4$

$$\text{Hence } g^{-1}(x) = 4 - x$$

has domain $x < 4$

and range $g^{-1}(x) > 0$

Although $g(x)$ and $g^{-1}(x)$ have identical equations they act on different numbers and so are not identical. See graph.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 4

Question:

For the following functions $g(x)$, sketch the graphs of $g(x)$ and $g^{-1}(x)$ on the same set of axes. Determine the equation of $g^{-1}(x)$, taking care with its domain.

$$(a) g(x) = \frac{1}{x} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \geq 3 \end{array} \right\}$$

$$(b) g(x) = 2x - 1 \quad \{ x \in \mathbb{R}, \quad x \geq 0 \}$$


$$(c) g(x) = \frac{3}{x-2} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x > 2 \end{array} \right\}$$

$$(d) g(x) = \sqrt{x-3} \quad \{ x \in \mathbb{R}, \quad x \geq 7 \}$$

$$(e) g(x) = x^2 + 2 \quad \{ x \in \mathbb{R}, \quad x > 4 \}$$

$$(f) g(x) = x^3 - 8 \quad \{ x \in \mathbb{R}, \quad x \leq 2 \}$$

Solution:




$$g(x) = \frac{1}{x} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \geq 3 \end{array} \right\}$$

has range $g(x) \in \mathbb{R}, \quad 0 < g(x) \leq \frac{1}{3}$

Changing the subject of the formula gives

$$g^{-1}(x) = \frac{1}{x} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad 0 < x \leq \frac{1}{3} \end{array} \right\}$$

(b)




$$g(x) = 2x - 1 \quad \{ x \in \mathbb{R}, \quad x \geq 0 \}$$

has range $g(x) \in \mathbb{R}, \quad g(x) \geq -1$

Changing the subject of the formula gives

$$g^{-1}(x) = \frac{x + 1}{2} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \geq -1 \end{array} \right\}$$



$$g(x) = \frac{3}{x-2} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x > 2 \end{array} \right\}$$

has range $g(x) \in \mathbb{R}, \quad g(x) > 0$

Changing the subject of the formula gives

$$y = \frac{3}{x-2}$$


$$y(x-2) = 3$$

$$x-2 = \frac{3}{y}$$

$$x = \frac{3}{y} + 2 \quad \left(\text{or } \frac{3+2y}{y} \right)$$

$$\text{Hence } g^{-1}(x) = \frac{3}{x} + 2 \quad \left(\text{or } \frac{3+2x}{x} \right)$$

$$\{ x \in \mathbb{R}, \quad x > 0 \}$$



$$g(x) = \sqrt{x - 3} \quad \{ x \in \mathbb{R}, x \geq 7 \}$$

has range $g(x) \in \mathbb{R}, g(x) \geq 2$


Changing the subject of the formula gives

$$y = \sqrt{x - 3}$$

$$y^2 = x - 3$$

$$x = y^2 + 3$$

Hence $g^{-1}(x) = x^2 + 3$ with domain $x \in \mathbb{R}, x \geq 2$




$$g(x) = x^2 + 2 \quad \{ x \in \mathbb{R}, x > 4 \}$$

has range $g(x) \in \mathbb{R}, g(x) > 18$

Changing the subject of the formula gives

$$g^{-1}(x) = \sqrt{x - 2} \text{ with domain } x \in \mathbb{R}, x > 18$$

(f)



$$g(x) = x^3 - 8 \quad \{ x \in \mathbb{R}, x \leq 2 \}$$

has range $g(x) \in \mathbb{R}, g(x) \leq 0$

Changing the subject of the formula gives

$$y = x^3 - 8$$

$$y + 8 = x^3$$

$$\sqrt[3]{y + 8} = x$$

$$\text{Hence } g^{-1}(x) = \sqrt[3]{x + 8} \text{ with domain } x \in \mathbb{R}, x \leq 0$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 5

Question:

The function $m(x)$ is defined by $m(x) = x^2 + 4x + 9 \quad \{ x \in \mathbb{R}, x > a \}$ for some constant a . If $m^{-1}(x)$ exists, state the least value of a and hence determine the equation of $m^{-1}(x)$. State its domain.

Solution:


$$m(x) = x^2 + 4x + 9 \quad \{ x \in \mathbb{R}, x > a \} .$$

$$\text{Let } y = x^2 + 4x + 9$$

$$y = (x+2)^2 - 4 + 9$$


$$y = (x+2)^2 + 5$$

This has a minimum value of $(-2, 5)$.



For $m(x)$ to have an inverse it must be one-to-one.

Hence the least value of a is -2 .



$m(x)$ would have a range of $m(x) \in \mathbb{R}, m(x) > 5$

Changing the subject of the formula gives

$$y = (x + 2)^2 + 5$$

$$y - 5 = (x + 2)^2$$

$$\sqrt{y - 5} = x + 2$$

$$\sqrt{y - 5} - 2 = x$$

Hence $m^{-1}(x) = \sqrt{x - 5} - 2$ with domain $x \in \mathbb{R}, x > 5$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 6

Question:

Determine $t^{-1}(x)$ if the function $t(x)$ is defined by $t(x) = x^2 - 6x + 5$
 $\{ x \in \mathbb{R}, x \geq 5 \}$.

Solution:

$$t(x) = x^2 - 6x + 5 \quad \{ x \in \mathbb{R}, x \geq 5 \}$$

Let $y = x^2 - 6x + 5$ (complete the square)

$$y = (x - 3)^2 - 9 + 5$$

$$y = (x - 3)^2 - 4$$

This has a minimum point at $(3, -4)$.

Note. Since $x \geq 5$ is the domain, $t(x)$ is a one-to-one function.


Change the subject of the formula to find $t^{-1}(x)$:

$$y = (x - 3)^2 - 4$$

$$y + 4 = (x - 3)^2$$

$$\sqrt{y + 4} = x - 3$$

$$\sqrt{y + 4} + 3 = x$$



$$t(x) = x^2 - 6x + 5 \quad \{ x \in \mathbb{R}, x \geq 5 \}$$

has range $t(x) \in \mathbb{R}, t(x) \geq 0$

So $t^{-1}(x) = \sqrt{x + 4} + 3$ and has domain $x \in \mathbb{R}, x \geq 0$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 7

Question:


The function $h(x)$ is defined by $h(x) = \frac{2x+1}{x-2} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \neq 2 \end{array} \right\}.$

- (a) What happens to the function as x approaches 2?
- (b) Find $h^{-1}(3)$.
- (c) Find $h^{-1}(x)$, stating clearly its domain.
- (d) Find the elements of the domain that get mapped to themselves by the function.

Solution:

(a) As $x \rightarrow 2$ $h(x) \rightarrow \frac{5}{0}$ and hence $h(x) \rightarrow \infty$

(b) To find $h^{-1}(3)$ we can find what element of the domain gets mapped to 3.



$$\text{So } h(a) = 3$$

$$\frac{2a+1}{a-2} = 3$$

$$2a + 1 = 3a - 6$$

$$7 = a$$

$$\text{So } h^{-1}(3) = 7$$

(c) Let $y = \frac{2x+1}{x-2}$ and find x as a function of y .

$$y(x - 2) = 2x + 1$$

$$yx - 2y = 2x + 1$$

$$yx - 2x = 2y + 1$$

$$x(y - 2) = 2y + 1$$

$$x = \frac{2y + 1}{y - 2}$$

$$\text{So } h^{-1}(x) = \frac{2x + 1}{x - 2} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \neq 2 \end{array} \right\}$$

Hence the inverse function has exactly the same equation as the function. **But** the elements don't get mapped to themselves, see part (b).

(d) For elements to get mapped to themselves

$$h(b) = b$$

$$\frac{2b + 1}{b - 2} = b$$

$$2b + 1 = b(b - 2)$$

$$2b + 1 = b^2 - 2b$$

$$0 = b^2 - 4b - 1$$

$$b = \frac{4 \pm \sqrt{16 + 4}}{2} = \frac{4 \pm \sqrt{20}}{2} = \frac{4 \pm 2\sqrt{5}}{2} = 2 \pm \sqrt{5}$$

The elements $2 + \sqrt{5}$ and $2 - \sqrt{5}$ get mapped to themselves by the function.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 8

Question:

The function $f(x)$ is defined by $f(x) = 2x^2 - 3 \quad \{ x \in \mathbb{R}, x < 0 \}$.

Determine

(a) $f^{-1}(x)$ clearly stating its domain

(b) the values of a for which $f(a) = f^{-1}(a)$.

Solution:

(a) Let $y = 2x^2 - 3$

$$y + 3 = 2x^2$$

$$\frac{y+3}{2} = x^2$$

$$\sqrt{\frac{y+3}{2}} = x$$


The domain of $f^{-1}(x)$ is the range of $f(x)$.

$f(x) = 2x^2 - 3 \quad \{ x \in \mathbb{R}, x < 0 \}$

has range $f(x) > -3$

Hence $f^{-1}(x)$ must be the **negative** square root

$f^{-1}(x) = -\sqrt{\frac{x+3}{2}}$ has domain $x \in \mathbb{R}, x > -3$



(b) If $f(a) = f^{-1}(a)$ then a is negative (see graph).
Solve $f(a) = a$

$$2a^2 - 3 = a$$

$$2a^2 - a - 3 = 0$$

$$(2a - 3)(a + 1) = 0$$

$$a = \frac{3}{2}, -1$$

Therefore $a = -1$

Solutionbank

Edexcel AS and A Level Modular Mathematics


Exercise F, Question 1

Question:


Categorise the following as




- (i) not a function
- (ii) a one-to-one function
- (iii) a many-to-one function.

(a)




(b)






(f)


**Solution:**

(a) not a function




x value a gets mapped to two values of y .
 x value b gets mapped to no values


(b) one-to-one function




(c) many-to-one function



(d) many-to-one function




(e) not a function



x value a doesn't get mapped to any value of y .
It could be redefined as a function if the domain is said to exclude point a .

(f) not a function



x values less than a don't get mapped anywhere.
Again we could define the domain to be $x \geq a$ and then it would be a function.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 2

Question:

The following functions $f(x)$, $g(x)$ and $h(x)$ are defined by

$$f(x) = 4(x - 2) \quad \{ x \in \mathbb{R}, x \geq 0 \}$$

$$g(x) = x^3 + 1 \quad \{ x \in \mathbb{R} \}$$

$$h(x) = 3^x \quad \{ x \in \mathbb{R} \}$$

- (a) Find $f(7)$, $g(3)$ and $h(-2)$.
- (b) Find the range of $f(x)$ and the range of $g(x)$.
- (c) Find $g^{-1}(x)$.
- (d) Find the composite function $fg(x)$.
- (e) Solve $gh(a) = 244$.

Solution:

$$(a) f(7) = 4(7 - 2) = 4 \times 5 = 20$$


$$g(3) = 3^3 + 1 = 27 + 1 = 28$$

$$h(-2) = 3^{-2} = \frac{1}{3^2} = \frac{1}{9}$$

$$(b) f(x) = 4(x - 2) = 4x - 8$$


This is a straight line with gradient 4 and intercept -8 .

The domain tells us that $x \geq 0$.



The range of $f(x)$ is $f(x) \in \mathbb{R}, f(x) \geq -8$

$$g(x) = x^3 + 1$$



The range of $g(x)$ is $g(x) \in \mathbb{R}$

(c) Let $y = x^3 + 1$ (change the subject of the formula)

$$y - 1 = x^3$$

$$\sqrt[3]{y - 1} = x$$

$$\text{Hence } g^{-1}(x) = \sqrt[3]{x - 1} \quad \{ x \in \mathbb{R} \}$$

$$(d) fg(x) = f(x^3 + 1) = 4(x^3 + 1 - 2) = 4(x^3 - 1)$$

(e) Find $gh(x)$ first.

$$gh(x) = g(3^x) = (3^x)^3 + 1 = 3^{3x} + 1$$

$$\text{If } gh(a) = 244$$

$$3^{3a} + 1 = 244$$

$$3^{3a} = 243$$

$$3^{3a} = 3^5$$

$$3a = 5$$

$$a = \frac{5}{3}$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 3

Question:


The function $n(x)$ is defined by

$$n(x) = \begin{cases} 5 - x & x \leq 0 \\ x^2 & x > 0 \end{cases}$$

(a) Find $n(-3)$ and $n(3)$.

(b) Find the value(s) of a such that $n(a) = 50$.

Solution:




$y = 5 - x$ is a straight line with gradient -1 passing through 5 on the y axis.

$y = x^2$ is a \cup -shaped quadratic passing through $(0, 0)$.

$$(a) n(-3) = 5 - (-3) = 5 + 3 = 8$$

$$n(3) = 3^2 = 9$$

(b) There are two values of a .



The negative value of a is where

$$5 - a = 50$$

$$a = 5 - 50$$

$$a = -45$$

The positive value of a is where

$$a^2 = 50$$

$$a = \sqrt{50}$$

$$a = 5\sqrt{2}$$

The values of a such that $n(a) = 50$ are -45 and $+5\sqrt{2}$.

Solutionbank

Edexcel AS and A Level Modular Mathematics


Exercise F, Question 4
Question:

The function $g(x)$ is defined as $g(x) = 2x + 7 \quad \{ x \in \mathbb{R}, x \geq 0 \}$.

- (a) Sketch $g(x)$ and find the range.
- (b) Determine $g^{-1}(x)$, stating its domain.
- (c) Sketch $g^{-1}(x)$ on the same axes as $g(x)$, stating the relationship between the two graphs.


Solution:

- (a) $y = 2x + 7$ is a straight line of gradient 2 passing through 7 on the y axis.
The domain is given as $x \geq 0$.




Hence the range is $g(x) \geq 7$

- (b) The domain of the inverse function is $x \geq 7$.
To find the equation of the inverse function use a flow chart.



$$g^{-1}(x) = \frac{x-7}{2} \text{ and has domain } x \geq 7$$



$g^{-1}(x)$ is the reflection of $g(x)$ in the line $y = x$.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 5

Question:

The functions f and g are defined by

$$f : x \rightarrow 4x - 1 \quad \{ x \in \mathbb{R} \}$$

$$g : x \rightarrow \frac{3}{2x - 1} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \neq \frac{1}{2} \end{array} \right\}$$

Find in its simplest form:

- (a) the inverse function f^{-1}
- (b) the composite function gf , stating its domain
- (c) the values of x for which $2f(x) = g(x)$, giving your answers to 3 decimal places.

[E]

Solution:

(a) $f : x \rightarrow 4x - 1$

Let $y = 4x - 1$ and change the subject of the formula.

$$\Rightarrow y + 1 = 4x$$

$$\Rightarrow x = \frac{y + 1}{4}$$

Hence $f^{-1} : x \rightarrow \frac{x + 1}{4}$

(b) $gf(x) = g(4x - 1) = \frac{3}{2(4x - 1) - 1} = \frac{3}{8x - 3}$

Hence $gf : x \rightarrow \frac{3}{8x - 3}$

The domain would include all the real numbers apart from $x = \frac{3}{8}$ (i.e. where $8x - 3 = 0$).

(c) If $2f(x) = g(x)$

$$2 \times (4x - 1) = \frac{3}{2x - 1}$$

$$8x - 2 = \frac{3}{2x - 1}$$

$$(8x - 2)(2x - 1) = 3$$

$$16x^2 - 12x + 2 = 3$$

$$16x^2 - 12x - 1 = 0$$

Use $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ with $a = 16$, $b = -12$ and $c = -1$.

$$\text{Then } x = \frac{12 \pm \sqrt{144 + 64}}{32} = \frac{12 \pm \sqrt{208}}{32} = 0.826, -0.076$$

Values of x are -0.076 and 0.826

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 6

Question:

The function $f(x)$ is defined by

$$f(x) = \begin{cases} -x & x \leq 1 \\ x - 2 & x > 1 \end{cases}$$


(a) Sketch the graph of $f(x)$ for $-2 \leq x \leq 6$.

(b) Find the values of x for which $f(x) = -\frac{1}{2}$.

[E]

Solution:

(a)



For $x \leq 1$, $f(x) = -x$

This is a straight line of gradient -1 .

At point $x = 1$, its y coordinate is -1 .


For $x > 1$, $f(x) = x - 2$

This is a straight line of gradient $+1$.

At point $x = 1$, its y coordinate is also -1 .

The graph is said to be **continuous**.

(b) There are two values at which $f(x) = -\frac{1}{2}$ (see graph).



Point a is where

$$-x = -\frac{1}{2} \Rightarrow x = \frac{1}{2}$$

Point b is where

$$x - 2 = -\frac{1}{2} \Rightarrow x = 1\frac{1}{2}$$

The values of x for which $f(x) = -\frac{1}{2}$ are $\frac{1}{2}$ and $1\frac{1}{2}$.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 7

Question:

The function f is defined by

$$f : x \rightarrow \frac{2x+3}{x-1} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x > 1 \end{array} \right\}$$

- (a) Find $f^{-1}(x)$.
- (b) Find (i) the range of $f^{-1}(x)$
 (ii) the domain of $f^{-1}(x)$.

[E]

Solution:

(a) To find $f^{-1}(x)$ change the subject of the formula.

$$\text{Let } y = \frac{2x+3}{x-1}$$

$$y(x-1) = 2x+3$$

$$yx - y = 2x + 3$$

$$yx - 2x = y + 3$$


$$x(y-2) = y+3$$

$$x = \frac{y+3}{y-2}$$

$$\text{Therefore } f^{-1} : x \rightarrow \frac{x+3}{x-2}$$

(b) $f(x)$ has domain $\{x \in \mathbb{R}, \quad x > 1\}$ and range $\{f(x) \in \mathbb{R}, \quad f(x) > 2\}$

$$\text{As } x \rightarrow \infty, \quad y \rightarrow \frac{2x}{x} = 2$$



So $f^{-1}(x)$ has domain $\{x \in \mathbb{R}, x > 2\}$ and range $\{f^{-1}(x)$

$$\in \mathbb{R}, f^{-1} \left(x \right) > 1 \}$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 8

Question:

The functions f and g are defined by

$$f : x \rightarrow \frac{x}{x-2} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \neq 2 \end{array} \right\}$$

$$g : x \rightarrow \frac{3}{x} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \neq 0 \end{array} \right\}$$

- (a) Find an expression for $f^{-1}(x)$.
- (b) Write down the range of $f^{-1}(x)$.
- (c) Calculate $gf(1.5)$.
- (d) Use algebra to find the values of x for which $g(x) = f(x) + 4$.

[E]

Solution:

- (a) To find $f^{-1}(x)$ change the subject of the formula.

$$\text{Let } y = \frac{x}{x-2}$$

$$y(x-2) = x$$

$$yx - 2y = x \quad (\text{rearrange})$$

$$yx - x = 2y$$

$$x(y-1) = 2y$$

$$x = \frac{2y}{y-1}$$

It must always be rewritten as a function in x :

$$f^{-1} \left(x \right) = \frac{2x}{x-1}$$

- (b) The range of $f^{-1}(x)$ is the domain of $f(x)$.

Hence range is $\{ f^{-1}(x) \in \mathbb{R}, \quad f^{-1}(x) \neq 2 \}$.

$$(c) gf(1.5) = g \left(\frac{1.5}{1.5-2} \right) = g \left(\frac{1.5}{-0.5} \right) = g(-3) = \frac{3}{-3} = -1$$

(d) If $g(x) = f(x) + 4$

$$\frac{3}{x} = \frac{x}{x-2} + 4 \quad \left[\times x \left(x-2 \right) \right]$$

$$3(x-2) = x \times x + 4x(x-2)$$

$$3x - 6 = x^2 + 4x^2 - 8x$$

$$0 = 5x^2 - 11x + 6$$

$$0 = (5x-6)(x-1)$$

$$\Rightarrow x = \frac{6}{5}, 1$$

The values of x for which $g(x) = f(x) + 4$ are $\frac{6}{5}$ and 1.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise F, Question 9

Question:

The functions f and g are given by

$$f : x \rightarrow \frac{x}{x^2 - 1} - \frac{1}{x + 1} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x > 1 \end{array} \right\}$$

$$g : x \rightarrow \frac{2}{x} \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x > 0 \end{array} \right\}$$

(a) Show that $f(x) = \frac{1}{(x-1)(x+1)}$.

(b) Find the range of $f(x)$.

(c) Solve $gf(x) = 70$.

[E]


Solution:

$$\begin{aligned} (a) f(x) &= \frac{x}{x^2 - 1} - \frac{1}{x + 1} \\ &= \frac{x}{(x+1)(x-1)} - \frac{1}{(x+1)} \\ &= \frac{x}{(x+1)(x-1)} - \frac{x-1}{(x+1)(x-1)} \\ &= \frac{x - (x-1)}{(x+1)(x-1)} \\ &= \frac{1}{(x+1)(x-1)} \end{aligned}$$

(b) The range of $f(x)$ is the set of values that y take.

By using a graphical calculator we can see that $y = f \left(\begin{array}{c} x \end{array} \right) \quad \left\{ \begin{array}{l} x \in \mathbb{R}, \quad x \neq -1, \quad x \neq 1 \end{array} \right\}$

$\left. \begin{array}{l} x \in \mathbb{R}, \quad x \neq -1, \quad x \neq 1 \end{array} \right\}$ is a symmetrical graph about the y axis.



For $x > 1$, $f(x) > 0$

$$(c) gf(x) = g \left[\frac{1}{(x-1)(x+1)} \right] = \frac{2}{\frac{1}{(x-1)(x+1)}} = 2 \times \frac{(x-1)(x+1)}{1} = 2 \begin{pmatrix} x-1 \\ x+1 \end{pmatrix}$$

If $gf(x) = 70$

$$2(x-1)(x+1) = 70$$

$$(x-1)(x+1) = 35$$

$$x^2 - 1 = 35$$

$$x^2 = 36$$

$$x = \pm 6$$