

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 1


Question:

A curve is given by the parametric equations $x = 2t^2$, $y = 4t$. $t \in \mathbb{R}$. Copy and complete the following table and draw a graph of the curve for $-4 \leq t \leq 4$.

t	-4	-3	-2	-1	-0.5	0	0.5	1	2	3	4
$x = 2t^2$	32					0	0.5				32
$y = 4t$	-16						2				16

Solution:

t	-4	-3	-2	-1	-0.5	0	0.5	1	2	3	4
$x = 2t^2$	32	18	8	2	0.5	0	0.5	2	8	18	32
$y = 4t$	-16	-12	-8	-4	-2	0	2	4	8	12	16



Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 2


Question:

A curve is given by the parametric equations $x = 3t^2$, $y = 6t$. $t \in \mathbb{R}$. Copy and complete the following table and draw a graph of the curve for $-3 \leq t \leq 3$.

t	-3	-2	-1	-0.5	0	0.5	1	2	3
$x = 3t^2$					0				
$y = 6t$					0				

Solution:

t	-3	-2	-1	-0.5	0	0.5	1	2	3
$x = 3t^2$	27	12	3	0.75	0	0.75	3	12	27
$y = 6t$	-18	-12	-6	-3	0	3	6	12	18



Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 3


Question:

A curve is given by the parametric equations $x = 4t$, $y = \frac{4}{t}$. $t \in \mathbb{R}$, $t \neq 0$. Copy and complete the following table and draw a graph of the curve for $-4 \leq t \leq 4$.

t	-4	-3	-2	-1	-0.5	0.5	1	2	3	4
$x = 4t$	-16				-2					
$y = \frac{4}{t}$	-1				-8					

Solution:

t	-4	-3	-2	-1	-0.5	0.5	1	2	3	4
$x = 4t$	-16	-12	-8	-4	-2	2	4	8	12	16
$y = \frac{4}{t}$	-1	$-\frac{4}{3}$	-2	-4	-8	8	4	2	$\frac{4}{3}$	1



Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 4

Question:

Find the Cartesian equation of the curves given by these parametric equations.

a $x = 5t^2$, $y = 10t$

b $x = \frac{1}{2}t^2$, $y = t$

c $x = 50t^2$, $y = 100t$

d $x = \frac{1}{5}t^2$, $y = \frac{2}{5}t$

e $x = \frac{5}{2}t^2$, $y = 5t$

f $x = \sqrt{3}t^2$, $y = 2\sqrt{3}t$

g $x = 4t$, $y = 2t^2$

h $x = 6t$, $y = 3t^2$

Solution:

a $y = 10t$

So $t = \frac{y}{10}$ (1)

$$x = 5t^2 \quad (2)$$

Substitute (1) into (2):

$$x = 5\left(\frac{y}{10}\right)^2$$

So $x = \frac{5y^2}{100}$ simplifies to $x = \frac{y^2}{20}$

Hence, the Cartesian equation is $y^2 = 20x$.

b $y = t$ (1)

$$x = \frac{1}{2}t^2 \quad (2)$$

Substitute (1) into (2):

$$x = \frac{1}{2}y^2$$

Hence, the Cartesian equation is $y^2 = 2x$.

c $y = 100t$

So $t = \frac{y}{100}$ (1)

$$x = 50t^2 \quad (2)$$

Substitute (1) into (2):

$$x = 50\left(\frac{y}{100}\right)^2$$

So $x = \frac{50y^2}{10000}$ simplifies to $x = \frac{y^2}{200}$

Hence, the Cartesian equation is $y^2 = 200x$.

d $y = \frac{2}{5}t$

So $t = \frac{5y}{2}$ (1)

$$x = \frac{1}{5}t^2 \quad (2)$$

Substitute (1) into (2):

$$x = \frac{1}{5}\left(\frac{5y}{2}\right)^2$$

So $x = \frac{25y^2}{20}$ simplifies to $x = \frac{5y^2}{4}$

Hence, the Cartesian equation is $y^2 = \frac{4}{5}x$.

e $y = 5t$

So $t = \frac{y}{5}$ (1)

$$x = \frac{5}{2}t^2 \quad (2)$$

Substitute (1) into (2):

$$x = \frac{5}{2}\left(\frac{y}{5}\right)^2$$

So $x = \frac{5y^2}{50}$ simplifies to $x = \frac{y^2}{10}$

Hence, the Cartesian equation is $y^2 = 10x$.

f $y = 2\sqrt{3}t$

So $t = \frac{y}{2\sqrt{3}}$ (1)

$$x = \sqrt{3}t^2 \quad (2)$$

Substitute **(1)** into **(2)**:

$$x = \sqrt{3} \left(\frac{y}{2\sqrt{3}} \right)^2$$

So $x = \frac{\sqrt{3}y^2}{12}$ gives $y = \frac{12x}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$

Hence, the Cartesian equation is $y^2 = 4\sqrt{3}x$.

g $x = 4t$

So $t = \frac{x}{4}$ **(1)**

$$y = 2t^2 \quad \text{(2)}$$

Substitute **(1)** into **(2)**:

$$y = 2 \left(\frac{x}{4} \right)^2$$

So $y = \frac{2x^2}{16}$ simplifies to $y = \frac{x^2}{8}$

Hence, the Cartesian equation is $x^2 = 8y$.

h $x = 6t$

So $t = \frac{x}{6}$ **(1)**

$$y = 3t^2 \quad \text{(2)}$$

Substitute **(1)** into **(2)**:

$$y = 3 \left(\frac{x}{6} \right)^2$$

So $y = \frac{3x^2}{36}$ simplifies to $y = \frac{x^2}{12}$

Hence, the Cartesian equation is $x^2 = 12y$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 5

Question:

Find the Cartesian equation of the curves given by these parametric equations.

a $x = t, y = \frac{1}{t}, t \neq 0$

b $x = 7t, y = \frac{7}{t}, t \neq 0$

c $x = 3\sqrt{5}t, y = \frac{3\sqrt{5}}{t}, t \neq 0$

d $x = \frac{t}{5}, y = \frac{1}{5t}, t \neq 0$

Solution:

a $xy = t \times \left(\frac{1}{t}\right)$

$$xy = \frac{t}{t}$$

Hence, the Cartesian equation is $xy = 1$.

b $xy = 7t \times \left(\frac{7}{t}\right)$

$$xy = \frac{49t}{t}$$

Hence, the Cartesian equation is $xy = 49$.

c $xy = 3\sqrt{5}t \times \left(\frac{3\sqrt{5}}{t}\right)$

$$xy = \frac{9(5)t}{t}$$

Hence, the Cartesian equation is $xy = 45$.

d $xy = \frac{t}{5} \times \left(\frac{1}{5t}\right)$

$$xy = \frac{t}{25t}$$

Hence, the Cartesian equation is $xy = \frac{1}{25}$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 6

Question:

A curve has parametric equations $x = 3t$, $y = \frac{3}{t}$, $t \in \mathbb{R}$, $t \neq 0$.

a Find the Cartesian equation of the curve.

b Hence sketch this curve.


Solution:

a $xy = 3t \times \left(\frac{3}{t}\right)$

$$xy = \frac{9t}{t}$$

Hence, the Cartesian equation is $xy = 9$.

b



Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise A, Question 7

Question:

A curve has parametric equations $x = \sqrt{2}t$, $y = \frac{\sqrt{2}}{t}$, $t \in \mathbb{R}$, $t \neq 0$.

a Find the Cartesian equation of the curve.

b Hence sketch this curve.


Solution:

a $xy = \sqrt{2}t \times \left(\frac{\sqrt{2}}{t}\right)$

$$xy = \frac{2t}{t}$$

Hence, the Cartesian equation is $xy = 2$.

b



© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise B, Question 1

Question:

Find an equation of the parabola with

a focus $(5, 0)$ and directrix $x + 5 = 0$,

b focus $(8, 0)$ and directrix $x + 8 = 0$,

c focus $(1, 0)$ and directrix $x = -1$,

d focus $\left(\frac{3}{2}, 0\right)$ and directrix $x = -\frac{3}{2}$,

e focus $\left(\frac{\sqrt{3}}{2}, 0\right)$ and directrix $x + \frac{\sqrt{3}}{2} = 0$.

Solution:

The focus and directrix of a parabola with equation $y^2 = 4ax$, are $(a, 0)$ and $x + a = 0$ respectively.

a focus $(5, 0)$ and directrix $x + 5 = 0$.

So $a = 5$ and $y^2 = 4(5)x$.

Hence parabola has equation $y^2 = 20x$.

b focus $(8, 0)$ and directrix $x + 8 = 0$.

So $a = 8$ and $y^2 = 4(8)x$.

Hence parabola has equation $y^2 = 32x$.

c focus $(1, 0)$ and directrix $x = -1$ giving $x + 1 = 0$.

So $a = 1$ and $y^2 = 4(1)x$.

Hence parabola has equation $y^2 = 4x$.

d focus $\left(\frac{3}{2}, 0\right)$ and directrix $x = -\frac{3}{2}$ giving $x + \frac{3}{2} = 0$.

So $a = \frac{3}{2}$ and $y^2 = 4\left(\frac{3}{2}\right)x$.

Hence parabola has equation $y^2 = 6x$.

e focus $\left(\frac{\sqrt{3}}{2}, 0\right)$ and directrix $x + \frac{\sqrt{3}}{2} = 0$.

So $a = \frac{\sqrt{3}}{2}$ and $y^2 = 4\left(\frac{\sqrt{3}}{2}\right)x$.

Hence parabola has equation $y^2 = 2\sqrt{3}x$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise B, Question 2

Question:

Find the coordinates of the focus, and an equation for the directrix of a parabola with these equations.

a $y^2 = 12x$

b $y^2 = 20x$

c $y^2 = 10x$

d $y^2 = 4\sqrt{3}x$

e $y^2 = \sqrt{2}x$

f $y^2 = 5\sqrt{2}x$

Solution:

The focus and directrix of a parabola with equation $y^2 = 4ax$, are $(a, 0)$ and $x + a = 0$ respectively.

a $y^2 = 12x$. So $4a = 12$, gives $a = \frac{12}{4} = 3$.

So the focus has coordinates $(3, 0)$ and the directrix has equation $x + 3 = 0$.

b $y^2 = 20x$. So $4a = 20$, gives $a = \frac{20}{4} = 5$.

So the focus has coordinates $(5, 0)$ and the directrix has equation $x + 5 = 0$.

c $y^2 = 10x$. So $4a = 10$, gives $a = \frac{10}{4} = \frac{5}{2}$.

So the focus has coordinates $\left(\frac{5}{2}, 0\right)$ and the directrix has equation $x + \frac{5}{2} = 0$.

d $y^2 = 4\sqrt{3}x$. So $4a = 4\sqrt{3}$, gives $a = \frac{4\sqrt{3}}{4} = \sqrt{3}$.

So the focus has coordinates $(\sqrt{3}, 0)$ and the directrix has equation $x + \sqrt{3} = 0$.

e $y^2 = \sqrt{2}x$. So $4a = \sqrt{2}$, gives $a = \frac{\sqrt{2}}{4}$.

So the focus has coordinates $\left(\frac{\sqrt{2}}{4}, 0\right)$ and the directrix has equation $x + \frac{\sqrt{2}}{4} = 0$.

f $y^2 = 5\sqrt{2}x$. So $4a = 5\sqrt{2}$, gives $a = \frac{5\sqrt{2}}{4}$.

So the focus has coordinates $\left(\frac{5\sqrt{2}}{4}, 0\right)$ and the directrix has equation $x + \frac{5\sqrt{2}}{4} = 0$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics


Quadratic Equations

Exercise B, Question 3

Question:

A point $P(x, y)$ obeys a rule such that the distance of P to the point $(3, 0)$ is the same as the distance of P to the straight line $x + 3 = 0$. Prove that the locus of P has an equation of the form $y^2 = 4ax$, stating the value of the constant a .

Solution:



From sketch the locus satisfies $SP = XP$.

Therefore, $SP^2 = XP^2$.

So, $(x - 3)^2 + (y - 0)^2 = (x - -3)^2$.

$$x^2 - 6x + 9 + y^2 = x^2 + 6x + 9$$

$$-6x + y^2 = 6x$$

which simplifies to $y^2 = 12x$.

So, the locus of P has an equation of the form $y^2 = 4ax$, where $a = 3$.

The (shortest) distance of P to the line $x + 3 = 0$ is the distance XP .

The distance SP is the same as the distance XP .

The line XP is horizontal and has distance $XP = x + 3$.

The locus of P is the curve shown.

This means the distance SP is the same as the distance XP .

Use $d^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$ on $SP^2 = XP^2$, where $S(3, 0)$, $P(x, y)$, and $X(-3, y)$.

This is in the form $y^2 = 4ax$.

So $4a = 12$, gives $a = \frac{12}{4} = 3$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics


Quadratic Equations

Exercise B, Question 4

Question:

A point $P(x, y)$ obeys a rule such that the distance of P to the point $(2\sqrt{5}, 0)$ is the same as the distance of P to the straight line $x = -2\sqrt{5}$. Prove that the locus of P has an equation of the form $y^2 = 4ax$, stating the value of the constant a .

Solution:



From sketch the locus satisfies $SP = XP$.

Therefore, $SP^2 = XP^2$.

So, $(x - 2\sqrt{5})^2 + (y - 0)^2 = (x - (-2\sqrt{5}))^2$.

$$x^2 - 4\sqrt{5}x + 20 + y^2 = x^2 + 4\sqrt{5}x + 20$$

$$-4\sqrt{5}x + y^2 = 4\sqrt{5}x$$

which simplifies to $y^2 = 8\sqrt{5}x$.

So, the locus of P has an equation of the form $y^2 = 4ax$, where $a = 2\sqrt{5}$.

The (shortest) distance of P to the line $x = -2\sqrt{5}$ or $x + 2\sqrt{5} = 0$ is the distance XP .

The distance SP is the same as the distance XP .

The line XP is horizontal and has distance $XP = x + 2\sqrt{5}$.

The locus of P is the curve shown.

This means the distance SP is the same as the distance XP .

Use $d^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$ on $SP^2 = XP^2$, where $S(2\sqrt{5}, 0)$, $P(x, y)$, and $X(-2\sqrt{5}, y)$.

This is in the form $y^2 = 4ax$.

So $4a = 8\sqrt{5}$, gives $a = \frac{8\sqrt{5}}{4} = 2\sqrt{5}$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise B, Question 5

Question:

A point $P(x, y)$ obeys a rule such that the distance of P to the point $(0, 2)$ is the same as the distance of P to the straight line $y = -2$.

a Prove that the locus of P has an equation of the form $y = kx^2$, stating the value of the constant k .


Given that the locus of P is a parabola,

b state the coordinates of the focus of P , and an equation of the directrix to P ,

c sketch the locus of P with its focus and its directrix.

Solution:

a



The (shortest) distance of P to the line $y = -2$ is the distance YP .

The distance SP is the same as the distance YP .

The line YP is vertical and has distance $YP = y + 2$.

The locus of P is the curve shown.

From sketch the locus satisfies $SP = YP$.

Therefore, $SP^2 = YP^2$.

So, $(x - 0)^2 + (y - 2)^2 = (y - (-2))^2$.

$$x^2 + y^2 - 4y + 4 = y^2 + 4y + 4$$

$$x^2 - 4y = 4y$$

which simplifies to $x^2 = 8y$ and then $y = \frac{1}{8}x^2$.

So, the locus of P has an equation of the form $y = \frac{1}{8}x^2$, where

This means the distance SP is the same as the distance YP .

Use $d^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$ on $SP^2 = YP^2$, where $S(0, 2)$, $P(x, y)$, and $Y(x, -2)$.


$$k = \frac{1}{8}.$$

b The focus and directrix of a parabola with equation $y^2 = 4ax$, are $(a, 0)$ and $x + a = 0$ respectively. Therefore it follows that the focus and directrix of a parabola with equation $x^2 = 4ay$, are $(0, a)$ and $y + a = 0$ respectively.

So the focus has coordinates $(0, 2)$ and the directrix has equation $y + 2 = 0$.

$x^2 = 8y$ is in the form $x^2 = 4ay$.
So $4a = 8$, gives $a = \frac{8}{4} = 2$.

c



© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 1

Question:

The line $y = 2x - 3$ meets the parabola $y^2 = 3x$ at the points P and Q .

Find the coordinates of P and Q .

Solution:

Line: $y = 2x - 3$ (1)

Curve: $y^2 = 3x$ (2)

Substituting (1) into (2) gives

$$\begin{aligned}(2x - 3)^2 &= 3x \\(2x - 3)(2x - 3) &= 3x \\4x^2 - 12x + 9 &= 3x \\4x^2 - 15x + 9 &= 0 \\(x - 3)(4x - 3) &= 0 \\x &= 3, \frac{3}{4}\end{aligned}$$

When $x = 3$, $y = 2(3) - 3 = 3$

When $x = \frac{3}{4}$, $y = 2\left(\frac{3}{4}\right) - 3 = -\frac{3}{2}$

Hence the coordinates of P and Q are $(3, 3)$ and $\left(\frac{3}{4}, -\frac{3}{2}\right)$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 2

Question:

The line $y = x + 6$ meets the parabola $y^2 = 32x$ at the points A and B . Find the exact length AB giving your answer as a surd in its simplest form.

Solution:

Line: $y = x + 6$ (1)

Curve: $y^2 = 32x$ (2)

Substituting (1) into (2) gives

$$(x + 6)^2 = 32x$$

$$(x + 6)(x + 6) = 32x$$

$$x^2 + 12x + 36 = 32x$$

$$x^2 - 20x + 36 = 0$$

$$(x - 2)(x - 18) = 0$$

$$x = 2, 18$$

When $x = 2$, $y = 2 + 6 = 8$.

When $x = 18$, $y = 18 + 6 = 24$.

Hence the coordinates of A and B are $(2, 8)$ and $(18, 24)$.

$$\begin{aligned} AB &= \sqrt{(18 - 2)^2 + (24 - 8)^2} \quad \text{Use } d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}. \\ &= \sqrt{16^2 + 16^2} \\ &= \sqrt{2(16)^2} \\ &= 16\sqrt{2} \end{aligned}$$

Hence the exact length AB is $16\sqrt{2}$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 3

Question:

The line $y = x - 20$ meets the parabola $y^2 = 10x$ at the points A and B . Find the coordinates of A and B . The mid-point of AB is the point M . Find the coordinates of M .

Solution:

Line: $y = x - 20$ (1)

Curve: $y^2 = 10x$ (2)

Substituting (1) into (2) gives

$$(x - 20)^2 = 10x$$

$$(x - 20)(x - 20) = 10x$$

$$x^2 - 40x + 400 = 10x$$

$$x^2 - 50x + 400 = 0$$

$$(x - 10)(x - 40) = 0$$

$$x = 10, 40$$

When $x = 10$, $y = 10 - 20 = -10$.

When $x = 40$, $y = 40 - 20 = 20$.

Hence the coordinates of A and B are $(10, -10)$ and $(40, 20)$.

The midpoint of A and B is $\left(\frac{10+40}{2}, \frac{-10+20}{2}\right) = (25, 5)$. Use $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$

Hence the coordinates of M are $(25, 5)$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 4

Question:

The parabola C has parametric equations $x = 6t^2$, $y = 12t$. The focus to C is at the point S .

- a** Find a Cartesian equation of C .
- b** State the coordinates of S and the equation of the directrix to C .
- c** Sketch the graph of C .

The points P and Q are both at a distance 9 units away from the directrix of the parabola.

- d** State the distance PS .
- e** Find the exact length PQ , giving your answer as a surd in its simplest form.
- f** Find the area of the triangle PQS , giving your answer in the form $k\sqrt{2}$, where k is an integer.

Solution:

a $y = 12t$

So $t = \frac{y}{12}$ (1)

$x = 6t^2$ (2)

Substitute (1) into (2):

$$x = 6\left(\frac{y}{12}\right)^2$$


So $x = \frac{6y^2}{144}$ simplifies to $x = \frac{y^2}{24}$

Hence, the Cartesian equation is $y^2 = 24x$.


b $y^2 = 24x$. So $4a = 24$, gives $a = \frac{24}{4} = 6$.

So the focus S , has coordinates $(6, 0)$ and the directrix has equation $x + 6 = 0$.

c



d



The (shortest) distance of P to the line $x + 6 = 0$ is the distance X_1P .

Therefore $X_1P = 9$.

The distance PS is the same as the distance X_1P , by the focus-directrix property.

Hence the distance $PS = 9$.

e Using diagram in (d), the x -coordinate of P and Q is $x = 9 - 6 = 3$.

When $x = 3$, $y^2 = 24(3) = 72$.

$$\begin{aligned} \text{Hence } y &= \pm\sqrt{72} \\ &= \pm\sqrt{36}\sqrt{2} \\ &= \pm 6\sqrt{2} \end{aligned}$$

So the coordinates of P and Q are $(3, 6\sqrt{2})$ and $(3, -6\sqrt{2})$.


As P and Q are vertically above each other then

$$\begin{aligned} PQ &= 6\sqrt{2} - (-6\sqrt{2}) \\ &= 12\sqrt{2}. \end{aligned}$$

Hence, the distance PQ is $12\sqrt{2}$.

f Drawing a diagram of the triangle PQS gives:

The x -coordinate of P and Q is 3 and the x -coordinate of S is 6.



Hence the height of the triangle is height
 $= 6 - 3 = 3$.

The length of the base is $12\sqrt{2}$.

$$\begin{aligned} \text{Area} &= \frac{1}{2}(12\sqrt{2})(3) \\ &= \frac{1}{2}(36\sqrt{2}) \\ &= 18\sqrt{2}. \end{aligned}$$

Therefore the area of the triangle is $18\sqrt{2}$, where $k = 18$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 5

Question:

The parabola C has equation $y^2 = 4ax$, where a is a constant. The point $\left(\frac{5}{4}t^2, \frac{5}{2}t\right)$ is a general point on C .

a Find a Cartesian equation of C .

The point P lies on C with y -coordinate 5.

b Find the x -coordinate of P .

The point Q lies on the directrix of C where $y = 3$. The line l passes through the points P and Q .

c Find the coordinates of Q .

d Find an equation for l , giving your answer in the form $ax + by + c = 0$, where a , b and c are integers.

Solution:

a $P\left(\frac{5}{4}t^2, \frac{5}{2}t\right)$. Substituting $x = \frac{5}{4}t^2$ and $y = \frac{5}{2}t$ into $y^2 = 4ax$ gives,

$$\left(\frac{5}{2}t\right)^2 = 4a\left(\frac{5}{4}t^2\right) \Rightarrow \frac{25t^2}{4} = 5at^2 \Rightarrow \frac{25}{4} = 5a \Rightarrow \frac{5}{4} = a$$

$$\text{When } a = \frac{5}{4}, y^2 = 4\left(\frac{5}{4}\right)x \Rightarrow y^2 = 5x$$

The Cartesian equation of C is $y^2 = 5x$.

b When $y = 5$, $(5)^2 = 5x \Rightarrow \frac{25}{5} = x \Rightarrow x = 5$.

The x -coordinate of P is 5.

c As $a = \frac{5}{4}$, the equation of the directrix of C is $x + \frac{5}{4} = 0$ or $x = -\frac{5}{4}$.

Therefore the coordinates of Q are $\left(-\frac{5}{4}, 3\right)$.

d The coordinates of P and Q are $(5, 5)$ and $\left(-\frac{5}{4}, 3\right)$.

$$m_l = m_{PQ} = \frac{3-5}{-\frac{5}{4}-5} = \frac{-2}{-\frac{25}{4}} = \frac{8}{25}$$

$$l: y - 5 = \frac{8}{25}(x - 5)$$

$$l: 25y - 125 = 8(x - 5)$$

$$l: 25y - 125 = 8x - 40$$

$$l: 0 = 8x - 25y - 40 + 125$$

$$l: 0 = 8x - 25y + 85$$

An equation for l is $8x - 25y + 85 = 0$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 6

Question:

A parabola C has equation $y^2 = 4x$. The point S is the focus to C .

a Find the coordinates of S .

The point P with y -coordinate 4 lies on C .

b Find the x -coordinate of P .

The line l passes through S and P .

c Find an equation for l , giving your answer in the form $ax + by + c = 0$, where a , b and c are integers.

The line l meets C again at the point Q .

d Find the coordinates of Q .

e Find the distance of the directrix of C to the point Q .

Solution:

a $y^2 = 4x$. So $4a = 4$, gives $a = \frac{4}{4} = 1$.

So the focus S , has coordinates $(1, 0)$.

Also note that the directrix has equation $x + 1 = 0$.

b Substituting $y = 4$ into $y^2 = 4x$ gives:

$$16 = 4x \Rightarrow x = \frac{16}{4} = 4.$$

The x -coordinate of P is 4.

c The line l goes through $S(1, 0)$ and $P(4, 4)$.

$$\text{Hence gradient of } l, m_l = \frac{4-0}{4-1} = \frac{4}{3}$$

$$\text{Hence, } y - 0 = \frac{4}{3}(x - 1)$$

$$3y = 4(x - 1)$$

$$3y = 4x - 4$$

$$0 = 4x - 3y - 4$$

The line l has equation $4x - 3y - 4 = 0$.

$$\text{d Line } l : 4x - 3y - 4 = 0 \quad \text{(1)}$$

$$\text{Curve : } y^2 = 4x \quad \text{(2)}$$

Substituting (2) into (1) gives

$$\begin{aligned}y^2 - 3y - 4 &= 0 \\(y - 4)(y + 1) &= 0 \\y &= 4, -1\end{aligned}$$


At P , it is already known that $y = 4$. So at Q , $y = -1$.

Substituting $y = -1$ into $y^2 = 4x$ gives

$$(-1)^2 = 4x \Rightarrow x = \frac{1}{4}.$$

Hence the coordinates of Q are $\left(\frac{1}{4}, -1\right)$.

e The directrix of C has equation $x + 1 = 0$ or $x = -1$. Q has coordinates $\left(\frac{1}{4}, -1\right)$.



From the diagram, distance $= 1 + \frac{1}{4} = \frac{5}{4}$.

Therefore the distance of the directrix of C to the point Q is $\frac{5}{4}$.

Solutionbank FP1


Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 7

Question:

The diagram shows the point P which lies on the parabola C with equation $y^2 = 12x$.



The point S is the focus of C . The points Q and R lie on the directrix to C . The line segment QP is parallel to the line segment RS as shown in the diagram. The distance of PS is 12 units.

a Find the coordinates of R and S .

b Hence find the exact coordinates of P and Q .

c Find the area of the quadrilateral $PQRS$, giving your answer in the form $k\sqrt{3}$, where k is an integer.

Solution:

a $y^2 = 12x$. So $4a = 12$, gives $a = \frac{12}{4} = 3$.

Therefore the focus S has coordinates $(3, 0)$ and an equation of the directrix of C is $x + 3 = 0$ or $x = -3$. The coordinates of R are $(-3, 0)$ as R lies on the x -axis.

b The directrix has equation $x = -3$. The (shortest) distance of P to the directrix is the distance PQ . The distance $SP = 12$. The focus-directrix property implies that $SP = PQ = 12$.


Therefore the x -coordinate of P is $x = 12 - 3 = 9$.

As P lies on C , when $x = 9$, $y^2 = 12(9) \Rightarrow y^2 = 108$

As $y > 0$, $y = \sqrt{108} = \sqrt{36} \sqrt{3} = 6\sqrt{3} \Rightarrow P(9, 6\sqrt{3})$

Hence the exact coordinates of P are $(9, 6\sqrt{3})$ and the coordinates of Q are $(-3, 6\sqrt{3})$.

c



$$\begin{aligned}\text{Area}(PQRS) &= \frac{1}{2}(6 + 12)6\sqrt{3} \\ &= \frac{1}{2}(18)(6\sqrt{3}) \\ &= (9)(6\sqrt{3}) \\ &= 54\sqrt{3}\end{aligned}$$

The area of the quadrilateral $PQRS$ is $54\sqrt{3}$ and $k = 54$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise C, Question 8

Question:

The points $P(16, 8)$ and $Q(4, b)$, where $b < 0$ lie on the parabola C with equation $y^2 = 4ax$.

a Find the values of a and b .

P and Q also lie on the line l . The mid-point of PQ is the point R .

b Find an equation of l , giving your answer in the form $y = mx + c$, where m and c are constants to be determined.

c Find the coordinates of R .

The line n is perpendicular to l and passes through R .

d Find an equation of n , giving your answer in the form $y = mx + c$, where m and c are constants to be determined.

The line n meets the parabola C at two points.

e Show that the x -coordinates of these two points can be written in the form $x = \lambda \pm \mu\sqrt{13}$, where λ and μ are integers to be determined.

Solution:

a $P(16, 8)$. Substituting $x = 16$ and $y = 8$ into $y^2 = 4ax$ gives,

$$(8)^2 = 4a(16) \Rightarrow 64 = 64a \Rightarrow a = \frac{64}{64} = 1.$$

$Q(4, b)$. Substituting $x = 4$, $y = b$ and $a = 1$ into $y^2 = 4ax$ gives,

$$b^2 = 4(1)(4) = 16 \Rightarrow b = \pm\sqrt{16} \Rightarrow b = \pm 4. \text{ As } b < 0, b = -4.$$

Hence, $a = 1$, $b = -4$.

b The coordinates of P and Q are $(16, 8)$ and $(4, -4)$.

$$m_l = m_{PQ} = \frac{-4 - 8}{4 - 16} = \frac{-12}{-12} = 1$$

$$l : y - 8 = 1(x - 16)$$

$$l : y = x - 8$$

l has equation $y = x - 8$.

c R has coordinates $\left(\frac{16+4}{2}, \frac{8+(-4)}{2}\right) = (10, 2)$.

d As n is perpendicular to l , $m_n = -1$

$$n : y - 2 = -1(x - 10)$$

$$n : y - 2 = -x + 10$$

$$n : y = -x + 12$$

n has equation $y = -x + 12$.

e Line n : $y = -x + 12$ **(1)**

Parabola C : $y^2 = 4x$ **(2)**

Substituting **(1)** into **(2)** gives

$$(-x + 12)^2 = 4x$$

$$x^2 - 12x - 12x + 144 = 4x$$

$$x^2 - 28x + 144 = 0$$

$$(x - 14)^2 - 196 + 144 = 0$$

$$(x - 14)^2 - 52 = 0$$

$$(x - 14)^2 = 52$$

$$x - 14 = \pm\sqrt{52}$$

$$x - 14 = \pm\sqrt{4}\sqrt{13}$$

$$x - 14 = \pm 2\sqrt{13}$$

$$x = 14 \pm 2\sqrt{13}$$

The x coordinates are $x = 14 \pm 2\sqrt{13}$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 1

Question:

Find the equation of the tangent to the curve

a $y^2 = 4x$ at the point (16, 8)

b $y^2 = 8x$ at the point (4, $4\sqrt{2}$)

c $xy = 25$ at the point (5, 5)

d $xy = 4$ at the point where $x = \frac{1}{2}$

e $y^2 = 7x$ at the point (7, -7)

f $xy = 16$ at the point where $x = 2\sqrt{2}$.

Give your answers in the form $ax + by + c = 0$.

Solution:

a As $y > 0$ in the coordinates (16, 8), then

$$y^2 = 4x \Rightarrow y = \sqrt{4x} = \sqrt{4} \sqrt{x} = 2x^{\frac{1}{2}}$$

$$\text{So } y = 2x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{1}{\sqrt{x}}$$

$$\text{At (16, 8), } m_T = \frac{dy}{dx} = \frac{1}{\sqrt{16}} = \frac{1}{4}.$$

$$\text{T: } y - 8 = \frac{1}{4}(x - 16)$$

$$\text{T: } 4y - 32 = x - 16$$

$$\text{T: } 0 = x - 4y - 16 + 32$$

$$\text{T: } x - 4y + 16 = 0$$

Therefore, the equation of the tangent is $x - 4y + 16 = 0$.

b As $y > 0$ in the coordinates (4, $4\sqrt{2}$), then

$$y^2 = 8x \Rightarrow y = \sqrt{8x} = \sqrt{8} \sqrt{x} = \sqrt{4} \sqrt{2} \sqrt{x} = 2\sqrt{2}x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{2}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{2}\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \sqrt{2}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{2}}{\sqrt{x}}$$

$$\text{At } (4, 4\sqrt{2}), m_T = \frac{dy}{dx} = \frac{\sqrt{2}}{\sqrt{4}} = \frac{\sqrt{2}}{2}.$$

$$\text{T: } y - 4\sqrt{2} = \frac{\sqrt{2}}{2}(x - 4)$$

$$\text{T: } 2y - 8\sqrt{2} = \sqrt{2}(x - 4)$$

$$\text{T: } 2y - 8\sqrt{2} = \sqrt{2}x - 4\sqrt{2}$$

$$\text{T: } 0 = \sqrt{2}x - 2y - 4\sqrt{2} + 8\sqrt{2}$$

$$\text{T: } \sqrt{2}x - 2y + 4\sqrt{2} = 0$$

Therefore, the equation of the tangent is $\sqrt{2}x - 2y + 4\sqrt{2} = 0$.

$$\text{c } xy = 25 \Rightarrow y = 25x^{-1}$$

$$\frac{dy}{dx} = -25x^{-2} = -\frac{25}{x^2}$$

$$\text{At } (5, 5), m_T = \frac{dy}{dx} = -\frac{25}{5^2} = -\frac{25}{25} = -1$$

$$\text{T: } y - 5 = -1(x - 5)$$

$$\text{T: } y - 5 = -x + 5$$

$$\text{T: } x + y - 5 - 5 = 0$$

$$\text{T: } x + y - 10 = 0$$

Therefore, the equation of the tangent is $x + y - 10 = 0$.

$$\text{d } xy = 4 \Rightarrow y = 4x^{-1}$$

$$\frac{dy}{dx} = -4x^{-2} = -\frac{4}{x^2}$$

$$\text{At } x = \frac{1}{2}, m_T = \frac{dy}{dx} = -\frac{4}{\left(\frac{1}{2}\right)^2} = -\frac{4}{\left(\frac{1}{4}\right)} = -16$$

$$\text{When } x = \frac{1}{2}, y = \frac{4}{\left(\frac{1}{2}\right)} = 8 \Rightarrow \left(\frac{1}{2}, 8\right)$$

$$\text{T: } y - 8 = -16\left(x - \frac{1}{2}\right)$$

$$\text{T: } y - 8 = -16x + 8$$

$$\mathbf{T: } 16x + y - 8 - 8 = 0$$

$$\mathbf{T: } 16x + y - 16 = 0$$

Therefore, the equation of the tangent is $16x + y - 16 = 0$.

e As $y < 0$ in the coordinates $(7, -7)$, then

$$y^2 = 7x \Rightarrow y = -\sqrt{7x} = -\sqrt{7} \sqrt{x} = -\sqrt{7} x^{\frac{1}{2}}$$

$$\text{So } y = -\sqrt{7} x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = -\sqrt{7} \left(\frac{1}{2}\right) x^{-\frac{1}{2}} = -\frac{\sqrt{7}}{2} x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = -\frac{\sqrt{7}}{2\sqrt{x}}$$

$$\text{At } (7, -7), m_T = \frac{dy}{dx} = -\frac{\sqrt{7}}{2\sqrt{7}} = -\frac{1}{2}.$$

$$\mathbf{T: } y + 7 = -\frac{1}{2}(x - 7)$$

$$\mathbf{T: } 2y + 14 = -1(x - 7)$$

$$\mathbf{T: } 2y + 14 = -x + 7$$

$$\mathbf{T: } x + 2y + 14 - 7 = 0$$

$$\mathbf{T: } x + 2y + 7 = 0$$

Therefore, the equation of the tangent is $x + 2y + 7 = 0$.

$$\mathbf{f} \text{ } xy = 16 \Rightarrow y = 16x^{-1}$$

$$\frac{dy}{dx} = -16x^{-2} = -\frac{16}{x^2}$$

$$\text{At } x = 2\sqrt{2}, m_T = \frac{dy}{dx} = -\frac{16}{(2\sqrt{2})^2} = -\frac{16}{8} = -2$$

$$\text{When } x = 2\sqrt{2}, y = \frac{16}{2\sqrt{2}} = \frac{8}{\sqrt{2}} = \frac{8\sqrt{2}}{\sqrt{2}\sqrt{2}} = 4\sqrt{2} \Rightarrow (2\sqrt{2}, 4\sqrt{2})$$

$$\mathbf{T: } y - 4\sqrt{2} = -2(x - 2\sqrt{2})$$

$$\mathbf{T: } y - 4\sqrt{2} = -2x + 4\sqrt{2}$$

$$\mathbf{T: } 2x + y - 4\sqrt{2} - 4\sqrt{2} = 0$$

$$\mathbf{T: } 2x + y - 8\sqrt{2} = 0$$

Therefore, the equation of the tangent is $2x + y - 8\sqrt{2} = 0$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 2

Question:

Find the equation of the normal to the curve

a $y^2 = 20x$ at the point where $y = 10$,

b $xy = 9$ at the point $\left(-\frac{3}{2}, -6\right)$.

Give your answers in the form $ax + by + c = 0$, where a , b and c are integers.

Solution:

a Substituting $y = 10$ into $y^2 = 20x$ gives

$$(10)^2 = 20x \Rightarrow x = \frac{100}{20} = 5 \Rightarrow (5, 10)$$

As $y > 0$, then

$$y^2 = 20x \Rightarrow y = \sqrt{20x} = \sqrt{20} \sqrt{x} = \sqrt{4} \sqrt{5} \sqrt{x} = 2\sqrt{5} x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{5} x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{5} \left(\frac{1}{2}\right) x^{-\frac{1}{2}} = \sqrt{5} x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{5}}{\sqrt{x}}$$

$$\text{At } (5, 10), m_T = \frac{dy}{dx} = \frac{\sqrt{5}}{\sqrt{5}} = 1.$$

Gradient of tangent at $(5, 10)$ is $m_T = 1$.

So gradient of normal is $m_N = -1$.

$$\text{N: } y - 10 = -1(x - 5)$$

$$\text{N: } y - 10 = -x + 5$$

$$\text{N: } x + y - 10 - 5 = 0$$

$$\text{N: } x + y - 15 = 0$$

Therefore, the equation of the normal is $x + y - 15 = 0$.

b $xy = 9 \Rightarrow y = 9x^{-1}$

$$\frac{dy}{dx} = -9x^{-2} = -\frac{9}{x^2}$$

$$\text{At } x = -\frac{3}{2}, m_T = \frac{dy}{dx} = -\frac{9}{\left(-\frac{3}{2}\right)^2} = -\frac{9}{\left(\frac{9}{4}\right)} = -\frac{36}{9} = -4$$

Gradient of tangent at $\left(-\frac{3}{2}, -6\right)$ is $m_T = -4$.

So gradient of normal is $m_N = \frac{-1}{-4} = \frac{1}{4}$.

$$\mathbf{N: } y + 6 = \frac{1}{4}\left(x + \frac{3}{2}\right)$$

$$\mathbf{N: } 4y + 24 = x + \frac{3}{2}$$

$$\mathbf{N: } 8y + 48 = 2x + 3$$

$$\mathbf{N: } 0 = 2x - 8y + 3 - 48$$

$$\mathbf{N: } 0 = 2x - 8y - 45$$

Therefore, the equation of the normal is $2x - 8y - 45 = 0$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 3

Question:

The point $P(4, 8)$ lies on the parabola with equation $y^2 = 4ax$. Find

- a** the value of a ,
- b** an equation of the normal to C at P .

The normal to C at P cuts the parabola again at the point Q . Find

- c** the coordinates of Q ,
- d** the length PQ , giving your answer as a simplified surd.

Solution:

a Substituting $x = 4$ and $y = 8$ into $y^2 = 4ax$ gives

$$(8)^2 = 4(a)(4) \Rightarrow 64 = 16a \Rightarrow a = \frac{64}{16} = 4$$

So, $a = 4$.

b When $a = 4$, $y^2 = 4(4)x \Rightarrow y^2 = 16x$.

For $P(4, 8)$, $y > 0$, so

$$y^2 = 16x \Rightarrow y = \sqrt{16x} = \sqrt{16} \sqrt{x} = 4\sqrt{x} = 4x^{\frac{1}{2}}$$

$$\text{So } y = 4x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 4\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = 2x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{2}{\sqrt{x}}$$

$$\text{At } P(4, 8), m_T = \frac{dy}{dx} = \frac{2}{\sqrt{4}} = \frac{2}{2} = 1.$$

Gradient of tangent at $P(4, 8)$ is $m_T = 1$.

So gradient of normal at $P(4, 8)$ is $m_N = -1$.

$$\text{N: } y - 8 = -1(x - 4)$$

$$\text{N: } y - 8 = -x + 4$$

$$\text{N: } y = -x + 4 + 8$$

$$\text{N: } y = -x + 12$$

Therefore, the equation of the normal to C at P is $y = -x + 12$.

c Normal N: $y = -x + 12$ (1)

Parabola: $y^2 = 16x$ (2)

Multiplying (1) by 16 gives

$$16y = -16x + 192$$

Substituting (2) into this equation gives

$$\begin{aligned} 16y &= -y^2 + 192 \\ y^2 + 16y - 192 &= 0 \\ (y + 24)(y - 8) &= 0 \\ y &= -24, 8 \end{aligned}$$

At P , it is already known that $y = 8$. So at Q , $y = -24$.

Substituting $y = -24$ into $y^2 = 16x$ gives

$$(-24)^2 = 16x \Rightarrow 576 = 16x \Rightarrow x = \frac{576}{16} = 36.$$

Hence the coordinates of Q are $(36, -24)$.

d The coordinates of P and Q are $(4, 8)$ and $(36, -24)$.

$$\begin{aligned} AB &= \sqrt{(36 - 4)^2 + (-24 - 8)^2} \quad \text{Use } d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}. \\ &= \sqrt{32^2 + (-32)^2} \\ &= \sqrt{2(32)^2} \\ &= \sqrt{2} \sqrt{(32)^2} \\ &= 32\sqrt{2} \end{aligned}$$

Hence the exact length AB is $32\sqrt{2}$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 4

Question:

The point $A(-2, -16)$ lies on the rectangular hyperbola H with equation $xy = 32$.

a Find an equation of the normal to H at A .

The normal to H at A meets H again at the point B .

b Find the coordinates of B .

Solution:

$$\mathbf{a} \quad xy = 32 \Rightarrow y = 32x^{-1}$$

$$\frac{dy}{dx} = -32x^{-2} = -\frac{32}{x^2}$$

$$\text{At } A(-2, -16), m_T = \frac{dy}{dx} = -\frac{32}{2^2} = -\frac{32}{4} = -8$$

Gradient of tangent at $A(-2, -16)$ is $m_T = -8$.

So gradient of normal at $A(-2, -16)$ is $m_N = \frac{-1}{-8} = \frac{1}{8}$.

$$\mathbf{N}: y + 16 = \frac{1}{8}(x + 2)$$

$$\mathbf{N}: 8y + 128 = x + 2$$

$$\mathbf{N}: 0 = x - 8y + 2 - 128$$

$$\mathbf{N}: 0 = x - 8y - 126$$

The equation of the normal to H at A is $x - 8y - 126 = 0$.

$$\mathbf{b} \text{ Normal } \mathbf{N}: x - 8y - 126 = 0 \quad (1)$$

$$\text{Hyperbola } H: xy = 32 \quad (2)$$

Rearranging (2) gives

$$y = \frac{32}{x}$$

Substituting this equation into (1) gives

$$x - 8\left(\frac{32}{x}\right) - 126 = 0$$

$$x - \left(\frac{256}{x}\right) - 126 = 0$$

Multiplying both sides by x gives

$$x^2 - 256 - 126x = 0$$

$$x^2 - 126x - 256 = 0$$

$$(x - 128)(x + 2) = 0$$

$$x = 128, -2$$

At A , it is already known that $x = -2$. So at B , $x = 128$.

Substituting $x = 128$ into $y = \frac{32}{x}$ gives

$$y = \frac{32}{128} = \frac{1}{4}$$

Hence the coordinates of B are $\left(128, \frac{1}{4}\right)$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 5

Question:

The points $P(4, 12)$ and $Q(-8, -6)$ lie on the rectangular hyperbola H with equation $xy = 48$.

a Show that an equation of the line PQ is $3x - 2y + 12 = 0$.

The point A lies on H . The normal to H at A is parallel to the chord PQ .

b Find the exact coordinates of the two possible positions of A .

Solution:

a The points P and Q have coordinates $P(4, 12)$ and $Q(-8, -6)$.

$$\text{Hence gradient of } PQ, m_{PQ} = \frac{-6 - 12}{-8 - 4} = \frac{-18}{-12} = \frac{3}{2}$$

$$\text{Hence, } y - 12 = \frac{3}{2}(x - 4)$$

$$2y - 24 = 3(x - 4)$$

$$2y - 24 = 3x - 12$$

$$0 = 3x - 2y - 12 + 24$$

$$0 = 3x - 2y + 12$$

The line PQ has equation $3x - 2y + 12 = 0$.

b From part (a), the gradient of the chord PQ is $\frac{3}{2}$.

The normal to H at A is parallel to the chord PQ , implies that the gradient of the normal to H at A is $\frac{3}{2}$.

It follows that the gradient of the tangent to H at A is

$$m_T = \frac{-1}{m_N} = \frac{-1}{\left(\frac{3}{2}\right)} = -\frac{2}{3}$$

$$H : xy = 48 \Rightarrow y = 48x^{-1}$$

$$\frac{dy}{dx} = -48x^{-2} = -\frac{48}{x^2}$$

$$\text{At } A, m_T = \frac{dy}{dx} = -\frac{48}{x^2} = -\frac{2}{3} \Rightarrow \frac{48}{x^2} = \frac{2}{3}$$

$$\text{Hence, } 2x^2 = 144 \Rightarrow x^2 = 72 \Rightarrow x = \pm\sqrt{72} \Rightarrow x = \pm 6\sqrt{2} \text{ Note: } \sqrt{72} = \sqrt{36} \sqrt{2} = 6\sqrt{2}.$$

$$\text{When } x = 6\sqrt{2} \Rightarrow y = \frac{48}{6\sqrt{2}} = \frac{8}{\sqrt{2}} = \frac{8\sqrt{2}}{\sqrt{2}\sqrt{2}} = 4\sqrt{2}.$$

$$\text{When } x = -6\sqrt{2} \Rightarrow y = \frac{48}{-6\sqrt{2}} = \frac{-8}{\sqrt{2}} = \frac{-8\sqrt{2}}{\sqrt{2}\sqrt{2}} = -4\sqrt{2}.$$

Hence the possible exact coordinates of A are $(6\sqrt{2}, 4\sqrt{2})$ or $(-6\sqrt{2}, -4\sqrt{2})$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 6

Question:

The curve H is defined by the equations $x = \sqrt{3}t$, $y = \frac{\sqrt{3}}{t}$, $t \in \mathbb{R}$, $t \neq 0$.

The point P lies on H with x -coordinate $2\sqrt{3}$. Find:

a a Cartesian equation for the curve H ,

b an equation of the normal to H at P .

The normal to H at P meets H again at the point Q .

c Find the exact coordinates of Q .

Solution:

$$\mathbf{a} \quad xy = \sqrt{3}t \times \left(\frac{\sqrt{3}}{t}\right)$$

$$xy = \frac{3t}{t}$$

Hence, the Cartesian equation of H is $xy = 3$.

$$\mathbf{b} \quad xy = 3 \Rightarrow y = 3x^{-1}$$

$$\frac{dy}{dx} = -3x^{-2} = -\frac{3}{x^2}$$

$$\text{At } x = 2\sqrt{3}, m_T = \frac{dy}{dx} = -\frac{3}{(2\sqrt{3})^2} = -\frac{3}{12} = -\frac{1}{4}$$

$$\text{Gradient of tangent at } P \text{ is } m_T = -\frac{1}{4}.$$

$$\text{So gradient of normal at } P \text{ is } m_N = \frac{-1}{\left(-\frac{1}{4}\right)} = 4.$$

$$\text{At } P, \text{ when } x = 2\sqrt{3}, \Rightarrow 2\sqrt{3} = \sqrt{3}t \Rightarrow t = \frac{2\sqrt{3}}{\sqrt{3}} = 2$$

$$\text{When } t = 2, y = \frac{\sqrt{3}}{2} \Rightarrow P\left(2\sqrt{3}, \frac{\sqrt{3}}{2}\right).$$

$$\mathbf{N}: y - \frac{\sqrt{3}}{2} = 4(x - 2\sqrt{3})$$

$$\mathbf{N}: 2y - \sqrt{3} = 8(x - 2\sqrt{3})$$

$$\mathbf{N}: 2y - \sqrt{3} = 8x - 16\sqrt{3}$$

$$\mathbf{N}: 0 = 8x - 2y - 16\sqrt{3} + \sqrt{3}$$

$$\mathbf{N}: 0 = 8x - 2y - 15\sqrt{3}$$

The equation of the normal to H at P is $8x - 2y - 15\sqrt{3} = 0$.

$$\mathbf{c \text{ Normal } N: } 8x - 2y - 15\sqrt{3} = 0 \quad (1)$$

$$\text{Hyperbola } H: \quad xy = 3 \quad (2)$$

Rearranging (2) gives

$$y = \frac{3}{x}$$

Substituting this equation into (1) gives

$$8x - 2\left(\frac{3}{x}\right) - 15\sqrt{3} = 0$$

$$8x - \left(\frac{6}{x}\right) - 15\sqrt{3} = 0$$

Multiplying both sides by x gives

$$8x^2 - \left(\frac{6}{x}\right) - 15\sqrt{3} = 0$$

$$8x^2 - 6 - 15\sqrt{3}x = 0$$

$$8x^2 - 15\sqrt{3}x - 6 = 0$$

At P , it is already known that $x = 2\sqrt{3}$, so $(x - 2\sqrt{3})$ is a factor of this quadratic equation. Hence,

$$(x - 2\sqrt{3})(8x + \sqrt{3}) = 0$$

$$x = 2\sqrt{3} \text{ (at } P) \text{ or } x = -\frac{\sqrt{3}}{8} \text{ (at } Q).$$

$$\text{At } P, \text{ when } x = -\frac{\sqrt{3}}{8}, \Rightarrow \frac{-\sqrt{3}}{8} = \sqrt{3}t \Rightarrow t = \frac{-\sqrt{3}}{8\sqrt{3}} = -\frac{1}{8}$$

$$\text{When } t = -\frac{1}{8}, y = \frac{\sqrt{3}}{\left(-\frac{1}{8}\right)} = -8\sqrt{3} \Rightarrow Q\left(-\frac{1}{8}\sqrt{3}, -8\sqrt{3}\right).$$

Hence the coordinates of Q are $\left(-\frac{1}{8}\sqrt{3}, -8\sqrt{3}\right)$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise D, Question 7

Question:

The point $P(4t^2, 8t)$ lies on the parabola C with equation $y^2 = 16x$. The point P also lies on the rectangular hyperbola H with equation $xy = 4$.

a Find the value of t , and hence find the coordinates of P .

The normal to H at P meets the x -axis at the point N .

b Find the coordinates of N .

The tangent to C at P meets the x -axis at the point T .

c Find the coordinates of T .

d Hence, find the area of the triangle NPT .

Solution:

a Substituting $x = 4t^2$ and $y = 8t$ into $xy = 4$ gives

$$(4t^2)(8t) = 4 \Rightarrow 32t^3 = 4 \Rightarrow t^3 = \frac{4}{32} = \frac{1}{8}.$$

$$\text{So } t = \sqrt[3]{\left(\frac{1}{8}\right)}.$$

$$\text{When } t = \frac{1}{2}, x = 4\left(\frac{1}{2}\right)^2 = 1.$$

$$\text{When } t = \frac{1}{2}, y = 8\left(\frac{1}{2}\right) = 4.$$

Hence the value of t is $\frac{1}{2}$ and P has coordinates $(1, 4)$.

$$\mathbf{b} \quad xy = 4 \Rightarrow y = 4x^{-1}$$

$$\frac{dy}{dx} = -4x^{-2} = -\frac{4}{x^2}$$

$$\text{At } P(1, 4), m_T = \frac{dy}{dx} = -\frac{4}{(1)^2} = -\frac{4}{1} = -4$$

Gradient of tangent at $P(1, 4)$ is $m_T = -4$.

$$\text{So gradient of normal at } P(1, 4) \text{ is } m_N = \frac{-1}{-4} = \frac{1}{4}.$$

$$\mathbf{N: } y - 4 = \frac{1}{4}(x - 1)$$

$$\mathbf{N: } 4y - 16 = x - 1$$

$$\mathbf{N}: 0 = x - 4y + 15$$

$$\mathbf{N} \text{ cuts } x\text{-axis} \Rightarrow y = 0 \Rightarrow 0 = x + 15 \Rightarrow x = -15$$

Therefore, the coordinates of N are $(-15, 0)$.

c For $P(1, 4)$, $y > 0$, so

$$y^2 = 16x \Rightarrow y = \sqrt{16x} = \sqrt{16} \sqrt{x} = 4\sqrt{x} = 4x^{\frac{1}{2}}$$

$$\text{So } y = 4x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 4\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = 2x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{2}{\sqrt{x}}$$

$$\text{At } P(1, 4), m_T = \frac{dy}{dx} = \frac{2}{\sqrt{1}} = \frac{2}{1} = 2.$$

Gradient of tangent at $P(1, 4)$ is $m_T = 2$.

$$\mathbf{T}: y - 4 = 2(x - 1)$$


$$\mathbf{T}: y - 4 = 2x - 2$$

$$\mathbf{T}: 0 = 2x - y + 2$$

$$\mathbf{T} \text{ cuts } x\text{-axis} \Rightarrow y = 0 \Rightarrow 0 = 2x + 2 \Rightarrow x = -1$$

Therefore, the coordinates of T are $(-1, 0)$.

d



$$\begin{aligned} \text{Using sketch drawn, Area } \Delta NPT &= \text{Area}(R + S) - \text{Area}(S) \\ &= \frac{1}{2}(16)(4) - \frac{1}{2}(2)(4) \\ &= 32 - 4 \\ &= 28 \end{aligned}$$

Therefore, Area $\Delta NPT = 28$

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 1

Question:

The point $P(3t^2, 6t)$ lies on the parabola C with equation $y^2 = 12x$.

a Show that an equation of the tangent to C at P is $yt = x + 3t^2$.

b Show that an equation of the normal to C at P is $xt + y = 3t^3 + 6t$.

Solution:

$$\mathbf{a} \ C: y^2 = 12x \Rightarrow y = \pm\sqrt{12x} = \pm\sqrt{4 \cdot 3} \sqrt{x} = \pm 2\sqrt{3} x^{\frac{1}{2}}$$

$$\text{So } y = \pm 2\sqrt{3} x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = \pm 2\sqrt{3} \left(\frac{1}{2}\right) x^{-\frac{1}{2}} = \pm\sqrt{3} x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \pm \frac{\sqrt{3}}{\sqrt{x}}$$

$$\text{At } P(3t^2, 6t), m_T = \frac{dy}{dx} = \pm \frac{\sqrt{3}}{\sqrt{3t^2}} = \pm \frac{\sqrt{3}}{\sqrt{3}t} = \frac{1}{t}.$$

$$\mathbf{T: } y - 6t = \frac{1}{t}(x - 3t^2)$$

$$\mathbf{T: } ty - 6t^2 = x - 3t^2$$

$$\mathbf{T: } yt = x - 3t^2 + 6t^2$$

$$\mathbf{T: } yt = x + 3t^2$$

The equation of the tangent to C at P is $yt = x + 3t^2$.

b Gradient of tangent at $P(3t^2, 6t)$ is $m_T = \frac{1}{t}$.

So gradient of normal at $P(3t^2, 6t)$ is $m_N = \frac{-1}{\left(\frac{1}{t}\right)} = -t$.

$$\mathbf{N: } y - 6t = -t(x - 3t^2)$$

$$\mathbf{N: } y - 6t = -tx + 3t^3$$

$$\mathbf{N: } xt + y = 3t^3 + 6t.$$

The equation of the normal to C at P is $xt + y = 3t^3 + 6t$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 2

Question:

The point $P\left(6t, \frac{6}{t}\right)$, $t \neq 0$, lies on the rectangular hyperbola H with equation $xy = 36$.

a Show that an equation of the tangent to H at P is $x + t^2y = 12t$.

b Show that an equation of the normal to H at P is $t^3x - ty = 6(t^4 - 1)$.

Solution:

a $H: xy = 36 \Rightarrow y = 36x^{-1}$

$$\frac{dy}{dx} = -36x^{-2} = -\frac{36}{x^2}$$

$$\text{At } P\left(6t, \frac{6}{t}\right), m_T = \frac{dy}{dx} = -\frac{36}{(6t)^2} = -\frac{36}{36t^2} = -\frac{1}{t^2}$$

$$\text{T: } y - \frac{6}{t} = -\frac{1}{t^2}(x - 6t) \quad (\text{Now multiply both sides by } t^2.)$$

$$\text{T: } t^2y - 6t = -(x - 6t)$$

$$\text{T: } t^2y - 6t = -x + 6t$$

$$\text{T: } x + t^2y = 6t + 6t$$

$$\text{T: } x + t^2y = 12t$$

The equation of the tangent to H at P is $x + t^2y = 12t$.

b Gradient of tangent at $P\left(6t, \frac{6}{t}\right)$ is $m_T = -\frac{1}{t^2}$.

So gradient of normal at $P\left(6t, \frac{6}{t}\right)$ is $m_N = \frac{-1}{\left(-\frac{1}{t^2}\right)} = t^2$.

$$\text{N: } y - \frac{6}{t} = t^2(x - 6t) \quad (\text{Now multiply both sides by } t.)$$

$$\text{N: } ty - 6 = t^3(x - 6t)$$

$$\text{N: } ty - 6 = t^3x - 6t^4$$

$$\text{N: } 6t^4 - 6 = t^3x - ty$$

$$\text{N: } 6(t^4 - 1) = t^3x - ty$$

The equation of the normal to H at P is $t^3x - ty = 6(t^4 - 1)$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 3

Question:

The point $P(5t^2, 10t)$ lies on the parabola C with equation $y^2 = 4ax$, where a is a constant and $t \neq 0$.

a Find the value of a .

b Show that an equation of the tangent to C at P is $yt = x + 5t^2$.

The tangent to C at P cuts the x -axis at the point X and the y -axis at the point Y . The point O is the origin of the coordinate system.

c Find, in terms of t , the area of the triangle OXY .

Solution:

a Substituting $x = 5t^2$ and $y = 10t$ into $y^2 = 4ax$ gives

$$(10t)^2 = 4(a)(5t^2) \Rightarrow 100t^2 = 20t^2a \Rightarrow a = \frac{100t^2}{20t^2} = 5$$

So, $a = 5$.

b When $a = 5$, $y^2 = 4(5)x \Rightarrow y^2 = 20x$.

$$C: y^2 = 20x \Rightarrow y = \pm\sqrt{20x} = \pm\sqrt{4}\sqrt{5}\sqrt{x} = \pm 2\sqrt{5}x^{\frac{1}{2}}$$

$$\text{So } y = \pm 2\sqrt{5}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = \pm 2\sqrt{5}\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \pm\sqrt{5}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \pm \frac{\sqrt{5}}{\sqrt{x}}$$

$$\text{At } P(5t^2, 10t), m_T = \frac{dy}{dx} = \frac{\sqrt{5}}{\sqrt{5t^2}} = \frac{\sqrt{5}}{\sqrt{5}t} = \frac{1}{t}$$

$$\text{T: } y - 10t = \frac{1}{t}(x - 5t^2)$$

$$\text{T: } ty - 10t^2 = x - 5t^2$$

$$\text{T: } yt = x - 5t^2 + 10t^2$$

$$\text{T: } yt = x + 5t^2$$

Therefore, the equation of the tangent to C at P is $yt = x + 5t^2$.

For $(at^2, 2at)$ on $y^2 = 4ax$

We always get $\frac{d}{dx}(y^2) = 4a$

$$2y \frac{dy}{dx} = 4a \frac{dy}{dx} = \frac{2a}{y} = \frac{2a}{2ac} = \frac{1}{t}$$


c T: $yt = x + 5t^2$

T cuts x -axis $\Rightarrow y = 0 \Rightarrow 0 = x + 5t^2 \Rightarrow x = -5t^2$

Hence the coordinates of X are $(-5t^2, 0)$.

T cuts y -axis $\Rightarrow x = 0 \Rightarrow yt = 5t^2 \Rightarrow y = 5t$

Hence the coordinates of Y are $(0, 5t)$.



Using sketch drawn, Area $\triangle OXY = \frac{1}{2}(5t^2)(5t)$
 $= \frac{25}{2}t^3$

Therefore, Area $\triangle OXY = \frac{25}{2}t^3$

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 4

Question:

The point $P(at^2, 2at)$, $t \neq 0$, lies on the parabola C with equation $y^2 = 4ax$, where a is a positive constant.

a Show that an equation of the tangent to C at P is $ty = x + at^2$.

The tangent to C at the point A and the tangent to C at the point B meet at the point with coordinates $(-4a, 3a)$.

b Find, in terms of a , the coordinates of A and the coordinates of B .

Solution:

$$\mathbf{a} \text{ } C: y^2 = 4ax \Rightarrow y = \pm\sqrt{4ax} = \sqrt{4}\sqrt{a}\sqrt{x} = 2\sqrt{a}x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{a}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{a}\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \sqrt{a}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{x}}$$

$$\text{At } P(at^2, 2at), m_T = \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{at^2}} = \frac{\sqrt{a}}{\sqrt{a}t} = \frac{1}{t}.$$

$$\mathbf{T}: y - 2at = \frac{1}{t}(x - at^2)$$

$$\mathbf{T}: ty - 2at^2 = x - at^2$$

$$\mathbf{T}: ty = x - at^2 + 2at^2$$

$$\mathbf{T}: ty = x + at^2$$

The equation of the tangent to C at P is $ty = x + at^2$.

b As the tangent \mathbf{T} goes through $(-4a, 3a)$, then substitute $x = -4a$ and $y = 3a$ into \mathbf{T} .

$$t(3a) = -4a + at^2$$

$$0 = at^2 - 3at - 4a$$

$$t^2 - 3t - 4 = 0$$

$$(t+1)(t-4) = 0$$

$$t = -1, 4$$

$$\text{When } t = -1, x = a(-1)^2 = a, y = 2a(-1) = -2a \Rightarrow (a, -2a).$$

$$\text{When } t = 4, x = a(4)^2 = 16a, y = 2a(4) = 8a \Rightarrow (16a, 8a).$$

The coordinates of A and B are $(a, -2a)$ and $(16a, 8a)$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 5

Question:

The point $P\left(4t, \frac{4}{t}\right)$, $t \neq 0$, lies on the rectangular hyperbola H with equation $xy = 16$.

a Show that an equation of the tangent to C at P is $x + t^2y = 8t$.

The tangent to H at the point A and the tangent to H at the point B meet at the point X with y -coordinate 5. X lies on the directrix of the parabola C with equation $y^2 = 16x$.

b Write down the coordinates of X .

c Find the coordinates of A and B .

d Deduce the equations of the tangents to H which pass through X . Give your answers in the form $ax + by + c = 0$, where a , b and c are integers.

Solution:

a H : $xy = 16 \Rightarrow y = 16x^{-1}$

$$\frac{dy}{dx} = -16x^{-2} = -\frac{16}{x^2}$$

$$\text{At } P\left(4t, \frac{4}{t}\right), m_T = \frac{dy}{dx} = -\frac{16}{(4t)^2} = -\frac{16}{16t^2} = -\frac{1}{t^2}$$

$$\mathbf{T}: y - \frac{4}{t} = -\frac{1}{t^2}(x - 4t) \quad (\text{Now multiply both sides by } t^2.)$$

$$\mathbf{T}: t^2y - 4t = -(x - 4t)$$

$$\mathbf{T}: t^2y - 4t = -x + 4t$$

$$\mathbf{T}: x + t^2y = 4t + 4t$$

$$\mathbf{T}: x + t^2y = 8t$$

The equation of the tangent to H at P is $x + t^2y = 8t$.

b $y^2 = 16x$. So $4a = 16$, gives $a = \frac{16}{4} = 4$.

So the directrix has equation $x + 4 = 0$ or $x = -4$.

Therefore at X , $x = -4$ and as stated $y = 5$.

The coordinates of X are $(-4, 5)$.

$$\mathbf{c} \mathbf{T}: x + t^2y = 8t$$

As the tangent \mathbf{T} goes through $(-4, 5)$, then substitute $x = -4$ and $y = 5$ into \mathbf{T} .

$$(-4) + t^2(5) = 8t$$

$$5t^2 - 4 = 8t$$

$$5t^2 - 8t - 4 = 0$$

$$(t-2)(5t+2) = 0$$

$$t = 2, -\frac{2}{5}$$

When $t = 2$, $x = 4(2) = 8$, $y = \frac{4}{2} = 2 \Rightarrow (8, 2)$.

When $t = -\frac{2}{5}$, $x = 4(-\frac{2}{5}) = -\frac{8}{5}$, $y = \frac{4}{(-\frac{2}{5})} = -10 \Rightarrow (-\frac{8}{5}, -10)$.

The coordinates of A and B are $(8, 2)$ and $(-\frac{8}{5}, -10)$.

d Substitute $t = 2$ and $t = -\frac{2}{5}$ into **T** to find the equations of the tangents to H that go through the point X .

When $t = 2$, **T**: $x + 4y = 16 \Rightarrow x + 4y - 16 = 0$

When $t = -\frac{2}{5}$, **T**: $x + \left(-\frac{2}{5}\right)^2 y = 8\left(-\frac{2}{5}\right)$

T: $x + \frac{4}{25}y = -\frac{16}{5}$

T: $25x + 4y = -80$

T: $25x + 4y + 80 = 0$

Hence the equations of the tangents are $x + 4y - 16 = 0$ and $25x + 4y + 80 = 0$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 6

Question:

The point $P(at^2, 2at)$ lies on the parabola C with equation $y^2 = 4ax$, where a is a constant and $t \neq 0$. The tangent to C at P cuts the x -axis at the point A .

a Find, in terms of a and t , the coordinates of A .

The normal to C at P cuts the x -axis at the point B .

b Find, in terms of a and t , the coordinates of B .

c Hence find, in terms of a and t , the area of the triangle APB .

Solution:

$$\mathbf{a} \text{ } C: y^2 = 4ax \Rightarrow y = \pm\sqrt{4ax} = \sqrt{4} \sqrt{a} \sqrt{x} = 2\sqrt{a} x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{a} x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{a} \left(\frac{1}{2}\right) x^{-\frac{1}{2}} = \sqrt{a} x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{x}}$$

$$\text{At } P(at^2, 2at), m_T = \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{at^2}} = \frac{\sqrt{a}}{\sqrt{a}t} = \frac{1}{t}.$$

$$\mathbf{T}: y - 2at = \frac{1}{t}(x - at^2)$$

$$\mathbf{T}: ty - 2at^2 = x - at^2$$

$$\mathbf{T}: ty = x - at^2 + 2at^2$$

$$\mathbf{T}: ty = x + at^2$$

\mathbf{T} cuts x -axis $\Rightarrow y = 0$. So,

$$0 = x + at^2 \Rightarrow x = -at^2$$

The coordinates of A are $(-at^2, 0)$.

b Gradient of tangent at $P(at^2, 2at)$ is $m_T = \frac{1}{t}$.

So gradient of normal at $P(at^2, 2at)$ is $m_N = \frac{-1}{\left(\frac{1}{t}\right)} = -t$.

$$\mathbf{N}: y - 2at = -t(x - at^2)$$

$$\mathbf{N:} \quad y - 2at = -tx + at^3$$

\mathbf{N} cuts x -axis $\Rightarrow y = 0$. So,


$$0 - 2at = -tx + at^3$$

$$tx = 2at + at^3$$

$$x = 2a + at^2$$

The coordinates of B are $(2a + at^2, 0)$.

c



$$\begin{aligned} \text{Using sketch drawn, Area } \triangle APB &= \frac{1}{2}(2a + 2at^2)(2at) \\ &= at(2a + 2at^2) \\ &= 2a^2t(1 + t^2) \end{aligned}$$

Therefore, Area $\triangle APB = 2a^2t(1 + t^2)$

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 7

Question:

The point $P(2t^2, 4t)$ lies on the parabola C with equation $y^2 = 8x$.

a Show that an equation of the normal to C at P is $xt + y = 2t^3 + 4t$.

The normals to C at the points R, S and T meet at the point $(12, 0)$.

b Find the coordinates of R, S and T .

c Deduce the equations of the normals to C which all pass through the point $(12, 0)$.

Solution:

$$\mathbf{a} \text{ } C: y^2 = 8x \Rightarrow y = \pm\sqrt{8x} = \sqrt{4}\sqrt{2}\sqrt{x} = 2\sqrt{2}x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{2}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{2}\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \sqrt{2}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{2}}{\sqrt{x}}$$

$$\text{At } P(2t^2, 4t), m_T = \frac{dy}{dx} = \frac{\sqrt{2}}{\sqrt{2t^2}} = \frac{\sqrt{2}}{\sqrt{2}t} = \frac{1}{t}.$$

$$\text{Gradient of tangent at } P(2t^2, 4t) \text{ is } m_T = \frac{1}{t}.$$

$$\text{So gradient of normal at } P(2t^2, 4t) \text{ is } m_N = \frac{-1}{\left(\frac{1}{t}\right)} = -t.$$

$$\mathbf{N}: y - 4t = -t(x - 2t^2)$$

$$\mathbf{N}: y - 4t = -tx + 2t^3$$

$$\mathbf{N}: xt + y = 2t^3 + 4t.$$

The equation of the normal to C at P is $xt + y = 2t^3 + 4t$.

b As the normals go through $(12, 0)$, then substitute $x = 12$ and $y = 0$ into \mathbf{N} .

$$(12)t + 0 = 2t^3 + 4t$$

$$12t = 2t^3 + 4t$$

$$0 = 2t^3 + 4t - 12t$$

$$0 = 2t^3 - 8t$$

$$t^3 - 4t = 0$$

$$t(t^2 - 4) = 0$$

$$t(t-2)(t+2) = 0$$

$$t = 0, 2, -2$$

When $t = 0$, $x = 2(0)^2 = 0$, $y = 4(0) = 0 \Rightarrow (0, 0)$.

When $t = 2$, $x = 2(2)^2 = 8$, $y = 4(2) = 8 \Rightarrow (8, 8)$.

When $t = -2$, $x = 2(-2)^2 = 8$, $y = 4(-2) = -8 \Rightarrow (8, -8)$.

The coordinates of R , S and T are $(0, 0)$, $(8, 8)$ and $(8, -8)$.

c Substitute $t = 0, 2, -2$ into $xt + y = 2t^3 + 4t$ to find the equations of the normals to H that go through the point $(12, 0)$.

When $t = 0$, **N:** $0 + y = 0 + 0 \Rightarrow y = 0$

When $t = 2$, **N:** $x(2) + y = 2(8) + 4(2)$

N: $2x + y = 24$

N: $2x + y - 24 = 0$

When $t = -2$, **N:** $x(-2) + y = 2(-8) + 4(-2)$

N: $-2x + y = -24$

N: $2x - y - 24 = 0$

Hence the equations of the normals are $y = 0$, $2x + y - 24 = 0$ and $2x - y - 24 = 0$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 8

Question:

The point $P(at^2, 2at)$ lies on the parabola C with equation $y^2 = 4ax$, where a is a positive constant and $t \neq 0$. The tangent to C at P meets the y -axis at Q .

a Find in terms of a and t , the coordinates of Q .

The point S is the focus of the parabola.

b State the coordinates of S .

c Show that PQ is perpendicular to SQ .

Solution:

$$\mathbf{a} \text{ } C: y^2 = 4ax \Rightarrow y = \sqrt{4ax} = \sqrt{4} \sqrt{a} \sqrt{x} = 2\sqrt{a}x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{a}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{a} \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \sqrt{a}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{x}}$$

$$\text{At } P(at^2, 2at), m_T = \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{at^2}} = \frac{\sqrt{a}}{\sqrt{a}t} = \frac{1}{t}.$$

$$\mathbf{T}: y - 2at = \frac{1}{t}(x - at^2)$$

$$\mathbf{T}: ty - 2at^2 = x - at^2$$

$$\mathbf{T}: ty = x - at^2 + 2at^2$$

$$\mathbf{T}: ty = x + at^2$$

\mathbf{T} meets y -axis $\Rightarrow x = 0$. So,

$$ty = 0 + at^2 \Rightarrow y = \frac{at^2}{t} \Rightarrow y = at$$

The coordinates of Q are $(0, at)$.

b The focus of a parabola with equation $y^2 = 4ax$ has coordinates $(a, 0)$.

So, the coordinates of S are $(a, 0)$.

c $P(at^2, 2at)$, $Q(0, at)$ and $S(a, 0)$.

$$m_{PQ} = \frac{at - 2at}{0 - at^2} = \frac{-at}{-at^2} = \frac{1}{t}.$$

$$m_{SQ} = \frac{0 - at}{a - 0} = -\frac{at}{a} = -t.$$

Therefore, $m_{PQ} \times m_{SQ} = \frac{1}{t} \times -t = -1$.

So PQ is perpendicular to SQ .

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise E, Question 9

Question:

The point $P(6t^2, 12t)$ lies on the parabola C with equation $y^2 = 24x$.

a Show that an equation of the tangent to the parabola at P is $ty = x + 6t^2$.

The point X has y -coordinate 9 and lies on the directrix of C .

b State the x -coordinate of X .

The tangent at the point B on C goes through point X .

c Find the possible coordinates of B .

Solution:

$$\mathbf{a} \ C: y^2 = 24x \Rightarrow y = \pm\sqrt{24x} = \sqrt{4} \cdot \sqrt{6} \cdot \sqrt{x} = 2\sqrt{6}x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{6}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{6} \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \sqrt{6}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{6}}{\sqrt{x}}$$

$$\text{At } P(6t^2, 12t), \ m_T = \frac{dy}{dx} = \frac{\sqrt{6}}{\sqrt{6t^2}} = \frac{\sqrt{6}}{\sqrt{6}t} = \frac{1}{t}.$$

$$\mathbf{T}: y - 12t = \frac{1}{t}(x - 6t^2)$$

$$\mathbf{T}: ty - 12t^2 = x - 6t^2$$

$$\mathbf{T}: ty = x - 6t^2 + 12t^2$$

$$\mathbf{T}: ty = x + 6t^2$$

The equation of the tangent to C at P is $ty = x + 6t^2$.

$$\mathbf{b} \ y^2 = 24x. \text{ So } 4a = 24, \text{ gives } a = \frac{24}{4} = 6.$$

So the directrix has equation $x + 6 = 0$ or $x = -6$.

Therefore at X , $x = -6$.

c T: $ty = x + 6t^2$ and the coordinates of X are $(-6, 9)$.

As the tangent \mathbf{T} goes through $(-6, 9)$, then substitute $x = -6$ and $y = 9$ into \mathbf{T} .

$$t(9) = -6 + 6t^2$$

$$0 = 6t^2 - 9t - 6$$

$$2t^2 - 3t - 2 = 0$$

$$(t-2)(2t+1) = 0$$

$$t = 2, -\frac{1}{2}$$

$$\text{When } t = 2, \quad x = 6(2)^2 = 24, \quad y = 12(2) = 24 \Rightarrow (24, 24).$$

$$\text{When } t = -\frac{1}{2}, \quad x = 6\left(-\frac{1}{2}\right)^2 = \frac{3}{2}, \quad y = 12\left(-\frac{1}{2}\right) = -6 \Rightarrow \left(\frac{3}{2}, -6\right).$$

The possible coordinates of B are $(24, 24)$ and $\left(\frac{3}{2}, -6\right)$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 1

Question:

A parabola C has equation $y^2 = 12x$. The point S is the focus of C .

a Find the coordinates of S .

The line l with equation $y = 3x$ intersects C at the point P where $y > 0$.

b Find the coordinates of P .

c Find the area of the triangle OPS , where O is the origin.

Solution:

a $y^2 = 12x$. So $4a = 12$, gives $a = \frac{12}{4} = 3$.

So the focus S , has coordinates $(3, 0)$.

b Line l : $y = 3x$ (1)

Parabola C : $y^2 = 12x$ (2)

Substituting (1) into (2) gives

$$(3x)^2 = 12x$$

$$9x^2 = 12x$$

$$9x^2 - 12x = 0$$

$$3x(3x - 4) = 0$$

$$x = 0, \frac{4}{3}$$


Substituting these values of x back into equation (1) gives

$$x = 0, y = 3(0) = 0 \Rightarrow (0, 0)$$

$$x = \frac{4}{3}, y = 3\left(\frac{4}{3}\right) = 4 \Rightarrow \left(\frac{4}{3}, 4\right)$$

As $y > 0$, the coordinates of P are $\left(\frac{4}{3}, 4\right)$.

c



$$\begin{aligned}\text{Using sketch drawn, Area } \triangle OPS &= \frac{1}{2}(3)(4) \\ &= \frac{1}{2}(12) \\ &= 6\end{aligned}$$

Therefore, Area $\triangle OPS = 6$

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 2

Question:

A parabola C has equation $y^2 = 24x$. The point P with coordinates $(k, 6)$, where k is a constant lies on C .

a Find the value of k .

The point S is the focus of C .

b Find the coordinates of S .

The line l passes through S and P and intersects the directrix of C at the point D .

c Show that an equation for l is $4x + 3y - 24 = 0$.

d Find the area of the triangle OPD , where O is the origin.

Solution:

a $(k, 6)$ lies on $y^2 = 24x$ gives

$$6^2 = 24k \Rightarrow 36 = 24k \Rightarrow \frac{36}{24} = k \Rightarrow k = \frac{3}{2}.$$

b $y^2 = 24x$. So $4a = 24$, gives $a = \frac{24}{4} = 6$.

So the focus S , has coordinates $(6, 0)$.

c The point P and S have coordinates $P\left(\frac{3}{2}, 6\right)$ and $S(6, 0)$.

$$m_l = m_{PS} = \frac{0-6}{6-\frac{3}{2}} = \frac{-6}{\frac{9}{2}} = -\frac{12}{9} = -\frac{4}{3}$$

$$l: y - 0 = -\frac{4}{3}(x - 6)$$

$$l: 3y = -4(x - 6)$$

$$l: 3y = -4x + 24$$

$$l: 4x + 3y - 24 = 0$$

Therefore an equation for l is $4x + 3y - 24 = 0$.

d From (b), as $a = 6$, an equation of the directrix is $x + 6 = 0$ or $x = -6$. Substituting $x = -6$ into l gives:


$$4(-6) + 3y - 24 = 0$$

$$3y = 24 + 24$$

$$3y = 48$$

$$y = 16$$

Hence the coordinates of D are $(-6, 16)$.



Using the sketch and the regions as labeled you can find the area required. Let $\text{Area } \triangle OPD = \text{Area}(R)$

Method 1

$$\begin{aligned}
 \text{Area}(R) &= \text{Area}(RST) - \text{Area}(S) - \text{Area}(T) \\
 &= \frac{1}{2}(16+6)\left(\frac{15}{2}\right) - \frac{1}{2}(6)(16) - \frac{1}{2}\left(\frac{3}{2}\right)(6) \\
 &= \frac{1}{2}(22)\left(\frac{15}{2}\right) - (3)(16) - \left(\frac{3}{2}\right)(3) \\
 &= \left(\frac{165}{2}\right) - 48 - \left(\frac{9}{2}\right) \\
 &= 30
 \end{aligned}$$

Therefore, $\text{Area } \triangle OPD = 30$

Method 2

$$\begin{aligned}
 \text{Area}(R) &= \text{Area}(RSTU) - \text{Area}(S) - \text{Area}(TU) \\
 &= \frac{1}{2}(12)(16) - \frac{1}{2}(6)(16) - \frac{1}{2}(6)(6) \\
 &= 96 - 48 - 18 \\
 &= 30
 \end{aligned}$$

Therefore, $\text{Area } \triangle OPD = 30$

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 3

Question:

The parabola C has parametric equations $x = 12t^2$, $y = 24t$. The focus to C is at the point S .

a Find a Cartesian equation of C .

The point P lies on C where $y > 0$. P is 28 units from S .

b Find an equation of the directrix of C .

c Find the exact coordinates of the point P .

d Find the area of the triangle OSP , giving your answer in the form $k\sqrt{3}$, where k is an integer.

Solution:

a $y = 24t$

So $t = \frac{y}{24}$ (1)

$$x = 12t^2 \quad (2)$$

Substitute (1) into (2):

$$x = 12\left(\frac{y}{24}\right)^2$$

So $x = \frac{12y^2}{576}$ simplifies to $x = \frac{y^2}{48}$

Hence, the Cartesian equation of C is $y^2 = 48x$.

b $y^2 = 48x$. So $4a = 48$, gives $a = \frac{48}{4} = 12$.

Therefore an equation of the directrix of C is $x + 12 = 0$ or $x = -12$.

c


From (b), as $a = 12$, the coordinates of S , the focus to C are $(12, 0)$. Hence, drawing a sketch gives,

The (shortest) distance of P to the line $x = -12$ is the distance XP .

The distance $SP = 28$.

The focus-directrix property implies that $SP = XP = 28$.

The directrix has equation $x = -12$.




Therefore the x -coordinate of P is
 $x = 28 - 12 = 16$.

$$\text{When } x = 16, y^2 = 48(16) \Rightarrow y^2 = 3(16)^2$$

$$\text{As } y > 0, \text{ then } y = \sqrt{3(16)^2} = 16\sqrt{3}.$$

Hence the exact coordinates of P are $(16, 16\sqrt{3})$.

d



Using the sketch and the regions as labeled you can find the area required. Let $\text{Area } \triangle OSP = \text{Area}(A)$

$$\begin{aligned} \text{Area}(A) &= \text{Area}(AB) - \text{Area}(B) \\ &= \frac{1}{2}(16)(16\sqrt{3}) - \frac{1}{2}(4)(16\sqrt{3}) \\ &= 128\sqrt{3} - 32\sqrt{3} \\ &= 96\sqrt{3} \end{aligned}$$

Therefore, $\text{Area } \triangle OSP = 96\sqrt{3}$ and $k = 96$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 4

Question:

The point $(4t^2, 8t)$ lies on the parabola C with equation $y^2 = 16x$. The line l with equation $4x - 9y + 32 = 0$ intersects the curve at the points P and Q .

a Find the coordinates of P and Q .

b Show that an equation of the normal to C at $(4t^2, 8t)$ is $xt + y = 4t^3 + 8t$.

c Hence, find an equation of the normal to C at P and an equation of the normal to C at Q .

The normal to C at P and the normal to C at Q meet at the point R .

d Find the coordinates of R and show that R lies on C .

e Find the distance OR , giving your answer in the form $k\sqrt{97}$, where k is an integer.

Solution:

a Method 1

Line: $4x - 9y + 32 = 0$ (1)

Parabola C : $y^2 = 16x$ (2)

Multiplying (1) by 4 gives

$$16x - 36y + 128 = 0 \quad (3)$$

Substituting (2) into (3) gives

$$\begin{aligned} y^2 - 36y + 128 &= 0 \\ (y - 4)(y - 32) &= 0 \\ y &= 4, 32 \end{aligned}$$

When $y = 4$, $4^2 = 16x \Rightarrow x = \frac{16}{16} = 1 \Rightarrow (1, 4)$.

When $y = 32$, $32^2 = 16x \Rightarrow x = \frac{1024}{16} = 64 \Rightarrow (64, 32)$.

The coordinates of P and Q are $(1, 4)$ and $(64, 32)$.

Method 2

Line: $4x - 9y + 32 = 0$ (1)

Parabola C : $x = 4t^2, y = 8t$ (2)

Substituting (2) into (1) gives

$$4(4t^2) - 9(8t) + 32 = 0$$

$$16t^2 - 72t + 32 = 0$$

$$2t^2 - 9t + 4 = 0$$

$$(2t-1)(t-4) = 0$$

$$t = \frac{1}{2}, 4$$

$$\text{When } t = \frac{1}{2}, \quad x = 4\left(\frac{1}{2}\right)^2 = 1, \quad y = 8\left(\frac{1}{2}\right) = 4 \Rightarrow (1, 4).$$

$$\text{When } t = 4, \quad x = 4(4)^2 = 64, \quad y = 8(4) = 32 \Rightarrow (64, 32).$$

The coordinates of P and Q are $(1, 4)$ and $(64, 32)$.

$$\mathbf{b} \text{ C: } y^2 = 16x \Rightarrow y = \sqrt{16x} = \sqrt{16} \cdot \sqrt{x} = 4x^{\frac{1}{2}}$$

$$\text{So } y = 4x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 4\left(\frac{1}{2}\right)x^{-\frac{1}{2}} = 2x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{2}{\sqrt{x}}$$

$$\text{At } (4t^2, 8t), m_T = \frac{dy}{dx} = \frac{2}{\sqrt{4t^2}} = \frac{2}{2t} = \frac{1}{t}.$$

$$\text{Gradient of tangent at } (4t^2, 8t) \text{ is } m_T = \frac{1}{t}.$$

$$\text{So gradient of normal at } (4t^2, 8t) \text{ is } m_N = \frac{-1}{\left(\frac{1}{t}\right)} = -t.$$

$$\mathbf{N: } y - 8t = -t(x - 4t^2)$$

$$\mathbf{N: } y - 8t = -tx + 4t^3$$

$$\mathbf{N: } xt + y = 4t^3 + 8t.$$

The equation of the normal to C at $(4t^2, 8t)$ is $xt + y = 4t^3 + 8t$.

\mathbf{c} Without loss of generality, from part (a) P has coordinates $(1, 4)$ when $t = \frac{1}{2}$ and Q has coordinates $(64, 32)$ when $t = 4$.

$$\text{When } t = \frac{1}{2},$$

$$\mathbf{N: } x\left(\frac{1}{2}\right) + y = 4\left(\frac{1}{2}\right)^3 + 8\left(\frac{1}{2}\right)$$

$$\mathbf{N: } \frac{1}{2}x + y = \frac{1}{2} + 4$$

$$\mathbf{N: } x + 2y = 1 + 8$$

$$\mathbf{N: } x + 2y - 9 = 0$$

$$\text{When } t = 4,$$

$$\mathbf{N:} \quad x(4) + y = 4(4)^3 + 8(4)$$

$$\mathbf{N:} \quad 4x + y = 256 + 32$$

$$\mathbf{N:} \quad 4x + y - 288 = 0$$

d The normals to C at P and at Q are $x + 2y - 9 = 0$ and $4x + y - 288 = 0$

$$\mathbf{N}_1: \quad x + 2y - 9 = 0 \quad \mathbf{(1)}$$

$$\mathbf{N}_2: \quad 4x + y - 288 = 0 \quad \mathbf{(2)}$$

Multiplying **(2)** by 2 gives

$$2 \times \mathbf{(2):} \quad 8x + 2y - 576 = 0 \quad \mathbf{(3)}$$

$$\mathbf{(3) - (1):} \quad 7x - 567 = 0$$

$$\Rightarrow 7x = 567 \Rightarrow x = \frac{567}{7} = 81$$

$$\mathbf{(2)} \Rightarrow \quad y = 288 - 4(81) = 288 - 324 = -36$$

The coordinates of R are $(81, -36)$.

$$\text{When } y = -36, \text{ LHS} = y^2 = (-36)^2 = 1296$$

$$\text{When } x = 81, \text{ RHS} = 16x = 16(81) = 1296$$

As $LHS = RHS$, R lies on C .

e The coordinates of O and R are $(0, 0)$ and $(81, -36)$.

$$\begin{aligned} OR &= \sqrt{(81-0)^2 + (-36-0)^2} \quad ? \\ &= \sqrt{81^2 + 36^2} \\ &= \sqrt{7857} \\ &= \sqrt{(81)(97)} \\ &= \sqrt{81} \sqrt{97} \\ &= 9\sqrt{97} \end{aligned}$$

Hence the exact distance OR is $9\sqrt{97}$ and $k = 9$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 5

Question:

The point $P(at^2, 2at)$ lies on the parabola C with equation $y^2 = 4ax$, where a is a positive constant. The point Q lies on the directrix of C . The point Q also lies on the x -axis.

a State the coordinates of the focus of C and the coordinates of Q .

The tangent to C at P passes through the point Q .

b Find, in terms of a , the two sets of possible coordinates of P .

Solution:

The focus and directrix of a parabola with equation $y^2 = 4ax$, are $(a, 0)$ and $x + a = 0$ respectively.

a Hence the coordinates of the focus of C are $(a, 0)$.

As Q lies on the x -axis then $y = 0$ and so Q has coordinates $(-a, 0)$.

$$\mathbf{b} \text{ } C: y^2 = 4ax \Rightarrow y = \sqrt{4ax} = \sqrt{4} \sqrt{a} \sqrt{x} = 2\sqrt{a} x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{a} x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{a} \left(\frac{1}{2}\right) x^{-\frac{1}{2}} = \sqrt{a} x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{x}}$$

$$\text{At } P(at^2, 2at), m_T = \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{at^2}} = \frac{\sqrt{a}}{\sqrt{at}} = \frac{1}{t}.$$

$$\mathbf{T: } y - 2at = \frac{1}{t}(x - at^2)$$

$$\mathbf{T: } ty - 2at^2 = x - at^2$$

$$\mathbf{T: } ty = x - at^2 + 2at^2$$

$$\mathbf{T: } ty = x + at^2$$

T passes through $(-a, 0)$, so substitute $x = -a$, $y = 0$ in T .

$$t(0) = -a + at^2 \Rightarrow 0 = -a + at^2 \Rightarrow 0 = -1 + t^2$$

$$\text{So, } t^2 - 1 = 0 \Rightarrow (t-1)(t+1) = 0 \Rightarrow t = 1, -1$$

$$\text{When } t = 1, \quad x = a(1)^2 = a, \quad y = 2a(1) = 2a \quad \Rightarrow (a, 2a).$$

$$\text{When } t = -1, \quad x = a(-1)^2 = a, \quad y = 2a(-1) = -2a \Rightarrow (a, -2a).$$

The possible coordinates of P are $(a, 2a)$ or $(a, -2a)$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 6

Question:

The point $P\left(ct, \frac{c}{t}\right)$, $c > 0$, $t \neq 0$, lies on the rectangular hyperbola H with equation $xy = c^2$.

a Show that the equation of the normal to H at P is $t^3x - ty = c(t^4 - 1)$.

b Hence, find the equation of the normal n to the curve V with the equation $xy = 36$ at the point $(12, 3)$. Give your answer in the form $ax + by = d$, where a , b and d are integers.

The line n meets V again at the point Q .

c Find the coordinates of Q .

Solution:

a H : $xy = c^2 \Rightarrow y = c^2x^{-1}$

$$\frac{dy}{dx} = -c^2x^{-2} = -\frac{c^2}{x^2}$$

At $P\left(ct, \frac{c}{t}\right)$, $m_T = \frac{dy}{dx} = -\frac{c^2}{(ct)^2} = -\frac{c^2}{c^2t^2} = -\frac{1}{t^2}$

Gradient of tangent at $P\left(ct, \frac{c}{t}\right)$ is $m_T = -\frac{1}{t^2}$.

So gradient of normal at $P\left(ct, \frac{c}{t}\right)$ is $m_N = \frac{-1}{\left(-\frac{1}{t^2}\right)} = t^2$.

N: $y - \frac{c}{t} = t^2(x - ct)$ (Now multiply both sides by t)

N: $ty - c = t^3(x - ct)$

N: $ty - c = t^3x - ct^4$

N: $ct^4 - c = t^3x - ty$

N: $t^3x - ty = ct^4 - c$

N: $t^3x - ty = c(t^4 - 1)$

The equation of the normal to H at P is $t^3x - ty = c(t^4 - 1)$.

b Comparing $xy = 36$ with $xy = c^2$ gives $c = 6$ and comparing the point $(12, 3)$ with $\left(ct, \frac{c}{t}\right)$ gives

$$ct = 12 \Rightarrow (6)t = 12 \Rightarrow t = 2. \text{ Therefore,}$$

n: $(2)^3x - (2)y = 6((2)^4 - 1)$

$$n: 8x - 2y = 6(15)$$

$$n: 8x - 2y = 90$$

$$n: 4x - y = 45$$

An equation for n is $4x - y = 45$.

$$\text{c Normal } n: \quad 4x - y = 45 \quad \text{(1)}$$

$$\text{Hyperbola } V: \quad xy = 36 \quad \text{(2)}$$

Rearranging (2) gives

$$y = \frac{36}{x}$$

Substituting this equation into (1) gives

$$4x - \left(\frac{36}{x}\right) = 45$$

Multiplying both sides by x gives

$$4x^2 - 36 = 45x$$

$$4x^2 - 45x - 36 = 0$$

$$(x - 12)(4x + 3) = 0$$

$$x = 12, -\frac{3}{4}$$

It is already known that $x = 12$. So at Q , $x = -\frac{3}{4}$.

Substituting $x = -\frac{3}{4}$ into $y = \frac{36}{x}$ gives

$$y = \frac{36}{\left(-\frac{3}{4}\right)} = -36\left(\frac{4}{3}\right) = -48.$$

Hence the coordinates of Q are $\left(-\frac{3}{4}, -48\right)$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 7

Question:

A rectangular hyperbola H has equation $xy = 9$. The lines l_1 and l_2 are tangents to H . The gradients of l_1 and l_2 are both $-\frac{1}{4}$. Find the equations of l_1 and l_2 .

Solution:

$$H: xy = 9 \Rightarrow y = 9x^{-1}$$

$$\frac{dy}{dx} = -9x^{-2} = -\frac{9}{x^2}$$

Gradients of tangent lines l_1 and l_2 are both $-\frac{1}{4}$ implies

$$-\frac{9}{x^2} = -\frac{1}{4}$$

$$\Rightarrow x^2 = 36$$

$$\Rightarrow x = \pm\sqrt{36}$$

$$\Rightarrow x = \pm 6$$

$$\text{When } x = 6, \quad 6y = 9 \Rightarrow y = \frac{9}{6} = \frac{3}{2} \Rightarrow \left(6, \frac{3}{2}\right).$$

$$\text{When } x = -6, \quad -6y = 9 \Rightarrow y = \frac{9}{-6} = -\frac{3}{2} \Rightarrow \left(-6, -\frac{3}{2}\right).$$

At $\left(6, \frac{3}{2}\right)$, $m_T = -\frac{1}{4}$ and

$$\mathbf{T: } y - \frac{3}{2} = -\frac{1}{4}(x - 6)$$

$$\mathbf{T: } 4y - 6 = -1(x - 6)$$

$$\mathbf{T: } 4y - 6 = -x + 6$$

$$\mathbf{T: } x + 4y - 12 = 0$$

At $\left(-6, -\frac{3}{2}\right)$, $m_T = -\frac{1}{4}$ and

$$\mathbf{T: } y + \frac{3}{2} = -\frac{1}{4}(x + 6)$$

$$\mathbf{T: } 4y + 6 = -1(x + 6)$$

$$\mathbf{T: } 4y + 6 = -x - 6$$

$$\mathbf{T: } x + 4y + 12 = 0$$

The equations for l_1 and l_2 are $x + 4y - 12 = 0$ and $x + 4y + 12 = 0$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 8

Question:

The point P lies on the rectangular hyperbola $xy = c^2$, where $c > 0$. The tangent to the rectangular hyperbola at the point $P\left(ct, \frac{c}{t}\right)$, $t > 0$, cuts the x -axis at the point X and cuts the y -axis at the point Y .

a Find, in terms of c and t , the coordinates of X and Y .

b Given that the area of the triangle OXY is 144, find the exact value of c .

Solution:

$$\mathbf{a} \text{ H: } xy = c^2 \Rightarrow y = c^2x^{-1}$$

$$\frac{dy}{dx} = -c^2x^{-2} = -\frac{c^2}{x^2}$$

$$\text{At } P\left(ct, \frac{c}{t}\right), m_T = \frac{dy}{dx} = -\frac{c^2}{(ct)^2} = -\frac{c^2}{c^2t^2} = -\frac{1}{t^2}$$

$$\mathbf{T: } y - \frac{c}{t} = -\frac{1}{t^2}(x - ct) \quad (\text{Now multiply both sides by } t^2.)$$

$$\mathbf{T: } t^2y - ct = -(x - ct)$$

$$\mathbf{T: } t^2y - ct = -x + ct$$


$$\mathbf{T: } x + t^2y = 2ct$$

$$\mathbf{T} \text{ cuts } x\text{-axis} \Rightarrow y = 0 \Rightarrow x + t^2(0) = 2ct \Rightarrow x = 2ct$$

$$\mathbf{T} \text{ cuts } y\text{-axis} \Rightarrow x = 0 \Rightarrow 0 + t^2y = 2ct \Rightarrow y = \frac{2ct}{t^2} = \frac{2c}{t}$$

So the coordinates are $X(2ct, 0)$ and $Y\left(0, \frac{2c}{t}\right)$.

b



$$\text{Using the sketch, area } \triangle OXY = \frac{1}{2}(2ct)\left(\frac{2c}{t}\right) = \frac{4c^2t}{2t} = 2c^2$$

As area $\Delta OXY = 144$, then $2c^2 = 144 \Rightarrow c^2 = 72$

As $c > 0$, $c = \sqrt{72} = \sqrt{36} \sqrt{2} = 6\sqrt{2}$.

Hence the exact value of c is $6\sqrt{2}$.

© Pearson Education Ltd 2008

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 9

Question:

The points $P(4at^2, 4at)$ and $Q(16at^2, 8at)$ lie on the parabola C with equation $y^2 = 4ax$, where a is a positive constant.

a Show that an equation of the tangent to C at P is $2ty = x + 4at^2$.

b Hence, write down the equation of the tangent to C at Q .

The tangent to C at P meets the tangent to C at Q at the point R .

c Find, in terms of a and t , the coordinates of R .

Solution:

$$\mathbf{a} \ C: y^2 = 4ax \Rightarrow y = \pm\sqrt{4ax} = \sqrt{4} \cdot \sqrt{a} \cdot \sqrt{x} = 2\sqrt{a}x^{\frac{1}{2}}$$

$$\text{So } y = 2\sqrt{a}x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 2\sqrt{a} \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \sqrt{a}x^{-\frac{1}{2}}$$

$$\text{So, } \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{x}}$$

$$\text{At } P(4at^2, 4at), m_T = \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{4at^2}} = \frac{\sqrt{a}}{2\sqrt{a}t} = \frac{1}{2t}.$$

$$\mathbf{T: } y - 4at = \frac{1}{2t}(x - 4at^2)$$

$$\mathbf{T: } 2ty - 8at^2 = x - 4at^2$$

$$\mathbf{T: } 2ty = x - 4at^2 + 8at^2$$

$$\mathbf{T: } 2ty = x + 4at^2$$

The equation of the tangent to C at $P(4at^2, 4at)$ is $2ty = x + 4at^2$.

b P has mapped onto Q by replacing t by $2t$, ie. $t \rightarrow 2t$

$$\text{So, } P(4at^2, 4at) \rightarrow Q(16at^2, 8at) = Q(4a(2t)^2, 4a(2t))$$

$$\text{At } Q, \mathbf{T} \text{ becomes } 2(2t)y = x + 4a(2t)^2$$

$$\mathbf{T: } 2(2t)y = x + 4a(2t)^2$$

$$\mathbf{T: } 4ty = x + 4a(4t^2)$$

$$\mathbf{T: } 4ty = x + 16at^2$$

The equation of the tangent to C at $Q(16at^2, 8at)$ is $4ty = x + 16at^2$.

$$\text{c } T_P: 2ty = x + 4at^2 \quad (1)$$

$$T_Q: 4ty = x + 16at^2 \quad (2)$$

(2) – (1) gives

$$2ty = 12at^2$$

$$\text{Hence, } y = \frac{12at^2}{2t} = 6at.$$

Substituting this into **(1)** gives,

$$2t(6at) = x + 4at^2$$

$$12at^2 = x + 4at^2$$

$$12at^2 - 4at^2 = x$$

$$\text{Hence, } x = 8at^2.$$

The coordinates of R are $(8at^2, 6at)$.

Solutionbank FP1

Edexcel AS and A Level Modular Mathematics

Quadratic Equations

Exercise F, Question 10

Question:

A rectangular hyperbola H has Cartesian equation $xy = c^2$, $c > 0$. The point $\left(ct, \frac{c}{t}\right)$, where $t \neq 0$, $t > 0$ is a general point on H .

a Show that an equation of the tangent to H at $\left(ct, \frac{c}{t}\right)$ is $x + t^2y = 2ct$.

The point P lies on H . The tangent to H at P cuts the x -axis at the point X with coordinates $(2a, 0)$, where a is a constant.

b Use the answer to part **a** to show that P has coordinates $\left(a, \frac{c^2}{a}\right)$.

The point Q , which lies on H , has x -coordinate $2a$.

c Find the y -coordinate of Q .

d Hence, find the equation of the line OQ , where O is the origin.

The lines OQ and XP meet at point R .

e Find, in terms of a , the x -coordinate of R .

Given that the line OQ is perpendicular to the line XP ,

f Show that $c^2 = 2a^2$,

g find, in terms of a , the y -coordinate of R .

Solution:

a H : $xy = c^2 \Rightarrow y = c^2x^{-1}$

$$\frac{dy}{dx} = -c^2x^{-2} = -\frac{c^2}{x^2}$$

$$\text{At } \left(ct, \frac{c}{t}\right), m_T = \frac{dy}{dx} = -\frac{c^2}{(ct)^2} = -\frac{c^2}{c^2t^2} = -\frac{1}{t^2}$$

$$\text{T: } y - \frac{c}{t} = -\frac{1}{t^2}(x - ct) \quad (\text{Now multiply both sides by } t^2.)$$

$$\text{T: } t^2y - ct = -(x - ct)$$

$$\text{T: } t^2y - ct = -x + ct$$

$$\text{T: } x + t^2y = 2ct$$

An equation of a tangent to H at $\left(ct, \frac{c}{t}\right)$ is $x + t^2y = 2ct$.

b **T** passes through $X(2a, 0)$, so substitute $x = 2a, y = 0$ into **T**.

$$(2a) + t^2(0) = 2ct \Rightarrow 2a = 2ct \Rightarrow \frac{2a}{2c} = t \Rightarrow t = \frac{a}{c}$$

Substitute $t = \frac{a}{c}$ into $\left(ct, \frac{c}{t}\right)$ gives

$$P\left(c\left(\frac{a}{c}\right), \frac{c}{\left(\frac{a}{c}\right)}\right) = P\left(a, \frac{c^2}{a}\right).$$

Hence P has coordinates $P\left(a, \frac{c^2}{a}\right)$.

c Substituting $x = 2a$ into the curve H gives

$$(2a)y = c^2 \Rightarrow y = \frac{c^2}{2a}.$$

The y-coordinate of Q is $y = \frac{c^2}{2a}$.

d The coordinates of O and Q are $(0, 0)$ and $\left(2a, \frac{c^2}{2a}\right)$.

$$m_{OQ} = \frac{\frac{c^2}{2a} - 0}{2a - 0} = \frac{c^2}{2a(2a)} = \frac{c^2}{4a^2}$$

$$OQ: y - 0 = \frac{c^2}{4a^2}(x - 0)$$

$$OQ: y = \frac{c^2x}{4a^2}. \quad \text{(1)}$$

The equation of OQ is $y = \frac{c^2x}{4a^2}$.

e The coordinates of X and P are $(2a, 0)$ and $\left(a, \frac{c^2}{a}\right)$.

$$m_{XP} = \frac{\frac{c^2}{a} - 0}{a - 2a} = \frac{\frac{c^2}{a}}{-a} = -\frac{c^2}{a^2}$$

$$XP: y - 0 = -\frac{c^2}{a^2}(x - 2a)$$

$$XP: y = -\frac{c^2}{a^2}(x - 2a) \quad \text{(2)}$$

Substituting **(1)** into **(2)** gives,

$$\frac{c^2x}{4a^2} = -\frac{c^2}{a^2}(x - 2a)$$

Cancelling $\frac{c^2}{a^2}$ gives,

$$\frac{x}{4} = -(x - 2a)$$

$$\frac{x}{4} = -x + 2a$$

$$\frac{5x}{4} = 2a$$

$$x = \frac{4(2a)}{5} = \frac{8a}{5}$$

The x -coordinate of R is $\frac{8a}{5}$.

f From earlier parts, $m_{OQ} = \frac{c^2}{4a^2}$ and $m_{XP} = -\frac{c^2}{a^2}$

OP is perpendicular to $XP \Rightarrow m_{OQ} \times m_{XP} = -1$, gives

$$m_{OQ} \times m_{XP} = \left(\frac{c^2}{4a^2}\right)\left(-\frac{c^2}{a^2}\right) = \frac{-c^4}{4a^4} = -1$$

$$-c^4 = -4a^4 \Rightarrow c^4 = 4a^4 \Rightarrow (c^2)^2 = 4a^4$$

$$c^2 = \sqrt{4a^4} = \sqrt{4} \sqrt{a^4} = 2a^2.$$

Hence, $c^2 = 2a^2$, as required.

g At R , $x = \frac{8a}{5}$. Substituting $x = \frac{8a}{5}$ into $y = \frac{c^2 x}{4a^2}$ gives,

$$y = \frac{c^2}{4a^2} \left(\frac{8a}{5}\right) = \frac{8ac^2}{20a^2}$$

and using the $c^2 = 2a^2$ gives,

$$y = \frac{8a(2a^2)}{20a^2} = \frac{16a^3}{20a^2} = \frac{4a}{5}.$$

The y -coordinate of R is $\frac{4a}{5}$.