2 Review Exercise Exercise A, Question 1

Question:



A particle of weight 24 N is held in equilibrium by two light inextensible strings. One string is horizontal. The other string is inclined at an angle of 30° to the horizontal, as shown. The tension in the horizontal string is Q newtons and the tension in the other string is P newtons. Find

a the value of P,

b the value of Q.



2 Review Exercise Exercise A, Question 2

Question:



A particle *P*, of mass 2 kg, is attached to one end of a light string, the other end of which is attached to a fixed point *O*. The particle is held in equilibrium, with *OP* horizontal, by a force of magnitude 30 N applied at an angle α to the horizontal, as shown.

a Find, to the nearest degree, the value of α .

b Find, in N to 3 significant figures, the magnitude of the tension in the string.



2g

30

Ja

representing each of the three forces.





2 Review Exercise Exercise A, Question 3

Question:



A body of mass 5 kg is held in equilibrium under gravity by two inextensible light ropes. One rope is horizontal, the other is at an angle α to the horizontal, as shown. The tension in the rope inclined at α to the horizontal is 72 N. Find

a the angle α , giving your answer to the nearest degree,

b the tension T in the horizontal rope, giving your answer to the nearest N.



Alternative solution using triangle of forces



b

$$\cos \alpha = \frac{T}{72}$$

$$\therefore T = 72 \cos \alpha$$

$$= 53 \,\mathrm{N}(2 \,\mathrm{s.f.})$$

2 Review Exercise Exercise A, Question 4

Question:



A particle *P* is attached to one end of a light inextensible string. The other end of the string is attached to a fixed point *O*. A horizontal force of magnitude 12 N is applied to *P*. The particle *P* is in equilibrium with the string taut and *OP* making an angle of 20° with the downward vertical, as shown. Find

a the tension in the string,

b the weight of *P*.



2 Review Exercise Exercise A, Question 5

Question:



A particle *P* is held in equilibrium under gravity by two light, inextensible strings. One string is inclined at an angle of 60° to the horizontal and has a tension of 40 N. The other string is inclined at an angle of 40° to the horizontal and has a tension of *T* newtons, as shown. Find, to three significant figures,

a the value of T,

b the weight of *P*.





2 Review Exercise Exercise A, Question 6

Question:



A smooth bead *B* is threaded on a light inextensible string. The ends of the string are attached to two fixed points *A* and *C* on the same horizontal level. The bead is held in equilibrium by a horizontal force of magnitude 6 N acting parallel to *AC*. The bead *B* is $_{3}$

vertically below C and $\angle BAC = \alpha$, as shown in the diagram. Given that $\tan \alpha = \frac{3}{4}$, find

a the tension in the string,

b the weight of the bead.



2 Review Exercise Exercise A, Question 7

Question:



Two small rings, A and B, each of mass 2m, are threaded on a rough horizontal pole. The coefficient of friction between each ring and the pole is μ . The rings are attached to the ends of a light inextensible string. A smooth ring C, of mass 3m, is threaded on the string and hangs in equilibrium below the pole. The rings A and B are in limiting equilibrium on the pole, with

 $\angle BAC = \angle ABC = \theta$, where $\tan \theta = \frac{3}{4}$, as shown in the diagram.

a Show that the tension in the string is $\frac{5}{2}mg$.

b Find the value of μ .



2 Review Exercise Exercise A, Question 8

Question:

30 C

A particle of weight *W* newtons is attached at *C* to the ends of two light inextensible strings *AC* and *BC*. The other ends of the strings are attached to two fixed points *A* and *B* on a horizontal ceiling. The particle hangs in equilibrium with *AC* and *BC* inclined to the horizontal at 30° and 60° respectively, as shown. Given the tension in *AC* is 50 N, calculate

a the tension in BC, to three significant figures,

b the value of *W*.



2 Review Exercise Exercise A, Question 9

Question:



A small box of mass 20 kg rests on a rough horizontal floor. The coefficient of friction between the box and the floor is 0.25. A light inextensible rope is tied to the box and pulled with a force of magnitude P newtons at 14° to the horizontal as shown in the diagram. Given that the box is on the point of sliding, find the value of P, giving your answer to 1 decimal place.

Solution:



2 Review Exercise Exercise A, Question 10

Question:



A smooth plane is inclined at an angle 10° to the horizontal. A particle *P* of mass 2 kg is held in equilibrium on the plane by a horizontal force of magnitude *F* newtons, as shown.

Find, to three significant figures,

a the normal reaction exerted by the plane on P.

b the value of *F*.



2 Review Exercise Exercise A, Question 11

Question:

XN. 20°

A particle *P* of mass 2.5 kg rests in equilibrium on a rough plane under the action of a force of magnitude *X* newtons acting up a line of greatest slope of the plane, as shown in the diagram. The plane is inclined at 20° to the horizontal. The coefficient of friction between *P* and the plane is 0.4. The particle is in limiting equilibrium and is on the point of moving up the plane. Calculate

a the normal reaction of the plane on P,

b the value of X.

The force of magnitude X newtons is now removed.

c Show that *P* remains in equilibrium on the plane.

a

Draw a diagram showing the normal reaction R N, the friction force down the plane F N and the weight 2 mg. 2.5 g 120° $R(\bar{\mathbb{N}})$ Resolve perpendicular to the plane first, as R is the only unknown in the resulting R $-2.5 g \cos 20^{\circ} = 0$ equation. $\therefore R = 2.5 g \cos 20^{\circ}$ 23(2s.f.) = Give your answer to 2 s.f. as you used The normal reaction is 23 N. g = 9.8 in your calculation. **b** R(↗) Resolve parallel to the plane to find X in $X - F - 2.5 g \sin 20^{\circ} = 0$ terms of F. $\therefore X = F + 2.5 g \sin 20^{\circ}$ As friction is limiting, F μR Use limiting friction and the value of R from part a to find the force F N. i.e. F = 0.4 Rand as R = 23.0, F = 9.21substitute into equation * Then X may be calculated. Again give XX = 17.6to 2 s.f. =18(2s.f.)

c The force X is removed and the friction force will now act up the plane.



2 Review Exercise Exercise A, Question 12

Question:



A parcel of weight 10 N lies on a rough plane inclined at an angle of 30° to the horizontal. A horizontal force of magnitude *P* newtons acts on the parcel, as shown. The parcel is in equilibrium and on the point of slipping up the plane. The normal reaction of the plane on the parcel is 18 N. The coefficient of friction between the parcel and the plane is μ . Find

a the value of *P*,

b the value of μ .

The horizontal force is removed.

c Determine whether or not the parcel moves.



The friction now acts up the plane and the normal reaction will be different. Let the normal reaction be *R*. $R(\nearrow)$





2 Review Exercise Exercise A, Question 13

Question:



A small ring of mass 0.25 kg is threaded on a fixed rough horizontal rod. The ring is pulled upwards by a light string which makes an angle 40° with the horizontal, as shown. The string and the rod are in the same vertical plane. The tension in the string is 1.2 N and the coefficient of friction between the ring and the rod is μ . Given that the ring is in limiting equilibrium, find

 ${\bf a}$ the normal reaction between the ring and the rod,

b the value of μ .

Solution:



2 Review Exercise Exercise A, Question 14

Question:



The diagram shows a boat B of mass 400 kg held at rest on a slipway by a rope. The boat is modelled as a particle and the slipway as a rough plane inclined at 15° to the horizontal. The coefficient of friction between B and the slipway is 0.2. The rope is modelled as a light, inextensible string, parallel to a line of greatest slope of the plane. The boat is in equilibrium and on the point of sliding down the slipway.

a Calculate the tension in the rope.

The boat is 50 m from the bottom of the slipway. The rope is detached from the boat and the boat slides down the slipway.

 ${\bf b}$ Calculate the time taken for the boat to slide to the bottom of the slipway.



Let T be the tension, F the friction and R the normal reaction. **a** $R(\stackrel{r}{\searrow})$





Time to slide down is 12.5 s.

2 Review Exercise Exercise A, Question 15

Question:



A heavy package is held in equilibrium on a slope by a rope. The package is attached to one end of the rope, the other end being held by a man standing at the top of the slope. The package is modelled as a particle of mass 20 kg. The slope is modelled as a rough plane inclined at 60° to the horizontal and the rope as a light inextensible string. The string is assumed to be parallel to a line of greatest slope of the plane, as shown in the diagram. At the contact between the package and the slope, the coefficient of friction is 0.4.

a Find the minimum tension in the rope for the package to stay in equilibrium on the slope.

The man now pulls the package up the slope. Given that the package moves at constant speed,

b find the tension in the rope.

c State how you have used, in your answer to part b, the fact that the package moves

i up the slope,

ii at constant speed.



Let T be the minimum tension, F the force of friction and R the normal reaction.

 $R(\nearrow)$ $R - 20 g \cos 60^{\circ} = 0$ $\therefore R = 20 g \cos 60^{\circ}$ = 98As the friction is limiting, $F = \mu R$.



 $\therefore F = 0.4 \times 98$ = 39.2 When the tension is a minimum the friction is limiting.

R(≦)



 $R(\nearrow)$ R = 98 as before


2 Review Exercise Exercise A, Question 16

Question:

4 Nα

A particle *P* of mass 0.5 kg is on a rough plane inclined at an angle α to the horizontal, where $\tan \alpha = \frac{3}{4}$. The particle is held at rest on the plane by the action of a force of magnitude 4 N acting up the plane in a direction parallel to a line of greatest slope of the plane, as shown. The particle is on the point of slipping up the plane.

a Find the coefficient of friction between P and the plane.

The force of magnitude 4 N is removed.

b Find the acceleration of *P* down the plane.



a Let the normal reaction be R N, and the friction be F N.

$$R(\swarrow)$$

$$R - 0.5 g \cos \alpha = 0$$

$$\therefore R = 0.5 g \times \frac{4}{5}$$

$$= 3.92 (3 \text{ s.f.})$$
Find *R* by resolving perpendicular to the plane.

$$R(\checkmark)$$

$$4 - F - 0.5 g \sin \alpha = 0$$

$$\therefore F = 4 - 0.5 g \sin \alpha$$

$$= 1.06 (3 \text{ s.f.})$$
Use $F = \mu R$, as the friction is limiting.

$$\therefore \mu = \frac{F}{R} = \frac{1.06}{3.92}$$
As the particle is on the point of slipping,
 $F = \mu R$.

b The force of 4 N is removed.

 $\therefore \mu = 0.270 (3 \text{ s.f.})$

The friction will now act up the plane





:. Acceleration is 3.76 m s⁻² (3 s.f.) down the plane or 3.8 m s⁻² (2 s.f.).

2 Review Exercise Exercise A, Question 17

Question:



A box of mass 30 kg is being pulled along rough horizontal ground at a constant speed using a rope. The rope makes an angle of 20° with the ground, as shown. The coefficient of friction between the box and the ground is 0.4. The box is modelled as a particle and the rope as a light, inextensible string. The tension in the rope is *P* newtons.

a Find the value of *P*.

The tension in the rope is now increased to 150 N.

b Find the acceleration of the box.





2 Review Exercise Exercise A, Question 18

Question:



A uniform rod AB has length 8 m and mass 12 kg. A particle of mass 8 kg is attached to the rod at B. The rod is supported at a point C and is in equilibrium in a horizontal position, as shown.

Find the length of AC.

Solution:



2 Review Exercise Exercise A, Question 19

Question:



A uniform beam AB has mass 12 kg and length 3 m. The beam rests in equilibrium in a horizontal position, resting on two smooth supports. One support is at end A, the other at a point C on the beam, where BC = 1 m, as shown in the diagram. The beam is modelled as a uniform rod.

a Find the reaction on the beam at *C*.

A woman of mass 48 kg stands on the beam at the point D. The beam remains in equilibrium. The reactions on the beam at A and C are now equal.

b Find the distance *AD*.



As the beam is uniform the weight acts through G, the mid-point of AB. Let the reaction at A be S N and the reaction at C be R N.



Let the distance AD be x m and let the reactions at A and C be R N.

R(1)



2 Review Exercise Exercise A, Question 20

Question:



Two men, Eric and Fred, set out to carry a water container across a desert, using a long uniform pole. The length of the pole is 4 m and its mass is 5 kg. The ends of the pole rest in equilibrium on the shoulders of the two men, with the pole horizontal. The water container has mass 16 kg and is suspended from the pole by means of a light rope, which is short enough to prevent the container reaching the ground, as shown. Eric has a sprained ankle, so Fred fixes the rope in such a way that the vertical force on his shoulder is twice as great as the vertical force on Eric's shoulder.

a Find the vertical force on Eric's shoulder.

 \mathbf{b} Find the distance from the centre of the pole to the point at which the rope is fixed.

Solution:



Let the force on Eric's shoulder be R and the force on Fred's shoulder be 2R.



Take moments about Fred's shoulder. Let distance from centre of pole to point

at which says is fine 1



2 Review Exercise Exercise A, Question 21

Question:



A uniform rod AB has length 1.5 m and mass 8 kg. A particle of mass m kg is attached to the rod at B. The rod is supported at the point C, where AC = 0.9 m, and the system is in equilibrium with AB horizontal, as shown.

a Show that m = 2.

A particle of mass 5 kg is now attached to the rod at A and the support is moved from C to a point D of the rod. The system, including both particles, is again in equilibrium with AB horizontal.

b Find the distance *AD*.



2 Review Exercise Exercise A, Question 22

Question:



A uniform steel girder *AB*, of mass 150 kg and length 10 m, rests horizontally on two supports at *A* and *B*. A man of mass 90 kg stands on the girder at the point *P*, where AP = 2 m, as shown. By modelling the girder as a uniform rod and the man as a particle.

a find the magnitude of the reaction at *B*.



The support *B* is moved to a point *Y* on the girder, where BY = x metres, as shown. The man remains on the girder at *P*. The magnitudes of the reactions at the two supports are now equal.

Find

b the magnitude of the reaction at each support,

c the value of x.



Let G be the mid-point of AB. As the girder is uniform the weight acts through G. Let the reaction at B be R N.



2 Review Exercise Exercise A, Question 23

Question:

A footbridge across a stream consists of a uniform horizontal plank *AB* of length 5 m and mass 140 kg, supported at the ends *A* and *B*.

A man of mass 100 kg is standing at a point C on the footbridge. Given that the magnitude of the force exerted by the support at A is twice the magnitude of the force exerted by the support at B, calculate

a the magnitude, in N, of the force exerted by the support at B,

b the distance *AC*.

Solution:



2 Review Exercise Exercise A, Question 24

Question:

A non-uniform thin straight rod *AB* has length 3*d* and mass 5*m*. It is in equilibrium resting horizontally on supports at the points *X* and *Y*, where AX = XY = YB = d. A particle of mass 2*m* is attached to the rod at *B*. Given that the rod is on the point of tilting about *Y*, find the distance of the centre of mass of the rod from *A*.

Solution:



2 Review Exercise Exercise A, Question 25

Question:



A uniform rod *AB* has weight 70 N and length 3 m. It rests in a horizontal position on two smooth supports placed at *P* and *Q*, where AP = 0.5 m as shown in the diagram. The reaction on the rod at *P* has magnitude 20 N. Find

a the magnitude of the reaction on the rod at Q,

b the distance AQ.

Solution:



2 Review Exercise Exercise A, Question 26

Question:



A uniform plank *AB* has weight 120 N and length 3 m. The plank rests horizontally in equilibrium on two smooth supports *C* and *D*, where AC = 1 m and CD = x m, as shown. The reaction of the support on the plank at *D* has magnitude 80 N. Modelling the plank as a rod.

a show that x = 0.75.

A rock is now placed at B and the plank is on the point of tilting about D. Modelling the rock as a particle, find

b the weight of the rock,

c the magnitude of the reaction of the support on the plank at D.

 \boldsymbol{d} State how you have used the model of the rock as a particle.



When the plank is on the point of tilting about D, the reaction at C is zero.



d The weight of the rock acts precisely at B.

file://C:\Users\Buba\kaz\ouba\m1_rev2_a_26.html

2 Review Exercise Exercise A, Question 27

Question:



A seesaw in a playground consists of a beam AB of length 4 m which is supported by a smooth pivot at its centre C. Jill has mass 25 kg and sits on the end A. David has mass 40 kg and sits at a distance x metres from C, as shown. The beam is initially modelled as a uniform rod. Using this model,

a find the value of x for which the seesaw can rest in equilibrium in a horizontal position.

 ${\bf b}$ State what is implied by the modelling assumptions that the beam is uniform.

David realises that the beam is not uniform as he finds he must sit at a distance 1.4 m from C for the seesaw to rest horizontally in equilibrium. The beam is now modelled as a non-uniform rod of mass 15 kg. Using this model,

c find the distance of the centre of mass of the beam from C.





 \therefore Distance of centre of mass from C is 0.4 m.

2 Review Exercise Exercise A, Question 28

Question:



A steel girder *AB* has weight 210 N. It is held in equilibrium in a horizontal position by two vertical cables. One cable is attached to the end *A*. The other cable is attached to the point *C* on the girder, where AC = 90 cm, as shown. The girder is modelled as a uniform rod, and the cables as light inextensible strings.

Given that the tension in the cable at C is twice the tension in the cable at A, find

a the tension in the cable at *A*,

b show that AB = 120 cm.

A small load of weight W newtons is attached to the girder at B. The load is modelled as a particle. The girder remains in equilibrium in a horizontal position. The tension in the cable at C is now three times the tension in the cable at A.

c Find the value of *W*.



[©] Pearson Education Ltd 2008

2 Review Exercise Exercise A, Question 29

Question:



A uniform rod *AB* has mass 8 kg and length 4 m. A particle of mass 4 kg is attached to the rod at *A* and a particle of mass *M* kg is attached to the rod at *B*. The rod is supported at the point *C*, where AC = 1.5 m, and rests in equilibrium in a horizontal position, as shown in the diagram.

a Find the value of *M*.

 ${\bf b}$ State how you used the information that the rod is uniform.

Solution:



2 Review Exercise Exercise A, Question 30

Question:



A uniform plank *ABC*, of length 12 m and mass 30 kg, is supported in a horizontal position at the points *A* and *B*, where AB = 8 m and BC = 4 m, as shown in the diagram. A woman of mass 60 kg stands on the plank at a distance of 2 m from *A*, and a rock of mass *M* kg is placed on the plank at the end *C*. The plank remains in equilibrium. The plank is modelled as a uniform rod, and the woman and the rock as particles.

Given that the forces exerted by the supports on the plank at A and B are equal in magnitude,

a find **i** the value of *M*, **ii** the magnitude of the force exerted by the support at *A* on the plank.

 ${\bf b}$ State how you used the modelling assumption that the rock is a particle.



2 Review Exercise Exercise A, Question 31

Question:



A light rod *AB* has length 10 m. It is suspended by two light vertical cables attached to the rod at the points *C* and *D*, where AC = 2 m, CD = 4 m, and DB = 4 m, as shown in the diagram. A load of weight 60 N is attached to the rod at *A* and a load of weight *X*N is attached to the rod at *B*. The rod is hanging in equilibrium in a horizontal position. Find, in terms of *X*,

a the tension in the cable at *C*,

b the tension in the cable at *D*,

c Hence show that $15 \le X \le 90$.

If the tension in either cable exceeds 120 N that cable breaks.

d Find the maximum possible value of X.



The maximum possible value of X is 75.

2 Review Exercise Exercise A, Question 32

Question:



A large uniform plank of wood of length 8 m and mass 30 kg is held in equilibrium by two small steel rollers A and B, ready to be pushed into a saw-mill. The centres of the rollers are 50 cm apart. One end of the plank presses against roller A from underneath, and the plank rests on top of roller B, as shown in the diagram. The rollers are adjusted so that the plank remains horizontal and the force exerted on the plank by each roller is vertical.

a Suggest a suitable model for the plank to determine the forces exerted by the rollers.

b Find the magnitude of the force exerted on the plank by the roller at B.

c Find the magnitude of the force exerted on the plank by the roller at A.

Solution:



2 Review Exercise Exercise A, Question 33

Question:



A plank of wood *AB* has length 5.4 m. It lies on a horizontal platform, with 1.2 m projecting over the edge, as shown in the diagram. When a girl of mass 50 kg stands at the point *C* on the plank, where BC = 0.3 m, the plank is on the point of tilting. By modelling the plank as a uniform rod and the girl as a particle,

a find the mass of the plank.

The girl places a rock on the end of the plank at A. By modelling the rock also as a particle,

b find, to two significant figures, the smallest mass of the rock which will enable the girl to stand on the plank at *B* without tilting.

c State briefly how you have used the modelling assumption that

i the plank is uniform,

ii the rock is a particle.



c i Plank is uniform \rightarrow weight acts through the mid-point.

ii Rock is a particle \rightarrow mass of rock acts through the end-point A.

2 Review Exercise Exercise A, Question 34

Question:

Three forces F_1 , F_2 and F_3 act on a particle.

 $F_1 = \ (\ -3\, \vec{z} + 7j \) \ \mathrm{N} \ , \ F_2 = \ (\ \vec{z} - j \) \ \mathrm{N} \ , \ F_3 = \ (\ p\vec{z} + qj \) \ \mathrm{N}$

a Given that the particle is in equilibrium, determine the value of p and the value of q.

The resultant of the forces F_1 and F_2 is **R**.

b Calculate, in N, the magnitude of **R**.

c Calculate to the nearest degree, the angle between the line of action of R and the vector $\boldsymbol{j}.$



2 Review Exercise Exercise A, Question 35

Question:



Two forces **P** and **Q**, act on a particle. The force **P** has magnitude 5 N and the force **Q** has magnitude 3 N. The angle between the directions of **P** and **Q** is 40° , as shown in the diagram. The resultant of **P** and **Q** is **F**.

 \mathbf{a} Find, to three significant figures, the magnitude of \mathbf{F} .

b Find, in degrees to one decimal place, the angle between the directions of ${\bf F}$ and ${\bf P}.$

a Draw a vector triangle:-


2 Review Exercise Exercise A, Question 36

Question:

Two forces **P** and **Q** act on a particle. The force **P** has magnitude 7 N and acts due north. The resultant of **P** and **Q** is a force of magnitude 10 N acting in a direction with bearing 120° . Find

a the magnitude of **Q**,

 \mathbf{b} the direction of \mathbf{Q} , giving your answer as a bearing.





2 Review Exercise Exercise A, Question 37

Question:

Two forces $F_1 = (2\vec{x} + 3j)$ N and $F_2 = (\lambda\vec{x} + \mu j)$ N, where λ and μ are scalars, act on a particle. The resultant of the two forces is **R**, where **R** is parallel to the vector $\vec{x} + 2j$.

a Find, to the nearest degree, the acute angle between the line of action of R and the vector i.

b Show that $2\lambda - \mu + 1 = 0$.

Given that the direction of F_2 is parallel to **j**,

 \boldsymbol{c} find, to three significant figures, the magnitude of $\boldsymbol{R}.$



2 Review Exercise Exercise A, Question 38

Question:

A force **R** acts on a particle, where $R = (7\vec{z} + 16j)$ N.

Calculate

 \mathbf{a} the magnitude of \mathbf{R} , giving your answers to one decimal place,

 \mathbf{b} the angle between the line of action of \mathbf{R} and \mathbf{i} , giving your answer to the nearest degree.

The force **R** is the resultant of two forces **P** and **Q**. The line of action of P is parallel to the vector $(\vec{x} + 4j)$ and the line of action of **Q** is parallel to the vector $(\vec{x} + j)$.

c Determine the forces P and Q expressing each in terms of i and j.



[©] Pearson Education Ltd 2008

2 Review Exercise Exercise A, Question 39

Question:

A particle *P* moves in a straight line with constant velocity. Initially *P* is at the point A with position vector $(2\vec{x} - j)$ m relative to a fixed origin *O*, and 2s later it is at the point *B* with position vector $(6\vec{x} + j)$ m.

a Find the velocity of *P*.

b Find, in degrees to one decimal place, the size of the angle between the direction of motion of *P* and the vector **i**.

Three second after it passes B the particle P reaches the point C.

c Find, in metres to one decimal place, the distance OC.



2 Review Exercise Exercise A, Question 40

Question:

A particle P is moving with constant velocity $(5t - 3j) \text{ ms}^{-1}$. At time t = 0, its position vector, with respect to a fixed origin O, is $(-2\mathcal{I} + j)$ m. Find, to three significant figures.

a the speed of P,

b the distance of *P* from *O* when t = 2s.

Solution:

a The velocity u = 5i - 3jThe speed

$ u = \sqrt{5^2 + (-3)^2}$ $ u = \sqrt{34}$ $= 5.83 \text{ M s}^{-1}$	Use the formula for magnitude of a vector.
b The displacement from $t = 0$ to $t = 2$ is $(5i - 3j) \times 2$	Find the displacement between $t = 0$ and $t = 2$, using velocity × time.
The position vector of P at $t = 2$ is $(-2\mathbf{i} + \mathbf{j}) + (10\mathbf{i} - 6\mathbf{j})$	Use the vector law of addition.
= 8 i − 5 j \therefore The distance of <i>P</i> from <i>O</i> is	
$ \sqrt{8^2 + (-5)^2} = \sqrt{64 + 25} = \sqrt{89} $	Use the formula for the magnitude of a vector.

© Pearson Education Ltd 2008

=

9.43

2 Review Exercise Exercise A, Question 41

Question:

A boat *B* is moving with constant velocity. At noon, *B* is at the point with position vector $(3\vec{x} - 4j)$ km with respect to a fixed origin *O*. At 1430 on the same day, *B* is at the point with position vector $(8\vec{x} + 11j)$ km.

a Find the velocity of *B*, giving your answer in the form $p\vec{z} + qj$.

At time t hours after noon, the position vector of B is **b** km.

b Find, in terms of *t*, an expression for **b**.

Another boat C is also moving with constant velocity. The position vector of C, \mathbf{c} km, at time t hours after noon, is given by

 $c = (-9\vec{z} + 20j) + t(6\vec{z} + -j)$, where λ is a constant.

Given that C intercepts B,

c find the value of λ ,

d show that, before C intercepts B, the boats are moving with the same speed.

a Displacement between noon and 14.30 is



2 Review Exercise Exercise A, Question 42

Question:

A particle *P* of mass 2 kg is moving under the action of a constant force **F** newtons. When t = 0, *P* has velocity $(3\ell + 2j) \text{ m s}^{-1}$ and at time t = 4s, *P* has velocity $(15\ell - 4j) \text{ m s}^{-1}$. Find

a the acceleration of *P* in terms of **i** and **j**,

b the magnitude of **F**,

c the velocity of *P* at time t = 6 s.

Solution:

a As the force is constant, the acceleration is constant. Given $u = 3\mathbf{i} + 2\mathbf{j}$, $v = 15\mathbf{i} - 4\mathbf{j}$ and t = 4, to find *a* use



2 Review Exercise Exercise A, Question 43

Question:

The horizontal unit vectors *i* and *j* are due east and due north respectively.

A ship *S* is moving with constant velocity $(-2.5\mathbf{i} + 6\mathbf{j})\text{kmh}^{-1}$. At time 1200, the position vector of *S* relative to a fixed origin *O* is $(16\mathbf{i} + 5\mathbf{j})\text{km}$. Find

a the speed of *S*,

b the bearing on which *S* is moving.

The ship is heading directly towards a submerged rock R. A radar tracking station calculates that, if S continues on the same course with the same speed, it will hit R at the time 1500.

c Find the position vector of *R*.

The tracking station warns the ship's captain of the situation. The captain maintains *S* on its course with the same speed until the time is 1400. He then changes course so that *S* moves due north at a constant speed of 5 km h^{-1} . Assuming that *S* continues to move with this new constant vector, find

d an expression for the position vector of the ship *t* hours after 1400.

e The time when *S* will be due east of *R*.

f The distance of *S* from *R* at the time 1500.





2 Review Exercise Exercise A, Question 44

Question:

The horizontal unit vectors *i* and *j* are due east and due north respectively.

A model boat *A* moves on a lake with constant velocity $(-\mathbf{i} + 6\mathbf{j})$ m s⁻¹. At time t = 0, *A* is at the point with position vector $(2\mathbf{i} - 10\mathbf{j})$ m. Find

a the speed of A,

b the direction in which A is moving, giving your answer as a bearing.

At time t = 0, a second boat *B* is at the point with position vector $(-26\mathbf{i} + 4\mathbf{j})\mathbf{m}$.

Given that the velocity of *B* is $(3\mathbf{i} + 4\mathbf{j})$ m s⁻¹,

c show that *A* and *B* will collide at a point *P* and find the position vector of *P*.

Given instead that *B* has speed 8 m s^{-1} and moves in the direction of the vector $(3\mathbf{i} + 4\mathbf{j})$.

d find the distance of *B* from *P* when t = 7s.





Edexcel AS and A Level Modular Mathematics

2 Review Exercise Exercise A, Question 45

Question:

The horizontal unit vectors *i* and *j* are due east and due north respectively.

At time t = 0, a football player kicks a ball from the point *A* with position vector $(2\vec{x} + j)$ m on a horizontal football field. The motion of the ball is modelled as that of a particle moving horizontally with constant velocity $(5\vec{x} + 8j)$ m s⁻¹. Find

a the speed of the ball,

b the position vector of the ball after *t* seconds.

The point *B* on the field has position vector $(10\vec{z} + 7j)$ m.

c Find the time when the ball is due north of *B*.

At time t = 0, another player starts running due north from *B* and moves with constant speed $v \text{ m s}^{-1}$. Given that he intercepts the ball,

d find the value of v.

e State one physical factor, other than air resistance, which would be needed in a refinement of the model of the ball's motion to make the model more realistic.

a

velocity =
$$5i + 8j \text{ m s}^{-1}$$

 \therefore speed = $\sqrt{5^2 + 8^2} \text{ m s}^{-1}$
= $\sqrt{89} \text{ m s}^{-1} = 9.43 \text{ m s}^{-1}$ (3 s.f.)

b After *t* seconds, position vector is

 $\mathbf{r} = 2\mathbf{i} + \mathbf{j} + t(5\mathbf{i} + 8\mathbf{j})$

c When the ball is due north of 10i + 7j

2+5t = 10 $\therefore 5t = 8 \Rightarrow t = 1.6 \text{ s}$

d At t = 1.6 ball is at $10\mathbf{i} + 13.8\mathbf{j}$ The second player moves from $10\mathbf{i} + 7\mathbf{j}$ to $10\mathbf{i} + 13.8\mathbf{j}$ in 1.6 s. His velocity is $6.8\mathbf{j} \div 1.6$ His speed is $6.8 \div 1.6 = 4.25$ m s⁻¹.

e Friction on the field – so velocity of ball not constant or vertical component of ball's motion

or

time for player to accelerate.

© Pearson Education Ltd 2008

Use speed = magnitude of velocity. Use formula for magnitude of vector.

Use displacement = velocity × time.

Equate the i component of the ball to 10.

Find the displacement of the second player. Use velocity = displacement ÷ time.

Any of these answers would be valid.

2 Review Exercise Exercise A, Question 46

Question:

A destroyer is moving due west at a constant speed of 10 km h^{-1} . It has radar on board which, at time t = 0, identifies a cruiser, 50 km due west and moving due north with a constant speed of 20 km h^{-1} . The unit vectors **i** and **j** are directed due east and north respectively, and the origin *O* is taken to be the initial position of the destroyer. Each vessel maintains its constant velocity.

a Write down the velocity of each vessel in vector form.

b Find the position vector of each vessel at time *t* hours.

c Show that the distance d km between the vessels at time t hours is given by

 $d^2 = 500t^2 - 1000t + 2500$

The radar on the cruiser detects vessels only up to a distance of 40 km. By finding the minimum value of d^2 , or otherwise,

d determine whether the destroyer will be detected by the cruiser's radar.



$$\mathbf{r}_{\text{destroyer}} = -10 t \mathbf{i} = \mathbf{d}$$

$$\mathbf{r}_{\text{rewiser}} = -50\mathbf{i} + 20 t \mathbf{j} = \mathbf{c}$$
Using $\mathbf{r}_{\text{new}} = \mathbf{r}_{\text{old}} + \mathbf{v}t$

c The vector $\overrightarrow{\text{CD}} = \mathbf{d} - \mathbf{c}$

$$= -10 t \mathbf{i} - (-50\mathbf{i} + 20 t \mathbf{j})$$

$$= (50 - 10 t) \mathbf{i} - 20 t \mathbf{j}$$
Using the triangle law, or vector subtraction.

$$\therefore |\overrightarrow{CD}| = \sqrt{(50 - 10 t)^2 + (-20 t)^2}$$

$$= \sqrt{2500 - 1000 t + 100 t^2 + 400 t^2}$$
Use Pythagoras' Theorem to find the magnitude of \overrightarrow{CD} .

d

$$d^{2} = 500(t^{2} - 2t + 5)$$

= 500((t-1)^{2} + 4)

The minimum value of d^2 is when t = 1 and

$$d^{2} = 500 \times 4$$
$$= 2000$$
$$\therefore d = \sqrt{2000}$$
$$= 44.72$$

The minimum value of d^2 can be found by completion of the square.

 \therefore As 44.72 > 40 cruiser will not be able to detect the destroyer.

An alternative method would be to attempt to solve $d^2 = 40^2$. This gives a quadratic with no real solutions.

2 Review Exercise Exercise A, Question 47

Question:

In this question the vectors **i** and **j** are horizontal unit vectors in the directions due east and due north respectively. Two boats A and B are moving with constant velocities. Boat A moves with velocity $9j \text{ km h}^{-1}$. Boat B moves with velocity $(3l + 5j) \text{ km h}^{-1}$.

a Find the bearings on which *B* is moving.

At noon A is at the point O and B is 10 km due west of O. At time t hours after noon, the position vectors of A and B relative to O are \mathbf{a} km and \mathbf{b} km respectively.

b Find expressions for **a** and **b** in terms of *t*, giving your answer in the form

 $p\mathbf{I} + q\mathbf{j}$

c Find the time when *B* is due south of *A*.

At time time t hours after noon, the distance between A and B is d km. By finding an expression for AB,

d show that $d^2 = 25t^2 - 60t + 100$.

At noon the boats are 10 km apart.

e Find the time after noon at which the boats are again 10 km apart.

a θ 31 Let the bearing on which B is moving be α . Then $\tan \alpha = \frac{3}{5}$ Use trigonometry to find α . The bearing is the angle measured clockwise from the North direction. $\therefore \alpha = \arctan(0.6)$ 30.96 : bearing is 031° (nearest degree) b using $\mathbf{r}_{new} = \mathbf{r}_{old} + \mathbf{v} t$. a = 0+9t j**b** = -10i + (3i + 5j)t= (3t - 10)i + 5tjWhen B is due south of Ac 3t - 10 = 0 $\therefore t = \frac{10}{3}$ As A has i component zero. = 3 h 20 min i.e. the time is 15.20 d $\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$ $= (3 t - 10)\mathbf{i} - 4 t\mathbf{j}$ ∴ $|\overline{AB}| = \sqrt{(3 t - 10)^2 + (-4 t)^2}$ = $\sqrt{(9 t^2 - 60 t + 100 + 16 t^2)}$ Using the 'triangle law' or vector subtraction. $d^2 = 25t^2 - 60t + 100$ Use Pythagoras to find the magnitude of AB. e When d = 10Form and solve a quadratic equation in t. $100 = 25t^2 - 60t + 100$ $\therefore 25 t^2 - 60 t = 0$ $\therefore 5t(5t-12) = 0$ $\therefore t = 0 \text{ or } t = \frac{12}{5}$ Express your answer as a time in hours and minutes. = 2.4 h So the boats are 10 km apart at 14.24.