

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 1

Question:

$$\frac{dy}{dx} = 2x$$


Solution:

$$\frac{dy}{dx} = 2x$$

$$\therefore y = \int 2x \, dx$$

$$\therefore y = x^2 + c \quad \text{where } c \text{ is constant}$$

Integrate and include the constant of integration.
Let the constant take values 1, 2, 3, 0, -1, -2 and draw solution curves.



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 2

Question:

$$\frac{dy}{dx} = y$$

Solution:

$$\frac{dy}{dx} = y$$

$$\therefore \int \frac{1}{y} dy = \int 1 dx$$


Separate the variables and integrate.
Include a constant of integration on one side of the equation.

$$\therefore \ln y = x + c \quad \text{where } c \text{ is constant}$$

$$y = e^{x+c}$$

$$= e^c \times e^x$$

$$y = Ae^x \quad \text{where } A \text{ is constant } (A = e^c)$$



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 3

Question:


$$\frac{dy}{dx} = x^2$$

Solution:

$$\frac{dy}{dx} = x^2$$

$$y = \int x^2 dx$$

$$y = \frac{x^3}{3} + c \quad \text{where } c \text{ is constant}$$



© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 4

Question:

$$\frac{dy}{dx} = \frac{1}{x}, x > 0$$

Solution:


$$\frac{dy}{dx} = \frac{1}{x}$$

$$\therefore y = \int \frac{1}{x} dx$$

$$= \ln x + c$$

$$= \ln x + \ln A$$

$$\therefore y = \ln Ax$$



© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 5

Question:

$$\frac{dy}{dx} = \frac{2y}{x}$$

Solution:

$$\frac{dy}{dx} = \frac{2y}{x}$$

$$\therefore \int \frac{1}{y} dy = \int \frac{2}{x} dx$$

Separate the variables and integrate.


$$\therefore \ln y = 2 \ln x + c$$

$$\therefore \ln y = \ln x^2 + \ln A$$

$$= \ln Ax^2$$

Express the constant of integration as $\ln A$ where A is constant and use laws of logs to simplify your answer.

$$\therefore y = Ax^2$$



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 6

Question:

$$\frac{dy}{dx} = \frac{x}{y}$$

Solution:


$$\frac{dy}{dx} = \frac{x}{y}$$

$$\therefore \int y \, dy = \int x \, dx$$

$$\therefore \frac{y^2}{2} = \frac{x^2}{2} + c$$

$$\text{or } y^2 - x^2 = 2c$$

$y^2 - x^2 = 0$ is a pair of straight lines.
These are $y = x$ and $y = -x$
 $y^2 - x^2 = 2c$, $c \neq 0$ is a hyperbola.



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 7

Question:

$$\frac{dy}{dx} = e^y$$

Solution:

$$\frac{dy}{dx} = e^y$$

$$\therefore \int \frac{1}{e^y} dy = \int 1 dx$$

To integrate $\frac{1}{e^y}$, express it as e^{-y} .


$$\therefore \int e^{-y} dy = \int 1 dx$$

$$\therefore -e^{-y} = x + c$$

$$\therefore -e^{-y} = -x - c$$

$$\therefore -y = \ln[-x - c]$$

$$y = -\ln[-x - c] \text{ or } \ln \frac{1}{(-x - c)}$$



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 8

Question:

$$\frac{dy}{dx} = \frac{y}{x(x+1)}, \quad x > 0$$

Solution:

$$\frac{dy}{dx} = \frac{y}{x(x+1)}$$


$$\therefore \int \frac{1}{y} dy = \int \frac{1}{x(x+1)} dx$$

Separate the variables,
then use partial fractions to
integrate the function of x .

$$\begin{aligned} \therefore \ln y &= \int \left(\frac{1}{x} - \frac{1}{(x+1)} \right) dx \\ &= \ln x - \ln(x+1) + c \end{aligned}$$

$$\begin{aligned} \therefore \ln y &= \ln \frac{x}{x+1} + \ln A \\ &= \ln \frac{Ax}{x+1} \end{aligned}$$

$$\therefore y = \frac{Ax}{x+1} \quad x > 0$$



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 9


Question:

$$\frac{dy}{dx} = \cos x$$

Solution:

$$\frac{dy}{dx} = \cos x$$

$$\therefore y = \sin x + c$$



© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 10

Question:

$$\frac{dy}{dx} = y \cot x, \quad 0 < x < \pi$$

Solution:

$$\frac{dy}{dx} = y \cot x \quad 0 < x < \pi$$


$$\therefore \int \frac{1}{y} dy = \int \frac{\cos x}{\sin x} dx$$

$$\therefore \ln|y| = \ln|\sin x| + \ln|A|$$

$$= \ln|A \sin x|$$

$$\therefore y = A \sin x$$

Express the constant of integration as $\ln|A|$ and combine logs to simplify your solution



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 11

Question:


$$\frac{dy}{dx} = \sec^2 t, \quad -\frac{\pi}{2} < t < \frac{\pi}{2}$$

Solution:

$$\frac{dx}{dt} = \sec^2 t \quad -\frac{\pi}{2} < t < \frac{\pi}{2}$$

$$\therefore x = \int \sec^2 t \, dt$$

$$\text{i.e. } x = \tan t + c \text{ for } -\frac{\pi}{2} < t < \frac{\pi}{2}$$



© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 12

Question:

$$\frac{dy}{dx} = x(1 - x), \quad 0 < x < 1$$

Solution:

$$\frac{dy}{dx} = x(1 - x)$$

$$\therefore \int \frac{1}{x(1-x)} dx = \int 1 dt$$

$$\therefore \int \left(\frac{1}{x} + \frac{1}{1-x} \right) dx = \int 1 dt$$

$$\therefore \ln x - \ln(1-x) = t + c$$


$$\therefore \ln \frac{x}{1-x} = t + c$$

$$\therefore \frac{x}{1-x} = e^{t+c} = Ae^t \quad \leftarrow \begin{array}{l} 0 < x < 1 \text{ implies that } A \text{ is} \\ \text{a positive constant.} \end{array}$$

$$\therefore x = Ae^t - xAe^t$$

$$\therefore x(1 + Ae^t) = Ae^t$$

$$x = \frac{Ae^t}{1 + Ae^t}$$



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 13

Question:

Given that a is an arbitrary constant, show that $y^2 = 4ax$ is the general solution of the differential equation $\frac{dy}{dx} = \frac{y}{2x}$.

- a** Sketch the members of the family of solution curves for which $a = \frac{1}{4}$, 1 and 4.
b Find also the particular solution, which passes through the point (1, 3), and add this curve to your diagram of solution curves.

Solution:

$$\frac{dy}{dx} = \frac{y}{2x}$$

$$\therefore \int \frac{1}{y} dy = \frac{1}{2} \int \frac{1}{x} dx$$


$$\therefore \ln y = \frac{1}{2} \ln x + c$$

$$\text{or } \ln y = \frac{1}{2} \ln x + \ln A$$

$$\therefore \ln y = \ln A \sqrt{x}$$

$$\text{i.e. } y = A \sqrt{x} \quad \text{or } y^2 = A^2 x \quad \text{or } y^2 = 4ax$$

- a** Sketch $y^2 = x$, $y^2 = 4x$ and $y^2 = 16x$



- b** $y^2 = 4ax$ passes through (1, 3)

$$\therefore 9 = 4a$$

$$\text{i.e. } a = \frac{9}{4} \text{ and } y^2 = 9x$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 14

Question:

Given that k is an arbitrary positive constant, show that $y^2 + kx^2 = 9k$ is the general solution of the differential equation $\frac{dy}{dx} = \frac{-xy}{9 - x^2} \quad |x| \leq 3$.

- a** Find the particular solution, which passes through the point $(2, 5)$.
- b** Sketch the family of solution curves for $k = \frac{1}{9}, \frac{4}{9}, 1$ and include your particular solution in the diagram.

Solution:

$$\frac{dy}{dx} = \frac{-xy}{9-x^2}$$

$$\therefore \int \frac{1}{y} dy = - \int \frac{x}{9-x^2} dx$$

$$\therefore \ln y = \frac{1}{2} \ln(9-x^2) + \ln A$$

$$\therefore 2 \ln y = \ln A^2 (9-x^2)$$

$$\therefore \ln y^2 = \ln A^2 (9-x^2)$$

$$\therefore y^2 = 9A^2 - A^2 x^2$$

$$\text{Let } A^2 = k$$

$$\text{Then } y^2 + kx^2 = 9k$$

The solution curves are all ellipses, except when $k = 1$ when the curve is a circle.


a If this curve passes through (2, 5) then

$$25 + 4k = 9k$$

$$\therefore 25 = 5k \rightarrow k = 5$$

$$\text{i.e. } y^2 + 5x^2 = 45$$

b When $y = 0$ $x = \pm 3$, when $x = 0$ $y = \pm \sqrt{9k}$



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 1

Question:

$$x \frac{dy}{dx} + y = \cos x$$

Solution:

$$x \frac{dy}{dx} + y = \cos x$$

$$\text{So } \frac{d}{dx}(xy) = \cos x$$

$$\begin{aligned} \therefore xy &= \int \cos x \, dx \\ &= \sin x + c \end{aligned}$$

Remember to add the constant of integration when you integrate – not at the end of the process.

$$\therefore y = \frac{1}{x} \sin x + \frac{c}{x}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 2

Question:

$$e^{-x} \frac{dy}{dx} - e^{-x} y = xe^x$$

Solution:

$$e^{-x} \frac{dy}{dx} - e^{-x} y = xe^x$$

$$\therefore \frac{d}{dx} (e^{-x} y) = xe^x$$

$$\therefore e^{-x} y = \int xe^x dx \quad \bullet \quad \text{Use integration by parts to integrate } xe^x.$$

$$= xe^x - \int e^x dx$$

$$= xe^x - e^x + c$$

$$\therefore y = xe^{2x} - e^{2x} + ce^x \quad \bullet \quad \text{Multiply integration by } e^x.$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 3

Question:

$$\sin x \frac{dy}{dx} + y \cos x = 3$$

Solution:

$$\sin x \frac{dy}{dx} + y \cos x = 3$$

$$\therefore \frac{d}{dx}(y \sin x) = 3$$

$$\therefore y \sin x = \int 3 \, dx$$

$$\therefore y \sin x = 3x + c$$

$$\begin{aligned}\therefore y &= \frac{3x}{\sin x} + \frac{c}{\sin x} \\ &= 3x \operatorname{cosec} x + c \operatorname{cosec} x\end{aligned}$$

© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 4

Question:

$$\frac{1}{x} \frac{dy}{dx} - \frac{1}{x^2} y = e^x$$

Solution:

$$\frac{1}{x} \frac{dy}{dx} - \frac{1}{x^2} y = e^x$$

$$\therefore \frac{d}{dx} \left(\frac{1}{x} y \right) = e^x$$

$$\begin{aligned} \therefore \frac{1}{x} y &= \int e^x dx \\ &= e^x + c \end{aligned}$$

$$\therefore y = x e^x + c x$$

© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 5

Question:

$$x^2 e^y \frac{dy}{dx} + 2x e^y = x$$

Solution:

$$x^2 e^y \frac{dy}{dx} + 2x e^y = x$$

This time the left hand side is $\frac{d}{dx}(x^2 f(y))$ not just $\frac{d}{dx}(x^2 y)$.

$$\therefore \frac{d}{dx}(x^2 e^y) = x$$

$$\begin{aligned} \therefore x^2 e^y &= \int x \, dx \\ &= \frac{x^2}{2} + c \end{aligned}$$

$$\therefore e^y = \frac{1}{2} + \frac{c}{x^2}$$

$$\text{or } y = \ln \left[\frac{1}{2} + \frac{c}{x^2} \right]$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 6

Question:

$$4xy \frac{dy}{dx} + 2y^2 = x^2$$

Solution:

$$4xy \frac{dy}{dx} + 2y^2 = x^2$$

Again the left hand side of the equation can be written $\frac{d}{dx} (2x f(y))$.

$$\therefore \frac{d}{dx} (2xy^2) = x^2$$

$$\begin{aligned} \therefore 2xy^2 &= \int x^2 dx \\ &= \frac{1}{3}x^3 + c \end{aligned}$$

$$\therefore y^2 = \frac{1}{6}x^2 + \frac{c}{2x}$$

Divide both sides by $2x$.

$$\text{or } y = \pm \sqrt{\left(\frac{1}{6}x^2 + \frac{c}{2x}\right)}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 7

Question:

a Find the general solution of the differential equation

$$x^2 \frac{dy}{dx} + 2xy = 2x + 1.$$

b Find the three particular solutions which pass through the points with coordinates $(-\frac{1}{2}, 0)$, $(-\frac{1}{2}, 3)$ and $(-\frac{1}{2}, 19)$ respectively and sketch their solution curves for $x < 0$.

Solution:



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 8

Question:

a Find the general solution of the differential equation

$$\ln x \frac{dy}{dx} + \frac{y}{x} = \frac{1}{(x+1)(x+2)}, \quad x > 1.$$

b Find the specific solution which passes through the point (2, 2).

Solution:

$$\mathbf{a} \quad \ln x \frac{dy}{dx} + \frac{y}{x} = \frac{1}{(x+1)(x+2)}$$

$$\therefore \frac{d}{dx} (\ln x \times y) = \frac{1}{(x+1)(x+2)}$$

$$\therefore y \ln x = \int \frac{1}{(x+1)(x+2)} dx$$

You will need to use partial fractions to do the integration.

$$= \int \left(\frac{1}{x+1} - \frac{1}{x+2} \right) dx$$

$$= \ln(x+1) - \ln(x+2) + c$$

$$\therefore y = \frac{\ln(x+1) - \ln(x+2) + \ln A}{\ln x}$$

$$\therefore y = \frac{\ln A(x+1)}{\ln x(x+2)} \text{ is the general solution}$$

b When $x = 2, y = 2$

$$\therefore 2 = \frac{\ln \frac{3}{4} A}{\ln 2}$$

$$\therefore \ln \frac{3}{4} A = 2 \ln 2 = \ln 4$$

$$\therefore A = \frac{16}{3}$$

$$\text{So } y = \frac{\ln \frac{16(x+1)}{3(x+2)}}{\ln x}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 1

Question:

$$\frac{dy}{dx} + 2y = e^x$$

Solution:

$$\frac{dy}{dx} + 2y = e^x$$

The integrating factor is $e^{\int 2 dx} = e^{2x}$ •

Find the integral factor $e^{\int p dx}$ and multiply the differential equation by it to give an exact equation.

$$\therefore e^{2x} \frac{dy}{dx} + 2e^{2x} y = e^{3x}$$

$$\therefore \frac{d}{dx} (e^{2x} y) = e^{3x}$$

$$\begin{aligned} \therefore e^{2x} y &= \int e^{3x} dx \\ &= \frac{1}{3} e^{3x} + c \end{aligned}$$

$$\therefore y = \frac{1}{3} e^x + ce^{-2x}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 2

Question:

$$\frac{dy}{dx} + y \cot x = 1$$

Solution:

$$\frac{dy}{dx} + y \cot x = 1$$

The integrating factor is $e^{\int p dx} = e^{\int \cot x dx}$

$$= e^{\ln \sin x}$$

$$= \sin x$$

The integrating factor $e^{\ln f(x)}$ can be simplified to $f(x)$.

Multiply differential equation by $\sin x$.

$$\therefore \sin x \frac{dy}{dx} + y \cos x = \sin x$$

$$\therefore \frac{d}{dx} (y \sin x) = \sin x$$

$$\therefore y \sin x = \int \sin x dx$$

$$= -\cos x + c$$

$$\therefore y = -\cot x + c \operatorname{cosec} x$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 3

Question:

$$\frac{dy}{dx} + y \sin x = e^{\cos x}$$

Solution:

$$\frac{dy}{dx} + y \sin x = e^{\cos x}$$

The integrating factor is $e^{\int \sin x \, dx} = e^{-\cos x}$

$$\therefore e^{-\cos x} \frac{dy}{dx} + y \sin x e^{-\cos x} = 1$$

$$\therefore \frac{d}{dx} (ye^{-\cos x}) = 1$$

$$\therefore ye^{-\cos x} = x + c$$

$$\therefore y = xe^{\cos x} + ce^{\cos x}$$

© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 4

Question:

$$\frac{dy}{dx} - y = e^{2x}$$

Solution:

$$\frac{dy}{dx} - y = e^{2x}$$

The integrating factor is $e^{\int -1 \, dx} = e^{-x}$ •

Remember that $P(x) = -1$ and the minus sign is important.

$$\therefore e^{-x} \frac{dy}{dx} - ye^{-x} = e^{2x} \times e^{-x}$$

$$\therefore \frac{d}{dx} (ye^{-x}) = e^x$$

$$\therefore ye^{-x} = \int e^x \, dx$$

$$= e^x + c$$

$$\therefore y = e^{2x} + ce^x$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 5

Question:

$$\frac{dy}{dx} + y \tan x = x \cos x$$

Solution:

$$\frac{dy}{dx} + y \tan x = x \cos x$$

The integrating factor is $e^{\int \tan x \, dx} = e^{\ln \sec x}$
 $= \sec x$

Find the integrating factor and simplify $e^{\ln f(x)}$ to give $f(x)$.

$$\therefore \sec x \frac{dy}{dx} + y \sec x \tan x = x$$

$$\therefore \frac{d}{dx} (y \sec x) = x$$

$$\therefore y \sec x = \int x \, dx$$

$$= \frac{1}{2} x^2 + c$$

$$\therefore y = \left(\frac{1}{2} x^2 + c \right) \cos x$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 6

Question:

$$\frac{dy}{dx} + \frac{y}{x} = \frac{1}{x^2}$$

Solution:

$$\frac{dy}{dx} + \frac{y}{x} = \frac{1}{x^2}$$

The integrating factor is $e^{\int \frac{1}{x} dx} = e^{\ln x} = x$

$$\therefore x \frac{dy}{dx} + y = \frac{1}{x}$$

$$\therefore \frac{d}{dx}(xy) = \frac{1}{x}$$

$$\begin{aligned} \therefore xy &= \int \frac{1}{x} dx \\ &= \ln x + c \end{aligned}$$

$$\therefore y = \frac{1}{x} \ln x + \frac{c}{x}$$

© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 7

Question:

$$x^2 \frac{dy}{dx} - xy = \frac{x^3}{x+2} \quad x > -2$$

Solution:

$$x^2 \frac{dy}{dx} - xy = \frac{x^3}{x+2}$$

Divide by x^2 •

First divide the equation through by x^2 , to give the correct form of equation.

$$\therefore \frac{dy}{dx} - \frac{1}{x}y = \frac{x}{x+2}$$

The integrating factor is $e^{\int -\frac{1}{x} dx} = e^{-\ln x} = e^{\ln \frac{1}{x}} = \frac{1}{x}$

Multiply the new equation by $\frac{1}{x}$

$$\therefore \frac{1}{x} \frac{dy}{dx} - \frac{1}{x^2}y = \frac{1}{x+2}$$

$$\therefore \frac{d}{dx} \left(\frac{1}{x}y \right) = \frac{1}{x+2}$$

$$\therefore \frac{1}{x}y = \int \frac{1}{x+2} dx$$

$$= \ln(x+2) + c$$

$$\therefore y = x \ln(x+2) + cx$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 8

Question:

$$3x \frac{dy}{dx} + y = x$$

Solution:

$$3x \frac{dy}{dx} + y = x$$

$$\therefore \frac{dy}{dx} + \frac{1}{3x}y = \frac{1}{3} \quad *$$

First divide equation through by $3x$, to get an equation of the correct form.

The integrating factor is $e^{\int \frac{1}{3x} dx} = e^{\frac{1}{3} \ln x}$

$$= e^{\ln x^{\frac{1}{3}}} = x^{\frac{1}{3}}$$

Multiply equation $*$ by $x^{\frac{1}{3}}$

$$\therefore x^{\frac{1}{3}} \frac{dy}{dx} + \frac{1}{3} x^{-\frac{2}{3}} y = \frac{1}{3} x^{\frac{1}{3}}$$

$$\therefore \frac{d}{dx} (x^{\frac{1}{3}} y) = \frac{1}{3} x^{\frac{1}{3}}$$

$$\begin{aligned} \therefore x^{\frac{1}{3}} y &= \int \frac{1}{3} x^{\frac{1}{3}} dx \\ &= \frac{1}{4} x^{\frac{4}{3}} + c \end{aligned}$$

$$\therefore y = \frac{1}{4} x + c x^{-\frac{1}{3}}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 9

Question:

$$(x + 2) \frac{dy}{dx} - y = (x + 2)$$

Solution:

$$(x + 2) \frac{dy}{dx} - y = (x + 2)$$

$$\therefore \frac{dy}{dx} - \frac{1}{(x + 2)} y = 1 \quad *$$

Divide equation by $(x + 2)$ before finding integrating factor.

The integrating factor is $e^{\int \frac{-1}{(x+2)} dx} = e^{-\ln(x+2)} = e^{\ln \frac{1}{x+2}}$

$$= \frac{1}{x + 2}$$

Multiply differential equation * by integrating factor.

$$\therefore \frac{1}{(x + 2)} \frac{dy}{dx} - \frac{1}{(x + 2)^2} y = \frac{1}{(x + 2)}$$

$$\therefore \frac{d}{dx} \left[\frac{1}{(x + 2)} y \right] = \frac{1}{x + 2}$$

$$\therefore \frac{1}{(x + 2)} y = \int \frac{1}{x + 2} dx$$

$$= \ln(x + 2) + c$$

$$\therefore y = (x + 2) \ln(x + 2) + c(x + 2)$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 10

Question:

$$x \frac{dy}{dx} + 4y = \frac{e^x}{x^2}$$

Solution:

$$x \frac{dy}{dx} + 4y = \frac{e^x}{x^2}$$

Divide throughout by x

$$\text{Then } \frac{dy}{dx} + \frac{4}{x}y = \frac{e^x}{x^3} \quad *$$

The integrating factor is $e^{\int \frac{4}{x} dx} = e^{4 \ln x} = e^{\ln x^4} = x^4$

$$\therefore x^4 \frac{dy}{dx} + 4x^3 y = xe^x \quad [\text{having multiplied } * \text{ by } x^4]$$

Integrate xe^x using integration by parts.

$$\therefore \frac{d}{dx}(x^4 y) = xe^x$$

$$\therefore x^4 y = \int xe^x dx$$

$$= xe^x - \int e^x dx$$

$$= xe^x - e^x + c$$

$$\therefore y = \frac{1}{x^3} e^x - \frac{1}{x^4} e^x + \frac{c}{x^4}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 11

Question:

Find y in terms of x given that

$$x \frac{dy}{dx} + 2y = e^x \text{ and that } y = 1 \text{ when } x = 1.$$

Solution:

$$x \frac{dy}{dx} + 2y = e^x$$

Divide throughout by x

$$\text{Then } \frac{dy}{dx} + \frac{2}{x}y = \frac{1}{x}e^x \quad *$$

The integrating factor is $e^{\int \frac{2}{x} dx} = e^{2 \ln x} = e^{\ln x^2} = x^2$

Multiply equation $*$ by x^2

$$\text{Then } x^2 \frac{dy}{dx} + 2xy = xe^x$$

$$\therefore \frac{d}{dx}(x^2y) = xe^x$$

$$\begin{aligned} \therefore x^2y &= \int xe^x dx \\ &= xe^x - \int e^x dx \\ &= xe^x - e^x + c \end{aligned}$$

$$\therefore y = \frac{1}{x}e^x - \frac{1}{x^2}e^x + \frac{c}{x^2}$$

Given also that $y = 1$ when $x = 1$

$$\text{Then } 1 = e - e + c$$

$$\therefore c = 1$$

$$\therefore y = \frac{1}{x}e^x - \frac{1}{x^2}e^x + \frac{1}{x^2}$$

Solve the differential equation then use the boundary condition $y = 1$ when $x = 1$ to find the constant of integration.

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 12

Question:

Solve the differential equation, giving y in terms of x , where

$$x^3 \frac{dy}{dx} - x^2y = 1 \text{ and } y = 1 \text{ at } x = 1.$$

Solution:

$$x^3 \frac{dy}{dx} - x^2y = 1$$

Divide throughout by x^3

$$\frac{dy}{dx} - \frac{1}{x}y = \frac{1}{x^3} \quad *$$

The integrating factor is $e^{-\int \frac{1}{x} dx} = e^{-\ln x} = e^{\ln \frac{1}{x}} = \frac{1}{x}$

Multiply equation $*$ by $\frac{1}{x}$

$$\text{Then } \frac{1}{x} \frac{dy}{dx} - \frac{1}{x^2}y = \frac{1}{x^4}$$

$$\therefore \frac{d}{dx} \left(\frac{1}{x}y \right) = \frac{1}{x^4}$$

$$\therefore \frac{1}{x}y = \int \frac{1}{x^4} dx$$

$$= \int x^{-4} dx$$

$$= -\frac{1}{3}x^{-3} + c$$

$$\therefore y = -\frac{1}{3}x^{-2} + cx$$

$$\text{So } y = -\frac{1}{3x^2} + cx$$

But $y = 1$, when $x = 1$

$$\therefore 1 = -\frac{1}{3} + c$$

$$\therefore c = \frac{4}{3}$$

$$\therefore y = -\frac{1}{3x^2} + \frac{4x}{3}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 13

Question:

Find the general solution of the differential equation

$$\left(x + \frac{1}{x}\right) \frac{dy}{dx} + 2y = 2(x^2 + 1)^2,$$

giving y in terms of x .

Find the particular solution which satisfies the condition that $y = 1$ at $x = 1$.

Solution:

$$\left(x + \frac{1}{x}\right) \frac{dy}{dx} + 2y = 2(x^2 + 1)^2$$

Divide equation by $\left(x + \frac{1}{x}\right)$.

$$\therefore \frac{dy}{dx} + \frac{2}{\left(x + \frac{1}{x}\right)} y = \frac{2(x^2 + 1)^2}{\left(x + \frac{1}{x}\right)}$$

$$\text{i.e. } \frac{dy}{dx} + \frac{2x}{x^2 + 1} \times y = 2x(x^2 + 1) \quad *$$

The integrating factor is $e^{\int \frac{2x}{x^2+1} dx} = e^{\ln(x^2+1)} = (x^2 + 1)$

Multiply $*$ by $(x^2 + 1)$

$$\text{Then } (x^2 + 1) \frac{dy}{dx} + 2xy = 2x(x^2 + 1)^2$$

$$\therefore \frac{d}{dx} [(x^2 + 1)y] = 2x(x^2 + 1)^2$$

$$\begin{aligned} \therefore y(x^2 + 1) &= \int 2x(x^2 + 1)^2 dx \\ &= \frac{1}{3}(x^2 + 1)^3 + c \end{aligned}$$

$$\therefore y = \frac{1}{3}(x^2 + 1)^2 + \frac{c}{(x^2 + 1)}$$

But $y = 1$, when $x = 1$

$$\therefore 1 = \frac{1}{3} \times 4 + \frac{1}{2}c$$

$$\therefore c = -\frac{2}{3}$$

$$\therefore y = \frac{1}{3}(x^2 + 1)^2 - \frac{2}{3(x^2 + 1)}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 14

Question:

Find the general solution of the differential equation

$$\cos x \frac{dy}{dx} + y = 1, \quad -\frac{\pi}{2} < x < \frac{\pi}{2}.$$

Find the particular solution which satisfies the condition that $y = 2$ at $x = 0$.

Solution:

$$\cos x \frac{dy}{dx} + y = 1$$

Divide throughout by $\cos x$

$$\therefore \frac{dy}{dx} + \sec x y = \sec x$$

The integrating factor is $e^{\int \sec x \, dx} = e^{\ln(\sec x + \tan x)}$
 $= \sec x + \tan x$

$$\int \sec x \, dx = \ln(\sec x + \tan x)$$

$$\therefore (\sec x + \tan x) \frac{dy}{dx} + (\sec^2 x + \sec x \tan x) y = \sec^2 x + \sec x \tan x$$

$$\therefore \frac{d}{dx} [(\sec x + \tan x)y] = \sec^2 x + \sec x \tan x$$

$$\therefore (\sec x + \tan x)y = \int \sec^2 x + \sec x \tan x \, dx$$

$$= \tan x + \sec x + c$$

$$\therefore y = 1 + \frac{c}{\sec x + \tan x}$$

Given also that $y = 2$, when $x = 0$

$$\therefore 2 = 1 + \frac{c}{1 + 0}$$

$$\therefore c = 1$$

$$\text{So } y = 1 + \frac{1}{\sec x + \tan x} \text{ or } y = 1 + \frac{\cos x}{1 + \sin x}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 1

Question:

$$\frac{dy}{dx} = \frac{y}{x} + \frac{x}{y}, \quad x > 0, y > 0$$

Solution:

$$z = \frac{y}{x} \Rightarrow y = xz$$

$$\therefore \frac{dy}{dx} = z + x \frac{dz}{dx}$$

Use the given substitution to express $\frac{dy}{dx}$ in terms of z , x and $\frac{dz}{dx}$.

Substitute into the equation:

$$\frac{dy}{dx} = \frac{y}{x} + \frac{x}{y}$$

$$\therefore z + x \frac{dz}{dx} = z + \frac{1}{z}$$

$$\therefore x \frac{dz}{dx} = \frac{1}{z}$$

Separate the variables:

$$\text{Then } \int z \, dz = \int \frac{1}{x} \, dx$$

$$\therefore \frac{z^2}{2} = \ln x + c$$

$$\therefore \frac{y^2}{2x^2} = \ln x + c, \text{ as } z = \frac{y}{x}$$

$$\therefore y^2 = 2x^2 (\ln x + c)$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 2

Question:

$$\frac{dy}{dx} = \frac{y}{x} + \frac{x^2}{y^2}, \quad x > 0$$

Solution:

$$\text{As } z = \frac{y}{x}, y = zx \text{ and } \frac{dy}{dx} = z + x \frac{dz}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{y}{x} + \frac{x^2}{y^2} \Rightarrow z + x \frac{dz}{dx} = z + \frac{1}{z^2}$$

$$\therefore x \frac{dz}{dx} = \frac{1}{z^2}$$

Separate the variables:

$$\text{Then } \int z^2 dz = \int \frac{1}{x} dx$$

$$\therefore \frac{z^3}{3} = \ln x + c$$

$$\text{But } z = \frac{y}{x}$$

$$\therefore \frac{y^3}{3x^3} = \ln x + c$$

$$\therefore y^3 = 3x^3 (\ln x + c)$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 3

Question:

$$\frac{dy}{dx} = \frac{y}{x} + \frac{y^2}{x^2}, \quad x > 0$$

Solution:

$$\text{As } z = \frac{y}{x}, y = zx \text{ and } \frac{dy}{dx} = z + x \frac{dz}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{y}{x} + \frac{y^2}{x^2} \Rightarrow z + x \frac{dz}{dx} = z + z^2$$

$$\therefore x \frac{dz}{dx} = z^2$$

Separate the variables:

$$\therefore \int \frac{1}{z^2} dz = \int \frac{1}{x} dx$$

$$\therefore -\frac{1}{z} = \ln x + c$$

$$\therefore z = \frac{-1}{\ln x + c}$$

$$\text{But } z = \frac{y}{x}$$

$$\therefore y = \frac{-x}{\ln x + c}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 4

Question:

$$\frac{dy}{dx} = \frac{x^3 + 4y^3}{3xy^2}, x > 0$$

Solution:

$$z = \frac{y}{x} \Rightarrow y = zx \text{ and } \frac{dy}{dx} = z + x \frac{dz}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{x^3 + 4y^3}{3xy^2} \Rightarrow z + x \frac{dz}{dx} = \frac{x^3 + 4z^3 x^3}{3x z^2 x^2}$$

$$\therefore \begin{aligned} x \frac{dz}{dx} &= \frac{1 + 4z^3}{3z^2} - z \\ &= \frac{1 + z^3}{3z^2} \end{aligned}$$

Separate the variables:

$$\therefore \int \frac{3z^2}{1+z^3} dz = \int \frac{1}{x} dx$$

$$\therefore \ln(1+z^3) = \ln x + \ln A, \text{ where } A \text{ is constant}$$

$$\therefore \ln(1+z^3) = \ln Ax$$

$$\text{So } 1+z^3 = Ax$$

$$\text{And } z^3 = Ax - 1. \text{ But } z = \frac{y}{x}$$

$$\therefore \frac{y^3}{x^3} = Ax - 1$$

$$\therefore y^3 = x^3 (Ax - 1), \text{ where } A \text{ is a positive constant}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 5

Question:

Use the substitution $z = y^{-2}$ to transform the differential equation

$$\frac{dy}{dx} + \left(\frac{1}{2} \tan x\right) y = -(2 \sec x) y^3, \quad -\frac{\pi}{2} < x < \frac{\pi}{2}$$

into a differential equation in z and x . By first solving the transformed equation, find the general solution of the original equation, giving y in terms of x .

Solution:

Given $z = y^{-2} \therefore y = z^{-\frac{1}{2}}$

and $\frac{dy}{dx} = -\frac{1}{2} z^{-\frac{3}{2}} \frac{dz}{dx}$

Find $\frac{dy}{dx}$ in terms of $\frac{dz}{dx}$ and z .

$$\therefore \frac{dy}{dx} + \left(\frac{1}{2} \tan x\right) y = -(2 \sec x) y^3$$

$$\Rightarrow -\frac{1}{2} z^{-\frac{3}{2}} \frac{dz}{dx} + \left(\frac{1}{2} \tan x\right) z^{-\frac{1}{2}} = -2 \sec x z^{-\frac{3}{2}}$$

$$\therefore \frac{dz}{dx} - z \tan x = 4 \sec x \quad *$$

This is a first order equation which can be solved by using an integrating factor.

The integrating factor is $e^{-\int \tan x \, dx} = e^{\ln \cos x}$
 $= \cos x$

The equation that you obtain needs an integrating factor to solve it.

Multiply the equation $*$ by $\cos x$

Then $\cos x \times \frac{dz}{dx} - z \sin x = 4$

$$\therefore \frac{d}{dx} (z \cos x) = 4$$

$$\therefore z \cos x = \int 4 \, dx$$

$$= 4x + c$$

$$\therefore z = \frac{4x + c}{\cos x}$$

As $y = z^{-\frac{1}{2}}, y = \sqrt{\frac{\cos x}{4x + c}}$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 6

Question:

Use the substitution $z = x^{\frac{1}{2}}$ to transform the differential equation

$$\frac{dx}{dt} + t^2x = t^2x^{\frac{3}{2}}$$

into a differential equation in z and t . By first solving the transformed equation, find the general solution of the original equation, giving x in terms of t .

Solution:

$$\text{Given that } z = x^{\frac{1}{2}}, x = z^2 \text{ and } \frac{dx}{dt} = 2z \frac{dz}{dt}$$

\therefore The equation $\frac{dx}{dt} + t^2x = t^2x^{\frac{3}{2}}$ becomes

$$2z \frac{dz}{dt} + t^2z^2 = t^2z$$

Divide through by $2z$

$$\text{Then } \frac{dz}{dt} + \frac{1}{2} t^2 z = \frac{1}{2} t^2$$

The integrating factor is $e^{\int \frac{1}{2} t^2 dt} = e^{\frac{1}{6} t^3}$

$$\therefore e^{\frac{1}{6} t^3} \frac{dz}{dt} + \frac{1}{2} t^2 e^{\frac{1}{6} t^3} z = \frac{1}{2} t^2 e^{\frac{1}{6} t^3}$$

$$\therefore \frac{d}{dt} (ze^{\frac{1}{6} t^3}) = \frac{1}{2} t^2 e^{\frac{1}{6} t^3}$$

$$\begin{aligned} \therefore ze^{\frac{1}{6} t^3} &= \int \frac{1}{2} t^2 e^{\frac{1}{6} t^3} dt \\ &= e^{\frac{1}{6} t^3} + c \end{aligned}$$

$$\therefore z = 1 + ce^{-\frac{1}{6} t^3}$$

$$\text{But } x = z^2 \therefore x = (1 + ce^{-\frac{1}{6} t^3})^2$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 7

Question:

Use the substitution $z = y^{-1}$ to transform the differential equation

$$\frac{dy}{dx} - \frac{1}{x}y = \frac{(x+1)^3}{x}y^2$$

into a differential equation in z and x . By first solving the transformed equation, find the general solution of the original equation, giving y in terms of x .

Solution:

Let $z = y^{-1}$, then $y = z^{-1}$ and $\frac{dy}{dx} = -z^{-2} \frac{dz}{dx}$

So $\frac{dy}{dx} - \frac{1}{x}y = \frac{(x+1)^3}{x}y^2$ becomes:

$$-z^{-2} \frac{dz}{dx} - \frac{1}{x}z^{-1} = \frac{(x+1)^3}{x}z^{-2}$$

Multiply through by $-z^2$

$$\text{Then } \frac{dz}{dx} + \frac{1}{x}z = -\frac{(x+1)^3}{x}$$

The integrating factor is $e^{\int P dx} = e^{\int \frac{1}{x} dx} = e^{\ln x} = x$

$$\therefore x \frac{dz}{dx} + z = -(x+1)^3$$

$$\text{i.e. } \frac{d}{dx}(xz) = -(x+1)^3$$

$$\begin{aligned} \therefore xz &= -\int (x+1)^3 dx \\ &= -\frac{1}{4}(x+1)^4 + c \end{aligned}$$

$$\therefore z = -\frac{1}{4x}(x+1)^4 + \frac{c}{x}$$

$$\therefore y = -\frac{4x}{4c - (x+1)^4}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 8

Question:

Use the substitution $z = y^2$ to transform the differential equation

$$2(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{y}$$

into a differential equation in z and x . By first solving the transformed equation,

a find the general solution of the original equation, giving y in terms of x .

b Find the particular solution for which $y = 2$ when $x = 0$.

Solution:

a Given that $z = y^2$, and so $y = z^{\frac{1}{2}}$ and $\frac{dy}{dx} = \frac{1}{2} z^{-\frac{1}{2}} \frac{dz}{dx}$

The equation $2(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{y}$ becomes

$$2(1 + x^2) \times \frac{1}{2} z^{-\frac{1}{2}} \frac{dz}{dx} + 2x z^{\frac{1}{2}} = z^{-\frac{1}{2}}$$

Multiply the equation by $\frac{z^{\frac{1}{2}}}{1 + x^2}$

$$\text{Then } \frac{dz}{dx} + \frac{2x}{1 + x^2} z = \frac{1}{1 + x^2}$$

The integrating factor is $e^{\int \frac{2x}{1+x^2} dx} = e^{\ln(1+x^2)} = 1 + x^2$

$$\therefore (1 + x^2) \frac{dz}{dx} + 2xz = 1$$

$$\therefore \frac{d}{dx} [(1 + x^2)z] = 1$$

$$\therefore (1 + x^2)z = \int 1 \, dx$$

$$= x + c$$

$$\therefore z = \frac{x + c}{(1 + x^2)}$$

$$\text{As } y = z^{\frac{1}{2}}, \quad y = \sqrt{\frac{x + c}{(1 + x^2)}}$$

b When $x = 0$, $y = 2$ $\therefore 2 = \sqrt{c} \Rightarrow c = 4$

$$\therefore y = \sqrt{\frac{x + 4}{1 + x^2}}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 9

Question:

Show that the substitution $z = y^{-(n-1)}$ transforms the general equation

$$\frac{dy}{dx} + Py = Qy^n,$$

where P and Q are functions of x , into the linear equation $\frac{dz}{dx} - P(n-1)z = -Q(n-1)$ (Bernoulli's equation)

Solution:

Given $z = y^{-(n-1)}$

$$\therefore y = z^{-\frac{1}{(n-1)}}$$

$$\frac{dy}{dx} = \frac{-1}{n-1} z^{-\frac{1}{n-1}-1} \frac{dz}{dx}$$

$$= \frac{-1}{n-1} z^{-\frac{n}{n-1}} \frac{dz}{dx}$$

$\therefore \frac{dy}{dx} + Py = Qy^n$ becomes

$$\frac{-1}{n-1} z^{-\frac{n}{n-1}} \frac{dz}{dx} + P z^{-\frac{1}{n-1}} = Q z^{-\frac{n}{n-1}}$$

Multiply each term by $-(n-1) z^{\frac{n}{n-1}}$

$$\text{Then } \frac{dz}{dx} - P(n-1) z^{\frac{n}{n-1}} z^{-\frac{1}{n-1}} = -Q(n-1) z^{\frac{n}{n-1}} z^{-\frac{n}{n-1}}$$

$$\text{i.e. } \frac{dz}{dx} - P(n-1) z = -Q(n-1)$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise D, Question 10

Question:

Use the substitution $u = y + 2x$ to transform the differential equation

$$\frac{dy}{dx} = \frac{-(1 + 2y + 4x)}{1 + y + 2x}$$

into a differential equation in u and x . By first solving this new equation, show that the general solution of the original equation may be written $4x^2 + 4xy + y^2 + 2y + 2x = k$, where k is a constant

Solution:

Given $u = y + 2x$ and so $y = u - 2x$ and $\frac{dy}{dx} = \frac{du}{dx} - 2$

\therefore the differential equation $\frac{dy}{dx} = -\frac{(1 + 2y + 4x)}{1 + y + 2x}$ becomes

Rearrange the given substitution to give y in terms of u and x , and $\frac{dy}{dx}$ in terms of $\frac{du}{dx}$.

$$\frac{du}{dx} - 2 = -\frac{1 + 2u}{1 + u}$$

$$\therefore \frac{du}{dx} = \frac{-(1 + 2u) + 2(1 + u)}{1 + u}$$

$$\therefore \frac{du}{dx} = \frac{1}{1 + u}$$

Separate the variables

$$\int (1 + u) du = \int 1 \times dx$$

$$\therefore u + \frac{u^2}{2} = x + c, \text{ where } c \text{ is constant}$$

$$\text{And } (y + 2x) + \frac{(y + 2x)^2}{2} = x + c$$

$$2y + 4x + y^2 + 4xy + 4x^2 = 2x + 2c$$

$$\text{i.e. } 4x^2 + 4xy + y^2 + 2y + 2x = k, \text{ where } k = 2c$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 1

Question:


Solve the equation $\frac{dy}{dx} = \frac{x}{\sqrt{x^2 + 16}}$ and sketch three solution curves.

Solution:

$$\frac{dy}{dx} = \frac{x}{\sqrt{x^2 + 16}}$$

$$\begin{aligned}\therefore y &= \int \frac{x}{\sqrt{x^2 + 16}} dx \\ &= (x^2 + 16)^{\frac{1}{2}} + c\end{aligned}$$

The integral is of the type $\int [f(x)]^n f'(x) dx$, which integrates to give $[f(x)]^{n+1} + c$.



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 2

Question:

Solve the equation $\frac{dy}{dx} = xy$ and sketch the solution curves which pass through

a (0, 1)

b (0, 2)

c (0, 3)

Solution:

$$\frac{dy}{dx} = xy$$

Separate the variables and integrate.

$$\therefore \int \frac{1}{y} dy = \int x dx$$

$$\therefore \ln y = \frac{1}{2}x^2 + c, \text{ where } c \text{ is constant}$$

$$\therefore y = e^{\frac{1}{2}x^2 + c}$$

$$= e^c e^{\frac{1}{2}x^2} = Ae^{\frac{1}{2}x^2}, \text{ where } A \text{ is } e^c$$

a The solution which satisfies $x = 0$ when $y = 1$

$$\text{is } y = Ae^{\frac{1}{2}x^2} \text{ where } 1 = Ae^0 \quad \text{i.e. } A = 1$$

$$\therefore y = e^{\frac{1}{2}x^2}$$


b The solution for which $y = 2$ when $x = 0$ is $y = Ae^{\frac{1}{2}x^2}$

$$\text{with } 2 = Ae^0 \quad \text{i.e. } A = 2$$

$$\therefore y = 2e^{\frac{1}{2}x^2}$$

c The solution for which $y = 3$ when $x = 0$ is $y = 3e^{\frac{1}{2}x^2}$

The solution curves are shown in the sketch.



Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 3

Question:

Solve the equation $\frac{dv}{dt} = -g - kv$ given that $v = u$ when $t = 0$, and that u , g and k are positive constants. Sketch the solution curve indicating the velocity which v approaches as t becomes large.

Solution:

$$\frac{dv}{dt} = -g - kv$$

You can separate the variables by dividing both sides by $(g + kv)$, or you could rearrange the equation as $\frac{dv}{dt} + kv = g$ and use the integrating factor e^{kt} .

$$\therefore \int \frac{dv}{g + kv} = - \int 1 dt$$

$$\therefore \frac{1}{k} \ln |g + kv| = -t + c \text{ where } c \text{ is a constant} \quad *$$

When $t = 0$, $v = u$

$$\therefore \frac{1}{k} \ln |g + ku| = c$$

\therefore Substituting c back into the equation $*$


$$\frac{1}{k} \ln |g + kv| = -t + \frac{1}{k} \ln |g + ku|$$

$$\therefore \frac{1}{k} [\ln |g + kv| - \ln |g + ku|] = -t$$

$$\therefore \ln \frac{g + kv}{g + ku} = -kt$$

$$\therefore g + kv = (g + ku) e^{-kt}$$

$$\therefore v = \frac{1}{k} [(g + ku) e^{-kt} - g]$$



The required velocity is $-\frac{g}{k} \text{ m s}^{-1}$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 4

Question:

Solve the equation $\frac{dy}{dx} + y \tan x = 2 \sec x$

Solution:

$$\frac{dy}{dx} + y \tan x = 2 \sec x$$

Use an integrating factor to solve this equation.

Use the integrating factor $e^{\int \tan x \, dx} = e^{\ln \sec x} = \sec x$

$$\therefore \sec x \frac{dy}{dx} + y \sec x \tan x = 2 \sec^2 x$$

$$\therefore \frac{d}{dx} (y \sec x) = 2 \sec^2 x$$

$$\therefore y \sec x = \int 2 \sec^2 x \, dx$$

$$= 2 \tan x + c$$

$$\therefore y = 2 \sin x + c \cos x$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 5

Question:

Solve the equation $(1 - x^2) \frac{dy}{dx} + xy = 5x$ $-1 < x < 1$

Solution:

$$(1 - x^2) \frac{dy}{dx} + xy = 5x$$

Divide through by $(1 - x^2)$, then find the integrating factor.

Divide through by $(1 - x^2)$

$$\therefore \frac{dy}{dx} + \frac{x}{1 - x^2} y = \frac{5x}{1 - x^2}$$

Use the integrating factor $e^{\int \frac{x}{1 - x^2} dx} = e^{-\frac{1}{2} \ln(1 - x^2)}$

$$= e^{\ln(1 - x^2)^{-\frac{1}{2}}} = \frac{1}{\sqrt{1 - x^2}}$$

$$\therefore \frac{1}{\sqrt{1 - x^2}} \frac{dy}{dx} + \frac{x}{(1 - x^2)^{\frac{3}{2}}} y = \frac{5x}{(1 - x^2)^{\frac{3}{2}}}$$

$$\therefore \frac{d}{dx} [(1 - x^2)^{-\frac{1}{2}} y] = \frac{5x}{(1 - x^2)^{\frac{3}{2}}}$$

$$\therefore (1 - x^2)^{-\frac{1}{2}} y = \int \frac{5x}{(1 - x^2)^{\frac{3}{2}}} dx$$

$$= 5(1 - x^2)^{-\frac{1}{2}} + c$$

$$\therefore y = 5 + c(1 - x^2)^{\frac{1}{2}}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 6

Question:

Solve the equation $x \frac{dy}{dx} + x + y = 0$

Solution:

$$x \frac{dy}{dx} + x + y = 0$$

Take the 'x' term to the other side of the equation.

$$\therefore x \frac{dy}{dx} + y = -x$$

This is an exact equation.

$$\text{So } \frac{d}{dx}(xy) = -x$$

$$\begin{aligned} \therefore xy &= -\int x \, dx \\ &= -\frac{1}{2}x^2 + c \end{aligned}$$

$$\therefore y = -\frac{1}{2}x + \frac{c}{x}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 7

Question:

Solve the equation $\frac{dy}{dx} + \frac{y}{x} = \sqrt{x}$

Solution:

$$\frac{dy}{dx} + \frac{y}{x} = \sqrt{x}$$

The integrating factor is $e^{\int \frac{1}{x} dx} = e^{\ln x} = x$

Multiply the differential equation by the integrating factor:

$$x \frac{dy}{dx} + y = x\sqrt{x}$$

$$\therefore \frac{d}{dx}(xy) = x^{\frac{3}{2}}$$

$$\therefore xy = \int x^{\frac{3}{2}} dx$$

$$= \frac{2}{5} x^{\frac{5}{2}} + c$$

$$\therefore y = \frac{2}{5} x^{\frac{3}{2}} + \frac{c}{x}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 8

Question:

Solve the equation $\frac{dy}{dx} + 2xy = x$

Solution:

$$\frac{dy}{dx} + 2xy = x$$

The integrating factor is $e^{\int 2x \, dx} = e^{x^2}$

Multiply the differential equation by e^{x^2}

$$\therefore e^{x^2} \frac{dy}{dx} + 2xe^{x^2}y = xe^{x^2}$$

$$\therefore \frac{d}{dx} (e^{x^2}y) = xe^{x^2}$$

$$\begin{aligned} \therefore ye^{x^2} &= \int xe^{x^2} \, dx \\ &= \frac{1}{2}e^{x^2} + c \end{aligned}$$

$$\therefore y = \frac{1}{2} + ce^{-x^2}$$

© Pearson Education Ltd 2009

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 9

Question:

Solve the equation $x(1 - x^2) \frac{dy}{dx} + (2x^2 - 1)y = 2x^3$ $0 < x < 1$

Solution:

$$x(1 - x^2) \frac{dy}{dx} + (2x^2 - 1)y = 2x^3$$

Divide through by $x(1 - x^2)$

$$\therefore \frac{dy}{dx} + \frac{2x^2 - 1}{x(1 - x^2)} y = \frac{2x^3}{x(1 - x^2)} \quad *$$

You will need to use partial fractions to integrate $\frac{2x^2 - 1}{x(1 - x^2)}$ and to find the integrating factor.

The integrating factor is $e^{\int \frac{2x^2 - 1}{x(1 - x^2)} dx}$

$$\begin{aligned} \int \frac{2x^2 - 1}{x(1 - x)(1 + x)} dx &= \int \left(-\frac{1}{x} + \frac{1}{2(1 - x)} - \frac{1}{2(1 + x)} \right) dx \\ &= -\ln x - \frac{1}{2} \ln(1 - x) - \frac{1}{2} \ln(1 + x) \\ &= -\ln x \sqrt{1 - x^2} \end{aligned}$$

So the integrating factor is $e^{-\ln x \sqrt{1 - x^2}} = e^{\ln \frac{1}{x\sqrt{1 - x^2}}} = \frac{1}{x\sqrt{1 - x^2}}$

Multiply the differential equation $*$ by $\frac{1}{x\sqrt{1 - x^2}}$

$$\therefore \frac{1}{x\sqrt{1 - x^2}} \frac{dy}{dx} + \frac{2x^2 - 1}{x^2(1 - x^2)^{\frac{3}{2}}} y = \frac{2x}{(1 - x^2)^{\frac{3}{2}}}$$

$$\therefore \frac{d}{dx} \left[\frac{1}{x\sqrt{1 - x^2}} y \right] = \frac{2x}{(1 - x^2)^{\frac{3}{2}}}$$

$$\begin{aligned} \therefore \frac{y}{x\sqrt{1 - x^2}} &= \int \frac{2x}{(1 - x^2)^{\frac{3}{2}}} dx \\ &= 2(1 - x^2)^{-\frac{1}{2}} + c \end{aligned}$$

$$\therefore y = 2x + cx\sqrt{1 - x^2}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 10

Question:

Solve the equation $R \frac{dq}{dt} + \frac{q}{c} = E$ when

a $E = 0$

b $E = \text{constant}$

c $E = \cos pt$

(R , c and p are constants)

Solution:

$$R \frac{dq}{dt} + \frac{q}{c} = E$$

$$\therefore \frac{dq}{dt} + \frac{1}{Rc} q = \frac{E}{R}$$

The integrating factor is $e^{\int \frac{1}{Rc} dt} = e^{\frac{t}{Rc}}$

$$\therefore e^{\frac{t}{Rc}} \frac{dq}{dt} + \frac{1}{Rc} e^{\frac{t}{Rc}} q = \frac{E}{R} e^{\frac{t}{Rc}}$$

$$\therefore \frac{d}{dt} (q e^{\frac{t}{Rc}}) = \frac{E}{R} e^{\frac{t}{Rc}}$$

$$\therefore q e^{\frac{t}{Rc}} = \int \frac{E}{R} e^{\frac{t}{Rc}} dt$$

a When $E = 0$

$$\therefore q e^{\frac{t}{Rc}} = k, \text{ where } k \text{ is constant.}$$

$$\therefore q = k e^{-\frac{t}{Rc}}$$

b When $E = \text{constant}$

$$q e^{\frac{t}{Rc}} = \int \frac{E}{R} e^{\frac{t}{Rc}} dt$$

$$= E c e^{\frac{t}{Rc}} + k, \text{ where } k \text{ is constant}$$

$$\therefore q = Ec + k e^{-\frac{t}{Rc}}$$

c When $E = \cos pt$

$$q e^{\frac{t}{Rc}} = \int \frac{1}{R} \cos pt e^{\frac{t}{Rc}} dt \quad *$$

$$\text{i.e. } \int \frac{1}{R} \cos pt e^{\frac{t}{Rc}} dt = c e^{\frac{t}{Rc}} \cos pt + \int c p e^{\frac{t}{Rc}} \sin pt dt \quad \leftarrow \text{Use integration by parts.}$$

$$\int \frac{1}{R} \cos pt e^{\frac{t}{Rc}} dt = c e^{\frac{t}{Rc}} \cos pt + R p c^2 e^{\frac{t}{Rc}} \sin pt - \int R p^2 c^2 e^{\frac{t}{Rc}} \cos pt dt \quad \leftarrow \text{Use 'parts' again.}$$

$$\therefore \int \left(\frac{1}{R} + R p^2 c^2 \right) e^{\frac{t}{Rc}} \cos pt dt = c e^{\frac{t}{Rc}} (\cos pt + R p c \sin pt) + k, \text{ where } k \text{ is a constant}$$

$$\therefore \frac{1}{R} \int e^{\frac{t}{Rc}} \cos pt dt = \frac{c}{(1 + R^2 p^2 c^2)} e^{\frac{t}{Rc}} (\cos pt + R p c \sin pt) + \frac{k}{(1 + R^2 p^2 c^2)}$$

From $*$

$$q e^{\frac{t}{Rc}} = \frac{c}{(1 + R^2 p^2 c^2)} e^{\frac{t}{Rc}} (\cos pt + R p c \sin pt) + \frac{k}{(1 + R^2 p^2 c^2)}$$

$$\therefore q = \frac{c}{(1 + R^2 p^2 c^2)} (\cos pt + R p c \sin pt) + k' e^{-\frac{t}{Rc}}, \text{ where } k' = \frac{k}{1 + R^2 p^2 c^2} \text{ is constant}$$

This is a difficult question – particularly part **c**.
You may decide to omit this question, unless
you want a challenge.

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 11

Question:

Find the general solution of the equation $\frac{dy}{dx} - ay = Q$, where a is a constant, giving your answer in terms of a , when

a $Q = ke^{\lambda x}$

b $Q = ke^{ax}$

c $Q = kx^n e^{ax}$.

(k , λ and n are constants).

Solution:

Given that $\frac{dy}{dx} - ay = Q$

The integrating factor is $e^{\int -a dx} = e^{-ax}$

Then $e^{-ax} \frac{dy}{dx} - ae^{-ax} y = Qe^{-ax}$

$\therefore \frac{d}{dx} (ye^{-ax}) = Qe^{-ax}$

$\therefore ye^{-ax} = \int Qe^{-ax} dx$

a When $Q = ke^{\lambda x}$

$$ye^{-ax} = \int ke^{(\lambda - a)x} dx$$

$$= \frac{k}{\lambda - a} e^{(\lambda - a)x} + c, \text{ where } c \text{ is constant}$$

$$\therefore y = \frac{k}{\lambda - a} e^{\lambda x} + ce^{ax}$$

When $\lambda \neq a$.
For $\lambda = a$, see part **b**.

b When $Q = ke^{ax}$

$$ye^{-ax} = \int k dx$$

$$= kx + c, \text{ where } c \text{ is constant}$$

$$\therefore y = (kx + c)e^{ax}$$

c When $Q = kx^n e^{ax}$

$$ye^{-ax} = \int kx^n dx$$

$$= \frac{kx^{n+1}}{n+1} + c, \text{ where } c \text{ is constant}$$

$$\therefore y = \frac{kx^{n+1}}{n+1} e^{ax} + ce^{ax}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 12

Question:

Use the substitution $z = y^{-1}$ to transform the differential equation $x \frac{dy}{dx} + y = y^2 \ln x$, into a linear equation. Hence obtain the general solution of the original equation.

Solution:

Given that $z = y^{-1}$, then $y = z^{-1}$ so $\frac{dy}{dx} = -z^{-2} \frac{dz}{dx}$.

The equation $x \frac{dy}{dx} + y = y^2 \ln x$ becomes

$$-xz^{-2} \frac{dz}{dx} + z^{-1} = z^{-2} \ln x$$

Divide through by $-xz^{-2}$

$$\therefore \frac{dz}{dx} - \frac{z}{x} = -\frac{\ln x}{x}$$

The integrating factor is $e^{-\int \frac{1}{x} dx} = e^{-\ln x} = e^{\ln \frac{1}{x}} = \frac{1}{x}$

$$\therefore \frac{1}{x} \frac{dz}{dx} - \frac{z}{x^2} = -\frac{\ln x}{x^2}$$

$$\therefore \frac{d}{dx} \left(\frac{1}{x} z \right) = -\frac{\ln x}{x^2}$$

$$\begin{aligned} \therefore \frac{1}{x} z &= -\int \frac{1}{x^2} \ln x \, dx \\ &= -\left[-\frac{1}{x} \ln x + \int \frac{1}{x^2} dx \right] \\ &= \frac{1}{x} \ln x + \frac{1}{x} + c \end{aligned}$$

$$\therefore z = \ln x + 1 + cx.$$

$$\text{As } y = z^{-1} \quad \therefore y = \frac{1}{1 + cx + \ln x}$$

Use the substitution to express y in terms of z and $\frac{dy}{dx}$ in terms of z and $\frac{dz}{dx}$.

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 13

Question:

Use the substitution $z = y^2$ to transform the differential equation

$2 \cos x \frac{dy}{dx} - y \sin x + y^{-1} = 0$, into a linear equation. Hence obtain the general solution of the original equation.

Solution:

Given that $z = y^2$, $y = z^{\frac{1}{2}}$ and $\frac{dy}{dx} = \frac{1}{2} z^{-\frac{1}{2}} \frac{dz}{dx}$

The differential equation

$$2 \cos x \frac{dy}{dx} - y \sin x + y^{-1} = 0 \text{ becomes}$$

$$\cos x z^{-\frac{1}{2}} \frac{dz}{dx} - z^{\frac{1}{2}} \sin x + z^{-\frac{1}{2}} = 0$$

Divide through by $z^{-\frac{1}{2}}$

$$\text{then } \cos x \frac{dz}{dx} - z \sin x = -1$$

This becomes an exact equation which can be solved directly.

$$\therefore \frac{d}{dx} (z \cos x) = -1$$

$$\therefore z \cos x = -\int 1 \, dx$$

$$= -x + c$$

$$\therefore z = \frac{c - x}{\cos x}$$

$$\therefore y = \sqrt{\frac{c - x}{\cos x}}$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 14

Question:

Use the substitution $z = \frac{y}{x}$ to transform the differential equation $(x^2 - y^2) \frac{dy}{dx} - xy = 0$, into a linear equation. Hence obtain the general solution of the original equation.

Solution:

Given that $z = \frac{y}{x}$, $y = zx$ so $\frac{dy}{dx} = z + x \frac{dz}{dx}$

The equation $(x^2 - y^2) \frac{dy}{dx} - xy = 0$ becomes

$$(x^2 - z^2x^2) \left(z + x \frac{dz}{dx} \right) - xzx = 0$$

$$\therefore (1 - z^2)z + (1 - z^2)x \frac{dz}{dx} - z = 0$$

$$\therefore x \frac{dz}{dx} = \frac{z}{1 - z^2} - z$$

$$\text{i.e.} \quad x \frac{dz}{dx} = \frac{z^3}{1 - z^2}$$

Separate the variables to give

$$\int \frac{1 - z^2}{z^3} dz = \int \frac{1}{x} dx$$

$$\therefore \int (z^{-3} - z^{-1}) dz = \int x^{-1} dx$$

$$\therefore \frac{z^{-2}}{-2} - \ln z = \ln x + c$$

$$\therefore -\frac{1}{2z^2} = \ln x + \ln z + c$$

$$= \ln xz + c$$

$$\text{But} \quad y = zx$$

$$\therefore (c + \ln y) = -\frac{x^2}{2y^2}$$

$$\therefore 2y^2 (\ln y + c) + x^2 = 0$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 15

Question:

Use the substitution $z = \frac{y}{x}$ to transform the differential equation $\frac{dy}{dx} = \frac{y(x+y)}{x(y-x)}$, into a linear equation. Hence obtain the general solution of the original equation.

Solution:

$$z = \frac{y}{x}, \quad \therefore y = xz \text{ and } \frac{dy}{dx} = z + x \frac{dz}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{y(x+y)}{x(y-x)} \text{ becomes } z + x \frac{dz}{dx} = \frac{xz(x+xz)}{x(xz-x)}$$

$$\therefore z + x \frac{dz}{dx} = \frac{z(1+z)}{(z-1)}$$

$$\begin{aligned} \text{So } x \frac{dz}{dx} &= \frac{z(1+z)}{z-1} - z \\ &= \frac{2z}{z-1} \end{aligned}$$

Separating the variables

$$\int \frac{(z-1)}{2z} dz = \int \frac{1}{x} dx$$

$$\therefore \int \left(\frac{1}{2} - \frac{1}{2z} \right) dz = \int \frac{1}{x} dx$$

$$\therefore \frac{1}{2}z - \frac{1}{2}\ln z = \ln x + c$$

$$\text{As } z = \frac{y}{x} \quad \therefore \frac{y}{2x} - \frac{1}{2}\ln \frac{y}{x} = \ln x + c$$

$$\therefore \frac{y}{2x} - \frac{1}{2}\ln y + \frac{1}{2}\ln x = \ln x + c$$

$$\therefore \frac{y}{2x} - \frac{1}{2}\ln y = \frac{1}{2}\ln x + c$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 16

Question:

Use the substitution $z = \frac{y}{x}$ to transform the differential equation $\frac{dy}{dx} = \frac{-3xy}{(y^2 - 3x^2)}$, into a linear equation. Hence obtain the general solution of the original equation.

Solution:

Given that $z = \frac{y}{x}$, so $y = zx$ and $\frac{dy}{dx} = z + x \frac{dz}{dx}$

The equation $\frac{dy}{dx} = \frac{-3xy}{y^2 - 3x^2}$ becomes

$$z + x \frac{dz}{dx} = \frac{-3x^2z}{z^2x^2 - 3x^2}$$

$$\begin{aligned} \text{i.e.} \quad x \frac{dz}{dx} &= \frac{-3z}{z^2 - 3} - z \\ &= \frac{-z^3}{z^2 - 3} \end{aligned}$$

Separate the variables:

$$\text{Then } \int \left(\frac{z^2 - 3}{z^3} \right) dz = - \int \frac{1}{x} dx.$$

$$\therefore \int \left(\frac{1}{z} - 3z^{-3} \right) dz = -\ln x + c$$

$$\therefore \ln z + \frac{3}{2} z^{-2} = -\ln x + c$$

$$\therefore \ln zx + \frac{3}{2z^2} = c$$

$$\text{But } zx = y \text{ and } z = \frac{y}{x}$$

$$\therefore \ln y + \frac{3x^2}{2y^2} = c$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 17

Question:

Use the substitution $u = x + y$ to transform the differential equation

$\frac{dy}{dx} = (x + y + 1)(x + y - 1)$ into a differential equation in u and x . By first solving this new equation, find the general solution of the original equation, giving y in terms of x .

Solution:

Let $u = x + y$, then $\frac{du}{dx} = 1 + \frac{dy}{dx}$ and so $\frac{dy}{dx} = (x + y + 1)(x + y - 1)$ becomes

$$\begin{aligned}\frac{du}{dx} - 1 &= (u + 1)(u - 1) \\ &= u^2 - 1\end{aligned}$$

$$\therefore \frac{du}{dx} = u^2$$

Separate the variables.

$$\text{Then} \quad \int \frac{1}{u^2} du = \int 1 dx$$

$$\therefore -\frac{1}{u} = x + c$$

$$\text{But } u = x + y \quad \therefore -\frac{1}{x + y} = x + c$$

$$\therefore y + x = \frac{-1}{x + c}$$

$$y = \frac{-1}{x + c} - x$$

Solutionbank FP2

Edexcel AS and A Level Modular Mathematics

Exercise E, Question 18

Question:

Use the substitution $u = y - x - 2$ to transform the differential equation $\frac{dy}{dx} = (y - x - 2)^2$ into a differential equation in u and x . By first solving this new equation, find the general solution of the original equation, giving y in terms of x .

Solution:

Given that $u = y - x - 2$, and so $\frac{du}{dx} = \frac{dy}{dx} - 1$

$$\therefore \frac{dy}{dx} = (y - x - 2)^2 \text{ becomes } \frac{du}{dx} + 1 = u^2$$

$$\text{i.e.} \quad \frac{du}{dx} = u^2 - 1$$

$$\therefore \int \frac{1}{u^2 - 1} du = \int 1 dx \quad \bullet \quad \begin{array}{l} \text{Factorise } \frac{1}{u^2 - 1} \text{ into } \frac{1}{(u - 1)(u + 1)} \\ \text{and use partial fractions.} \end{array}$$

$$\therefore \int \left(\frac{1}{2(u - 1)} - \frac{1}{2(u + 1)} \right) du = x + c \text{ where } c \text{ is constant}$$

$$\therefore \frac{1}{2} \ln(u - 1) - \frac{1}{2} \ln(u + 1) = x + c$$

$$\therefore \frac{1}{2} \ln \frac{u - 1}{u + 1} = x + c$$

$$\therefore \frac{u - 1}{u + 1} = e^{2x + 2c} = Ae^{2x} \text{ where } A = e^{2c} \text{ is a constant}$$

$$\therefore u - 1 = Ae^{2x} + Ae^{2x}$$

$$\therefore u(1 - Ae^{2x}) = (1 + Ae^{2x})$$

$$\therefore u = \frac{1 + Ae^{2x}}{1 - Ae^{2x}}$$

$$\text{But } u = y - x - 2$$

$$\therefore y = x + 2 + \frac{1 + Ae^{2x}}{1 - Ae^{2x}}$$