

Solutionbank M5

Edexcel AS and A Level Modular Mathematics


Exercise A, Question 1

Question:


A rocket is launched vertically upwards under gravity from rest at time $t = 0$. The rocket propels itself upwards by ejecting burnt fuel vertically downwards at a constant speed u relative to the rocket. At time t seconds after the launch the rocket has velocity v and mass $(M - kt)$. Derive the equation of motion for the rocket. Ignore air resistance.

Solution:

At time t



After an interval δt :



Change in momentum: $(m + \delta m)(v + \delta v) + (-\delta m)(v - u) - mv = -mg\delta t$

$$\Rightarrow m \frac{dv}{dt} + u \frac{dm}{dt} = -mg$$

$$m = M - kt \Rightarrow \frac{dm}{dt} = -k, (M - kt) \frac{dv}{dt} - ku = -(M - kt)g$$

$$\frac{dv}{dt} = \frac{ku}{M - kt} - g$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 2

Question:

A spaceship is moving in deep space with no external forces acting on it. At time t the spaceship has total mass m and is moving with velocity v . The spaceship reduces its speed by ejecting fuel from its front end with a speed c relative to itself and in the same direction as its own motion.


a Show that $\frac{dv}{dm} = \frac{c}{m}$.

Initially the spaceship is moving with speed V and has total mass M . Its speed is reduced to $\frac{1}{2}V$.

b Find the mass of fuel ejected.

Solution:

a At time t



After interval δt



$$\text{Change in momentum} \Rightarrow (m + \delta m)(v + \delta v) + (-\delta m)(v + c) - mv = 0$$

$$mv + m\delta v + v\delta m + \delta m\delta v - v\delta m - c\delta m - mv = 0$$

$$m\delta v + \delta m\delta v - c\delta m = 0$$

$$\Rightarrow m \frac{\delta v}{\delta m} + \delta v - c = 0, \Rightarrow \frac{dv}{dm} = \frac{c}{m}$$

b $\frac{dv}{dm} = \frac{c}{m} \Rightarrow \int_v^{\frac{V}{2}} \frac{1}{m} dm = \int_M^m \frac{1}{m} dm$

$$-\frac{V}{2} = c \left[\ln m \right]_M^m = c \ln \left(\frac{m}{M} \right)$$

$$\Rightarrow -\frac{V}{2c} = \ln \left(\frac{m}{M} \right), e^{-\frac{V}{2c}} = \frac{m}{M}, m = Me^{-\frac{V}{2c}}$$

$$\text{Mass of fuel ejected} = M \left(1 - e^{-\frac{V}{2c}} \right)$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 3


Question:

A rocket is launched vertically upwards from rest. The initial mass of the rocket and its fuel is 1000 kg. The rocket burns fuel at the rate of 20 kg s^{-1} . The burnt fuel is ejected vertically downwards with a speed of 2000 m s^{-1} relative to the rocket, and burning stops after 30 seconds. At time t seconds ($t < 30$) after the launch, the speed of the rocket is $v \text{ m s}^{-1}$. Air resistance may be assumed to be negligible.

- a Show that $-g(50-t) = (50-t) \frac{dv}{dt} - 2000$.
 b Find the speed of the rocket when the burning stops.

Solution:

- a At time t



After interval δt



Considering the change in momentum:

$$(m + \delta m)(v + \delta v) + (-\delta m)(v - 2000) - mv = -mg\delta t$$

$$\Rightarrow m \frac{dv}{dt} + 2000 \frac{dm}{dt} = -mg$$

At time t , $m = 1000 - 20t$

$$\Rightarrow (1000 - 20t) \frac{dv}{dt} + 2000 \times -20 = -(1000 - 20t)g$$

$$\text{Dividing by } 20 \Rightarrow (50 - t) \frac{dv}{dt} - 2000 = -g(50 - t)$$

- b $\frac{dv}{dt} = -g + \frac{2000}{50 - t}$
 $\Rightarrow \int_0^v 1 dv = \int_0^{30} -g + \frac{2000}{50 - t} dt$
 $V = [-gt - 2000 \ln(50 - t)]_0^{30}$
 $= -30g - 2000 \ln 20 + 0 + 2000 \ln 50 \approx 1540 \text{ m s}^{-1}$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 4

Question:


A rocket is launched vertically upwards from rest. The rocket expels burnt fuel vertically downwards with speed u relative to the rocket. Initially the rocket has mass

M . At time t the rocket has speed v and mass $M\left(1 - \frac{1}{3}t\right)$. Ignore air resistance.


- a Show that $\frac{dv}{dt} = \frac{u}{3-t} - g$.
- b Find the speed of the rocket when $t = 1$.
- c Find the height of the rocket above the launch site when $t = 1$.

Solution:

a At time t



After interval δt



$$(m + \delta m)(v + \delta v) + (-\delta m)(v - u) - mv = -mg \delta t$$

$$m\delta v + \delta m\delta v + u\delta m = -mg \delta t$$

$$\Rightarrow m \frac{dv}{dt} + u \frac{dm}{dt} = -mg$$

$$m = M \left(1 - \frac{1}{3}t \right)$$

$$\Rightarrow M \left(1 - \frac{1}{3}t \right) \frac{dv}{dt} + u \left(-\frac{1}{3}M \right) = -M \left(1 - \frac{1}{3}t \right) g$$

$$\frac{dv}{dt} = \frac{\frac{1}{3}u}{1 - \frac{1}{3}t} - g = \frac{u}{3-t} - g$$

b $\frac{dv}{dt} = \frac{u}{3-t} - g \Rightarrow v = \int \frac{u}{3-t} - g dt = -u \ln |3-t| - gt + C$

$$t = 0, v = 0 \Rightarrow 0 = -u \ln 3 + C; v = u \ln \left| \frac{3}{3-t} \right| - gt = u \ln \frac{3}{2} - g \text{ when } t = 1$$

c $v = \frac{dx}{dt} = u \ln \left(\frac{3}{3-t} \right) - gt, t < 3$

Using integration by parts

$$\Rightarrow x = \int u \ln \left(\frac{3}{3-t} \right) - gt dt = \int u \ln 3 - u \ln(3-t) - gt dt$$

$$= (u \ln 3)t + u(3-t) \ln(3-t) - u(3-t) - \frac{1}{2}gt^2 + C$$

$$t = 0, x = 0 \Rightarrow 0 = 0 + 3u \ln 3 - 3u + C, C = 3u - 3u \ln 3$$

$$\text{When } t = 1, x = u \ln 3 + 2u \ln 2 - 2u - \frac{g}{2} + 3u - 3u \ln 3 = 2u \ln \frac{2}{3} + u - \frac{g}{2}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 5

Question:

A spherical hailstone is falling under gravity in still air. At time t the hailstone has speed v . The radius r increases by condensation. Given that $\frac{dr}{dt} = kr$, where k is a constant, and neglecting air resistance,

a show that $\frac{dv}{dt} = g - 3kv$,


b find the time taken for the speed of the hailstone to increase from $\frac{g}{9k}$ to $\frac{g}{6k}$.

Solution:

a At time t



After time δt



$$[(m + \delta m)(v + \delta v)] - [mv + \delta m \times 0] = (m + \delta m)g \delta t$$

$$\Rightarrow m \frac{\delta v}{\delta t} + v \frac{\delta m}{\delta t} + \frac{\delta m \delta v}{\delta t} = mg + g \delta m$$

$$\text{so } m \frac{dv}{dt} + v \frac{dm}{dt} = mg$$

The mass of the hailstone is $\lambda \times \frac{4}{3} \pi r^3$

$$\Rightarrow \frac{dm}{dt} = 4 \lambda \pi r^2 \frac{dr}{dt} = 4 \lambda \pi r^2 \times kr = 4k \lambda \pi r^3$$

$$\Rightarrow \lambda \times \frac{4}{3} \pi r^3 \frac{dv}{dt} + v \times 4k \lambda \pi r^3 = \lambda \times \frac{4}{3} \pi r^3 g$$

And therefore $\frac{dv}{dt} = g - 3kv$

$$\begin{aligned} \text{b } \frac{dv}{dt} = g - 3kv &\Rightarrow t = \int_{\frac{g}{9k}}^{\frac{g}{6k}} \frac{1}{g - 3kv} dv = \left[-\frac{1}{3k} \ln |g - 3kv| \right]_{\frac{g}{9k}}^{\frac{g}{6k}} \\ &= -\frac{1}{3k} \ln \left(\frac{g - \frac{3kg}{6k}}{g - \frac{3kg}{9k}} \right) = -\frac{1}{3k} \ln \frac{g \times 3}{2 \times 2g} = \frac{1}{3k} \ln \frac{4}{3} \end{aligned}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics


Exercise A, Question 6

Question:


A spaceship is moving in deep space with no external forces acting on it. Initially it has total mass M and is moving with velocity V . The spaceship reduces its speed to $\frac{3}{5}V$ by ejecting fuel from its front end with a speed c relative to itself and in the same direction as its own motion. Find the mass of fuel ejected.

Solution:

At time t



After interval δt



$$\begin{aligned}\text{Change in momentum} &\Rightarrow (m + \delta m)(v + \delta v) + (-\delta m)(v + c) - mv = 0 \\ mv + m\delta v + v\delta m + \delta m\delta v - v\delta m - c\delta m - mv &= 0 \\ m\delta v + \delta m\delta v - c\delta m &= 0 \\ \Rightarrow m \frac{\delta v}{\delta m} + \delta v - c &= 0, \Rightarrow \frac{dv}{dm} = \frac{c}{m}\end{aligned}$$

$$\text{Speed reduced from } V \text{ to } \frac{3}{5}V: \int_V^{\frac{3}{5}V} 1 dv = c \int_M^m \frac{1}{m} dm$$

$$-\frac{2V}{5} = c [\ln m]_M^m = c \ln \left(\frac{m}{M} \right)$$

$$\Rightarrow -\frac{2V}{5c} = \ln \left(\frac{m}{M} \right), c \frac{2V}{5c} = \frac{m}{M}, m = Me^{\frac{2V}{5c}}$$

$$\text{Mass of fuel ejected} = M \left(1 - e^{\frac{2V}{5c}} \right)$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics


Exercise A, Question 7

Question:


A rocket is launched vertically upwards from rest. The initial mass of the rocket and its fuel is 1500 kg . The rocket burns fuel at the rate of 15 kg s^{-1} . The burnt fuel is ejected vertically downwards with a speed of 2000 m s^{-1} relative to the rocket, and burning stops after 60 seconds. Air resistance may be assumed to be negligible. Find the speed of the rocket when the burning stops.

Solution:

At time t



After interval δt



Considering the change in momentum:

$$(m + \delta m)(v + \delta v) + (-\delta m)(v - 2000) - mv = -mg \delta t$$

$$\Rightarrow m \frac{dv}{dt} + 2000 \frac{dm}{dt} = -mg$$

At time t , $m = 1500 - 15t$

$$\Rightarrow (1500 - 15t) \frac{dv}{dt} + 2000 \times -15 = -(1500 - 15t)g$$

Dividing by 15 $\Rightarrow (100 - t) \frac{dv}{dt} - 2000 = -g(100 - t)$

$$\frac{dv}{dt} = -g + \frac{2000}{100 - t}$$

$$\Rightarrow \int_0^v 1 dv = \int_0^{60} -g + \frac{2000}{100 - t} dt$$

$$V = [-gt - 2000 \ln(100 - t)]_0^{60}$$

$$= -60g - 2000 \ln 40 + 0 + 2000 \ln 100 \approx 1240 \text{ m s}^{-1}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 8


Question:

A rocket is launched vertically upwards from rest. The initial mass of the rocket and its fuel is 1200 kg. The rocket burns fuel at the rate of 24 kg s^{-1} . The burnt fuel is ejected vertically downwards with a speed of 2000 m s^{-1} relative to the rocket, and burning stops after 30 seconds. Air resistance may be assumed to be negligible.

- a Find the speed of the rocket when the burning stops.
- b Find the height of the rocket above the launch pad when the burning stops.

Solution:

a At time t



After interval δt



Considering the change in momentum:

$$(m + \delta m)(v + \delta v) + (-\delta m)(v - 2000) - mv = -mg\delta t$$

$$\Rightarrow m \frac{dv}{dt} + 2000 \frac{dm}{dt} = -mg$$

At time t , $m = 1200 - 24t$

$$\Rightarrow (1200 - 24t) \frac{dv}{dt} + 2000 \times -24 = -(1200 - 24t)g$$

$$\text{Dividing by } 24 \Rightarrow (50 - t) \frac{dv}{dt} - 2000 = -g(50 - t)$$

$$\frac{dv}{dt} = -g + \frac{2000}{50 - t} \Rightarrow \int_0^v 1 dv = \int_0^{30} -g + \frac{2000}{50 - t} dt$$

$$V = \left[-gt - 2000 \ln(50 - t) \right]_0^{30}$$

$$= -30g - 2000 \ln 20 + 0 + 2000 \ln 50 \approx 1540 \text{ m s}^{-1}$$

b After time t ,

$$v = \left[-gt - 2000 \ln(50 - t) \right]_0^t = -gt - 2000 \ln \left(\frac{50 - t}{50} \right) = -gt - 2000 \ln \left(1 - \frac{t}{50} \right)$$

so, using integration by parts,

$$x = \int_0^{30} -gt - 2000 \ln \left(1 - \frac{t}{50} \right) dt = \left[-\frac{g}{2} t^2 + 100\,000 \left(1 - \frac{t}{50} \right) \ln \left(1 - \frac{t}{50} \right) - 100\,000 \left(1 - \frac{t}{50} \right) \right]_0^{30}$$

$$= -\frac{900g}{2} + 100\,000 \times \frac{2}{5} \ln \frac{2}{5} - 100\,000 \times \frac{2}{5} + 0 - 0 + 100\,000 \approx 18\,900 \text{ m}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 9

Question:


A rocket is launched vertically upwards from rest. The rocket expels burnt fuel vertically downwards with speed u relative to the rocket. Initially the rocket has mass

M . At time t the rocket has speed v and mass $M\left(1 - \frac{1}{4}t\right)$. Ignore air resistance.


- a Find the speed of the rocket when $t = 2$.
- b Find the height of the rocket above the launch site when $t = 2$.

Solution:

a At time t



After interval δt



$$(m + \delta m)(v + \delta v) + (-\delta m)(v - u) - mv = -mg \delta t$$

$$m \delta v + \delta m \delta v + u \delta m = -mg \delta t$$

$$\Rightarrow m \frac{dv}{dt} + u \frac{dm}{dt} = -mg$$

$$m = M \left(1 - \frac{1}{4}t \right)$$

$$\Rightarrow M \left(1 - \frac{1}{4}t \right) \frac{dv}{dt} + u \left(-\frac{1}{4}M \right) = -M \left(1 - \frac{1}{4}t \right) g$$

$$\frac{dv}{dt} = \frac{\frac{1}{4}u}{1 - \frac{1}{4}t} - g = \frac{u}{4-t} - g$$

$$\frac{dv}{dt} = \frac{u}{4-t} - g \Rightarrow v = \int \frac{u}{4-t} - g dt = -u \ln |4-t| - gt + C$$

$$t = 0, v = 0 \Rightarrow 0 = -u \ln 4 + C$$

$$v = u \ln \left| \frac{4}{4-t} \right| - gt = u \ln \frac{4}{2} - 2g = u \ln 2 - 2g \text{ when } t = 2$$

b $v = \frac{dx}{dt} = u \ln \left(\frac{4}{4-t} \right) - gt, t < 4$

Using integration by parts

$$\Rightarrow x = \int u \ln \left(\frac{4}{4-t} \right) - gt dt = \int u \ln 4 - u \ln (4-t) - gt dt$$

$$= (u \ln 4)t + u(4-t) \ln (4-t) - u(4-t) - \frac{1}{2}gt^2 + C$$

$$t = 0, x = 0 \Rightarrow 0 = 0 + 4u \ln 4 - 4u + C, C = 4u - 4u \ln 4$$

$$\text{When } t = 2, x = 2 \times u \ln 4 + 2u \ln 2 - 2u - 2g + 4u - 4u \ln 4 = -2u \ln 2 + 2u - 2g$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 10


Question:

A rocket uses fuel at a rate $\lambda \text{ kg s}^{-1}$. The rocket moves forwards by expelling used fuel backwards from the rocket with speed 2500 m s^{-1} relative to the rocket. At time t the rocket is moving with speed v and the combined mass of the rocket and its fuel is m . The rocket starts from rest at time $t = 0$ with a total mass $10\,000 \text{ kg}$ and reaches a final speed 5000 m s^{-1} after 200 seconds. Given that no external forces act on the rocket


- a show that $m \frac{dv}{dt} = 2500 \lambda$,
- b find the value of $\lambda \text{ kg s}^{-1}$.

Solution:

a At time t



After an interval δt



$$\begin{aligned}(m + \delta m)(v + \delta v) + (-\delta m)(v - 2500) - mv &= 0 \\ \Rightarrow m\delta v + \delta m\delta v + \delta m2500 &= 0 \\ m\frac{dv}{dt} + 2500\frac{dm}{dt} &= 0\end{aligned}$$

but we are told that $\frac{dm}{dt} = -\lambda$

$$\text{so } m\frac{dv}{dt} - 2500\lambda = 0, m\frac{dv}{dt} = 2500\lambda$$

b The initial mass is 10 000 and $\frac{dm}{dt} = -\lambda$, so

$$(10\,000 - \lambda t)\frac{dv}{dt} = 2500\lambda$$

Separating the variables

$$\begin{aligned}\Rightarrow \int_0^{5000} \frac{1}{v} dv &= \int_0^{200} \frac{2500\lambda}{10\,000 - \lambda t} dt \\ 5000 &= \left[-2500 \ln(10\,000 - \lambda t) \right]_0^{200} = -2500 \ln \left(\frac{10\,000 - 200\lambda}{10\,000} \right) \\ \Rightarrow 2 &= \ln \left(\frac{50}{50 - \lambda} \right), e^2 = \frac{50}{50 - \lambda}, 50 - \lambda = 50e^{-2}, \\ \lambda &= 50(1 - e^{-2}) \approx 43.2\end{aligned}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 11

Question:


A rocket uses fuel at a rate λ . The rocket moves forwards by expelling used fuel backwards from the rocket with speed 2000 m s^{-1} relative to the rocket. At time t the rocket is moving with speed v and the combined mass of the rocket and its fuel is m . The rocket starts from rest at time $t = 0$ with a total mass $12\,000 \text{ kg}$ and reaches a speed of 5000 m s^{-1} after 3 minutes.

Given that no external forces act on the rocket

- a show that $m \frac{dv}{dt} = 2000 \lambda$,
- b find the greatest and the least acceleration of the vehicle during these three minutes.

Solution:

a At time t



After an interval δt



$$\begin{aligned}(m + \delta m)(v + \delta v) + (-\delta m)(v - 2000) - mv &= 0 \\ \Rightarrow m\delta v + \delta m\delta v + \delta m2000 &= 0 \\ m\frac{dv}{dt} + 2000\frac{dm}{dt} &= 0\end{aligned}$$

but we are told that $\frac{dm}{dt} = -\lambda$

$$\text{so } m\frac{dv}{dt} - 2000\lambda = 0, m\frac{dv}{dt} = 2000\lambda$$

b The initial mass is 12 000 and $\frac{dm}{dt} = -\lambda$, so

$$(12000 - \lambda t)\frac{dv}{dt} = 2000\lambda$$

$$\int_0^{5000} 1 dv = \int_0^{180} \frac{2000\lambda}{12000 - \lambda t} dt$$

$$5000 = \left[-2000 \ln(12000 - \lambda t) \right]_0^{180} = -2000 \ln \left(\frac{12000 - 180\lambda}{12000} \right)$$

$$\Rightarrow \frac{5}{2} = \ln \left(\frac{200}{200 - 3\lambda} \right), e^{\frac{5}{2}} = \frac{200}{200 - 3\lambda}, 200 - 3\lambda = 200e^{-\frac{5}{2}},$$

$$\lambda = \frac{200}{3} \left(1 - e^{-\frac{5}{2}} \right) \approx 61.2$$

$$\frac{dv}{dt} = \frac{2000\lambda}{m} \Rightarrow \text{min acceleration} = \frac{2000 \times 61.2}{12000} = 10.2 \text{ m s}^{-2}$$

$$\text{max acceleration} = \frac{2000 \times 61.2}{12000 - 180 \times 61.2} = 124 \text{ m s}^{-2}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 12




Question:

A particle falls from rest under gravity through a stationary cloud. At time t the particle has fallen a distance x , has mass m and speed v . The mass of the particle increases by accretion from the cloud at a rate of kmv , where k is a constant. Ignore air resistance. Show that

a $kv^2 = g(1 - e^{-2kx})$,

b $x = \frac{1}{k} \ln \left[\cosh \left(\sqrt{kg} t \right) \right]$.

Solution:

a At time t After an interval δt :

$$[(m + \delta m)(v + \delta v)] - [mv + \delta m \times 0] = (m + \delta m)g \delta t$$

$$\Rightarrow m \frac{dv}{dt} + v \frac{dm}{dt} = mg$$

But we are told that $\frac{dm}{dt} = mkv$, so $m \frac{dv}{dt} + v \times mkv = mg$

$$\frac{dv}{dt} = g - kv^2, \Rightarrow v \frac{dv}{dx} = g - kv^2$$

$$\int \frac{v}{g - kv^2} dv = \int 1 dx \Rightarrow -\frac{1}{2k} \ln(g - kv^2) = x + C$$

$$x = 0, v = 0 \Rightarrow -\frac{1}{2k} \ln g = C \Rightarrow x = -\frac{1}{2k} \ln \left(\frac{g - kv^2}{g} \right)$$

$$e^{2kx} = \frac{g}{g - kv^2}, \quad (g - kv^2)e^{2kx} = g, \quad kv^2 = g(1 - e^{-2kx})$$

$$\text{b} \quad v^2 = \frac{g}{k}(1 - e^{-2kx}), \quad v = \sqrt{\frac{g}{k}(1 - e^{-2kx})} = \frac{dx}{dt}$$

$$\int \sqrt{\frac{g}{k}} dt = \int \frac{1}{\sqrt{1 - e^{-2kx}}} dx = \int \frac{e^{kx}}{\sqrt{e^{2kx} - 1}} dx$$

$$\Rightarrow \text{by using the substitution } \cosh u = e^{kx}, \sinh u \cdot \frac{du}{dx} = k e^{kx}$$

$$\sqrt{\frac{g}{k}} t = \int \frac{e^{kx}}{\sqrt{e^{2kx} - 1}} dx = \frac{1}{k} \int \frac{\sinh u}{\sqrt{\cosh^2 u - 1}} du = \frac{1}{k} \int 1 du = \frac{u}{k} + C$$

$$t = 0, x = 0 \Rightarrow \cosh u = 1, u = \cosh^{-1} 1 = 0, \Rightarrow C = 0$$

$$\Rightarrow \sqrt{kg} t = u, \quad \cosh(\sqrt{kg} t) = e^{kx}, \quad kx = \ln[\cosh(\sqrt{kg} t)], \quad x = \frac{1}{k} \ln[\cosh(\sqrt{kg} t)]$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 13

Question:

A raindrop falls through a stationary cloud. When the raindrop has fallen distance x it has mass m and speed v . The mass increases uniformly by accretion so that $m = M(1+kx)$.


Given that $v=0$ when $x=0$, find an expression, in terms of M , k and x for the kinetic energy of the raindrop when it has fallen a distance x . Ignore air resistance.

Solution:

At time t



After an interval δt :



Impulse momentum: $[(m + \delta m)(v + \delta v)] - [mv] = (m + \delta m)g\delta t$

$$m \frac{\delta v}{\delta x} + v \frac{\delta m}{\delta x} + \frac{\delta m \delta v}{\delta x} = mg \frac{\delta t}{\delta x} + \delta mg \frac{\delta t}{\delta x}$$

$$m \frac{dv}{dx} + v \frac{dm}{dx} = mg \frac{dt}{dx} = \frac{mg}{v}, \quad mv \frac{dv}{dx} + v^2 \frac{dm}{dx} = mg$$

Substituting for m :

$$M(1+kx)v \frac{dv}{dx} + v^2 kM = M(1+kx)g$$

$$v \frac{dv}{dx} + v^2 \frac{k}{1+kx} = g, \quad 2v \frac{dv}{dx} + \frac{2k}{1+kx} v^2 = 2g$$

A linear differential equation in v^2 .

Integrating factor $e^{\int \frac{2k}{1+kx} dx} = e^{2 \ln(1+kx)} = (1+kx)^2$

$$\Rightarrow \frac{d}{dx} [v^2(1+kx)^2] = 2g(1+kx)^2, \quad v^2(1+kx)^2 = \frac{2g}{3k}(1+kx)^3 + C$$

$$x=0, v=0 \Rightarrow 0 = \frac{2g}{3k} + C, \quad v^2 = \frac{2g}{3k}(1+kx) - \frac{2g}{3k(1+kx)^2},$$

$$\text{so K.E.} = \frac{1}{2}mv^2 = \frac{1}{2}M(1+kx) \left[\frac{2g}{3k}(1+kx) - \frac{2g}{3k(1+kx)^2} \right]$$

$$= \frac{Mg}{3k} \left[(1+kx)^2 - \frac{1}{(1+kx)} \right]$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 14

Question:

A rocket is on the ground facing vertically upwards. When launched it propels itself by ejecting mass backwards with speed u relative to the rocket at a constant rate k per unit time. The initial mass of the rocket is M . Ignore air resistance.

a Explain why it is necessary for $ku > Mg$.


Given that $ku > Mg$,

b show that the velocity of the rocket after time t is $-u \ln \left(1 - \frac{kt}{M} \right) - gt$,


c find the height of the rocket above the ground when the mass of the rocket has reduced by one third of its initial value.

Solution:

a At time t



After an interval δt



$$\text{Change in momentum} = (m + \delta m)(v + \delta v) + (-\delta m)(v - u) - mv = -mg\delta t$$

$$\Rightarrow m \frac{dv}{dt} + u \frac{dm}{dt} = -mg$$

$$m = M - kt \Rightarrow (M - kt) \frac{dv}{dt} + u(-k) = -(M - kt)g$$

$$\frac{dv}{dt} = \frac{ku}{M - kt} - g$$

If the rocket is to be able to launch then when $t = 0$, $\frac{dv}{dt} > 0$

$$\frac{ku}{M} - g > 0, \text{ i.e. } ku > Mg$$

$$\text{b } \frac{dv}{dt} = \frac{ku}{M - kt} - g \Rightarrow v = -u \ln(M - kt) - gt + C$$

$$t = 0, v = 0 \Rightarrow 0 = -u \ln M + C$$

$$\begin{aligned} \Rightarrow v &= -u \ln(M - kt) - gt + u \ln M = -u \ln \left(\frac{M - kt}{M} \right) - gt \\ &= -u \ln \left(1 - \frac{kt}{M} \right) - gt \end{aligned}$$

$$\text{c } v = -u \ln \left(1 - \frac{kt}{M} \right) - gt = \frac{dx}{dt}$$

$$\Rightarrow x = \int -u \ln \left(1 - \frac{kt}{M} \right) - gt dt = \frac{uM}{k} \left[\left(1 - \frac{kt}{M} \right) \ln \left(1 - \frac{kt}{M} \right) - \left(1 - \frac{kt}{M} \right) \right] - \frac{gt^2}{2} + C$$

(using integration by parts of $\ln \left(1 - \frac{kt}{M} \right)$)

$$t = 0, x = 0 \Rightarrow 0 = \frac{uM}{k} \times -1 + C$$

$$m = \frac{2}{3}M = M - kt, t = \frac{M}{3k}$$

$$x = \frac{uM}{k} \left[\frac{2}{3} \ln \frac{2}{3} - \frac{2}{3} \right] - \frac{gM^2}{18k^2} + \frac{uM}{k} = \frac{uM}{3k} \left[2 \ln \frac{2}{3} + 1 - \frac{gM}{6ku} \right]$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 15

Question:


At time $t = 0$ a particle is projected vertically upwards. Initially the particle has mass M and speed gT , where T is a constant. At time t the speed of the particle is v and its mass is $Me^{\frac{t}{2T}}$. Ignore air resistance. If the added material is at rest when it is acquired, show that

a $\frac{d}{dt} \left(Mve^{\frac{t}{2T}} \right) = -Mge^{\frac{t}{2T}},$


b the particle has mass $\frac{3M}{2}$ at its highest point.

Solution:

a At time t v



After an interval δt



Change in momentum: $[(m + \delta m)(v + \delta v)] - [mv + \delta m \times 0] = -(m + \delta m)g \delta t$

Taking the limit as $\delta t \rightarrow 0$

$$m \frac{dv}{dt} + v \frac{dm}{dt} = -mg, \text{ i.e. } \frac{d}{dt}(mv) = -mg$$

$$\frac{d}{dt} \left(Mve^{\frac{t}{2T}} \right) = -Me^{\frac{t}{2T}} g = -Mge^{\frac{t}{2T}}$$

$$\text{b } \frac{d}{dt} \left(Mve^{\frac{t}{2T}} \right) = -Me^{\frac{t}{2T}} g, \left(Mve^{\frac{t}{2T}} \right) = \int -Mge^{\frac{t}{2T}} dt$$

$$\Rightarrow Mve^{\frac{t}{2T}} = -2MgTe^{\frac{t}{2T}} + C$$

$$t = 0, v = gT, C = 3MgT$$

$$\Rightarrow Mve^{\frac{t}{2T}} = -2MgTe^{\frac{t}{2T}} + 3MgT, \Rightarrow ve^{\frac{t}{2T}} = -2gTe^{\frac{t}{2T}} + 3gT$$

At the highest point, $v = 0$, so $0 = -2gTe^{\frac{t}{2T}} + 3gT, e^{\frac{t}{2T}} = \frac{3}{2}$ and

$$\text{mass} = Me^{\frac{t}{2T}} = \frac{3M}{2}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics


Exercise A, Question 16

Question:


At time $t = 0$ a particle is projected vertically upwards from the ground. Initially the particle has mass M and speed $2gT$, where T is a constant. At time t the mass of the particle is $Me^{\frac{t}{T}}$. If the added material is at rest when it is acquired, show that the highest point reached by the particle is $gT^2(2 - \ln 3)$ above the ground. Ignore air resistance.

Solution:

At time t



After an interval δt



Change in momentum: $[(m + \delta m)(v + \delta v)] - [mv + \delta m \times 0] = -(m + \delta m)g\delta t$

Taking the limit as $\delta t \rightarrow 0$

$$m \frac{dv}{dt} + v \frac{dm}{dt} = -mg, \text{ i.e. } \frac{d}{dt}(mv) = -mg$$

$$\frac{d}{dt} \left(Mve^{\frac{t}{T}} \right) = -Me^{\frac{t}{T}}g = -Mge^{\frac{t}{T}}$$

$$\frac{d}{dt} \left(Mve^{\frac{t}{T}} \right) = -Mge^{\frac{t}{T}}, \quad \left(Mve^{\frac{t}{T}} \right) = \int -Mge^{\frac{t}{T}} dt$$

$$\Rightarrow Mve^{\frac{t}{T}} = -MgTe^{\frac{t}{T}} + C$$

$$t = 0, v = 2gT, C = 3MgT$$

$$\Rightarrow Mve^{\frac{t}{T}} = -MgTe^{\frac{t}{T}} + 3MgT, \Rightarrow v = -gT + 3gTe^{-\frac{t}{T}}$$

$$\Rightarrow \frac{dx}{dt} = -gT + 3gTe^{-\frac{t}{T}}, x = -gTt - 3gT^2e^{-\frac{t}{T}} + C$$

$$t = 0, x = 0, \Rightarrow C = 3gT^2, x = -gTt - 3gT^2e^{-\frac{t}{T}} + 3gT^2$$

At the highest point, $v = 0, \Rightarrow e^{-\frac{t}{T}} = \frac{1}{3}, -\frac{t}{T} = \ln \frac{1}{3}, t = T \ln 3$

$$\Rightarrow x = -gT \cdot T \ln 3 - 3gT^2 \cdot \frac{1}{3} + 3gT^2$$

$$= gT^2(2 - \ln 3)$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics


Exercise A, Question 17

Question:


At time $t = 0$ a particle is projected vertically upwards. Initially the particle has mass M and speed gT , where T is a constant. At time t the mass of the particle is $Me^{\frac{t}{T}}$. If the added material is falling with constant speed gT when it is acquired, show that the particle has mass $\frac{3M}{2}$ at its highest point. Ignore air resistance.

Solution:

At time t



After an interval δt



Change in momentum:

$$[(m + \delta m)(v + \delta v)] - [mv - \delta m \times gT] = -(m + \delta m)g\delta t$$

$$\Rightarrow v \frac{dm}{dt} + m \frac{dv}{dt} = -mg - gT \frac{dm}{dt}$$

$$m = Me^{\frac{t}{T}} \Rightarrow \frac{dm}{dt} = \frac{M}{T} e^{\frac{t}{T}}, \frac{d}{dt}(mv) = -mg - gMe^{\frac{t}{T}}$$

$$\frac{d}{dt} \left(Me^{\frac{t}{T}} v \right) = -Me^{\frac{t}{T}} g - gMe^{\frac{t}{T}} = -2Mge^{\frac{t}{T}}$$

$$\Rightarrow Me^{\frac{t}{T}} v = \int -2gMe^{\frac{t}{T}} dt = -2gTMe^{\frac{t}{T}} + C$$

$$t = 0, v = gT \Rightarrow MgT = -2MgT + C, C = 3MgT$$

$$\Rightarrow Me^{\frac{t}{T}} v = -2gTMe^{\frac{t}{T}} + 3MgT$$

$$v = 0, \Rightarrow 2gTMe^{\frac{t}{T}} = 3MgT, e^{\frac{t}{T}} = \frac{3}{2}, \Rightarrow m = \frac{3M}{2}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 18


Question:

A particle of mass M is projected vertically upwards in a cloud. During the motion the particle absorbs moisture from the stationary cloud so that when the particle is at distance x above the point of projection, moving with speed v , it has mass $M(1 + \alpha x)$, where α is a constant. The initial speed of the particle is $\sqrt{2gk}$. Ignore air resistance.


- a** Show that $2v \frac{dv}{dx} + \frac{2\alpha}{1 + \alpha x} v^2 = -2g$.
- b** Show that at the greatest height, $h, (1 + \alpha h)^3 = 1 + 3k\alpha$.

Solution:

a At time t



After an interval δt :



Change in momentum: $[(m + \delta m)(v + \delta v)] - [mv + \delta m \times 0] = -(m + \delta m)g\delta t$

Taking the limit as $\delta t \rightarrow 0$

$$m \frac{dv}{dt} + v \frac{dm}{dt} = -mg$$

$$m = M(1 + \alpha x) \Rightarrow M(1 + \alpha x) \frac{dv}{dt} + vM\alpha \frac{dx}{dt} = -M(1 + \alpha x)g$$

Using $\frac{dv}{dt} = v \frac{dv}{dx}$ and $\frac{dx}{dt} = v$

$$\frac{dv}{dt} + v \frac{\alpha}{(1 + \alpha x)} \frac{dx}{dt} = -g, v \frac{dv}{dx} + v^2 \frac{\alpha}{(1 + \alpha x)} = -g$$

$$\Rightarrow 2v \frac{dv}{dx} + \frac{2\alpha}{(1 + \alpha x)} v^2 = -2g$$

b Multiply through the differential equation by the integrating factor (since the differential equation is linear differential equation in v^2)

$$\text{I.F.} = e^{\int \frac{2\alpha}{(1 + \alpha x)} dx} = e^{2 \ln(1 + \alpha x)} = (1 + \alpha x)^2$$

$$\begin{aligned} \Rightarrow \frac{d}{dx} [v^2 (1 + \alpha x)^2] &= -2g(1 + \alpha x)^2, v^2 (1 + \alpha x)^2 = \int -2g(1 + \alpha x)^2 dx \\ &= -\frac{2g}{3\alpha} (1 + \alpha x)^3 + C \end{aligned}$$

$$x = 0, v = \sqrt{2gk} \Rightarrow 2gk = -\frac{2g}{3\alpha} + C, \quad C = 2g \left(k + \frac{1}{3\alpha} \right)$$

$$\text{At the highest point, } v = 0 \Rightarrow 0 = -\frac{2g}{3\alpha} (1 + \alpha h)^3 + 2g \left(k + \frac{1}{3\alpha} \right)$$

$$\frac{1}{3\alpha} (1 + \alpha h)^3 = \left(k + \frac{1}{3\alpha} \right)$$

$$\text{and therefore } (1 + \alpha h)^3 = 1 + 3k\alpha$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 19

Question:


A body of mass $3M$ contains combustible and non-combustible material in the ratio $2 : 1$. The body is initially at rest and falls freely under gravity. At time t the body has speed v .

The combustible part burns at a constant rate of λM per second, where λ is a constant. The burning material is ejected vertically upwards with constant speed u relative to the body. Assuming that air resistance may be neglected,

- a** show that $\frac{dv}{dt} = \frac{\lambda u}{3 - \lambda t} + g$,
- b** find how far the body has fallen when all the combustible material has been used up.

Solution:

a At time t



After an interval δt :



$$\Rightarrow (m + \delta m)(v + \delta v) + (-\delta m)(v - u) - mv = mg \delta t$$

$$m \delta v + \delta m \delta v + \delta m u = mg \delta t, \quad m \frac{dv}{dt} + u \frac{dm}{dt} = mg$$

$$m = M(3 - \lambda t), \Rightarrow M(3 - \lambda t) \frac{dv}{dt} + u(-\lambda M) = M(3 - \lambda t)g$$

$$(3 - \lambda t) \frac{dv}{dt} - \lambda u = (3 - \lambda t)g, \quad \frac{dv}{dt} = \frac{\lambda u}{3 - \lambda t} + g$$

b $\frac{dv}{dt} = \frac{\lambda u}{3 - \lambda t} + g \Rightarrow v = \int \frac{\lambda u}{3 - \lambda t} + g dt = -u \ln(3 - \lambda t) + gt + C, (\lambda t < 3)$

$$t = 0, v = 0 \Rightarrow 0 = -u \ln 3 + C$$

$$\Rightarrow v = -u \ln \left(\frac{3 - \lambda t}{3} \right) + gt = -u \ln \left(1 - \frac{\lambda t}{3} \right) + gt = \frac{dx}{dt}$$

$$\Rightarrow x = \int -u \ln \left(1 - \frac{\lambda t}{3} \right) + gt dt = \frac{3u}{\lambda} \left[\left(1 - \frac{\lambda t}{3} \right) \ln \left(1 - \frac{\lambda t}{3} \right) - \left(1 - \frac{\lambda t}{3} \right) \right] + \frac{gt^2}{2} + C$$

$$t = 0, x = 0 \Rightarrow 0 = -\frac{3u}{\lambda} + C$$

All combustible material used $\Rightarrow m = M(3 - \lambda t) = M, \quad t = \frac{2}{\lambda}$

$$\begin{aligned} \Rightarrow x &= \frac{3u}{\lambda} \left[\left(1 - \frac{\lambda t}{3} \right) \ln \left(1 - \frac{\lambda t}{3} \right) - \left(1 - \frac{\lambda t}{3} \right) \right] + \frac{gt^2}{2} + \frac{3u}{\lambda} \\ &= \frac{3u}{\lambda} \left[\left(1 - \frac{\lambda 2}{3\lambda} \right) \ln \left(1 - \frac{\lambda 2}{3\lambda} \right) - \left(1 - \frac{\lambda 2}{3\lambda} \right) \right] + \frac{g4}{2\lambda^2} + \frac{3u}{\lambda} \\ &= \frac{3u}{\lambda} \left[\frac{1}{3} \ln \frac{1}{3} - \frac{1}{3} \right] + \frac{2g}{\lambda^2} + \frac{3u}{\lambda} = \frac{3u}{\lambda} \left[\frac{1}{3} \ln \frac{1}{3} + \frac{2}{3} \right] + \frac{2g}{\lambda^2} \\ &= \frac{u}{\lambda} \left[\ln \frac{1}{3} + 2 \right] + \frac{2g}{\lambda^2} = \frac{u}{\lambda} (2 - \ln 3) + \frac{2g}{\lambda^2} \end{aligned}$$

Solutionbank M5

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 20

Question:

A spherical hailstone falls vertically through a stationary cloud from rest under gravity. The initial radius of the hailstone is a . As the hailstone falls its volume increases through condensation. When the radius of the hailstone is r , the rate of increase of volume is $4\pi r^2\lambda$ and the hailstone is falling with speed v . Ignore air resistance.

a Show that, at time t , $r = a + \lambda t$.

b Show that $\frac{dv}{dt} = g - \frac{3\lambda v}{r}$.

c Find the speed of the particle when $t = \frac{a}{2\lambda}$.


Solution:

- a For the sphere, $\frac{dV}{dt} = 4\pi r^2 \lambda$, but


$$V = \frac{4}{3}\pi r^3 \Rightarrow 4\pi r^2 \frac{dr}{dt} = 4\pi r^2 \lambda$$

$$\Rightarrow \frac{dr}{dt} = \lambda, r = a + \lambda t$$

- b At time t



After time δt :



$$[(m + \delta m)(v + \delta v)] - [mv + \delta m \times 0] = (m + \delta m)g \delta t$$

$$\Rightarrow m \frac{\delta v}{\delta t} + v \frac{\delta m}{\delta t} + \frac{\delta m \delta v}{\delta t} = mg + g \delta m, \text{ so } m \frac{dv}{dt} + v \frac{dm}{dt} = mg$$

The mass of the hailstone is $\rho \times \frac{4}{3}\pi r^3$ since mass is proportional to volume

$$\Rightarrow \frac{dm}{dt} = 4\rho\pi r^2 \frac{dr}{dt} = 4\rho\pi r^2 \times \lambda$$

$$\Rightarrow \rho \times \frac{4}{3}\pi r^3 \frac{dv}{dt} + v \times 4\rho\lambda\pi r^2 = \rho \times \frac{4}{3}\pi r^3 g$$

and therefore $\frac{dv}{dt} = g - \frac{3\lambda v}{r}$

c $\frac{dv}{dt} = g - \frac{3\lambda v}{r} = g - \frac{3\lambda v}{a + \lambda t}, \frac{dv}{dt} + \frac{3\lambda v}{a + \lambda t} = g$

Using the integrating factor $e^{\int \frac{3\lambda}{a + \lambda t} dt} = e^{3\ln(a + \lambda t)} = (a + \lambda t)^3$:

$$v(a + \lambda t)^3 = \int g(a + \lambda t)^3 dt = \frac{g}{4\lambda}(a + \lambda t)^4 + C$$

$$t = 0, v = 0, 0 = \frac{ga^4}{4\lambda} + C, \quad v(a + \lambda t)^3 = \frac{g}{4\lambda}(a + \lambda t)^4 - \frac{ga^4}{4\lambda}$$

$$v = \frac{g(a + \lambda t)}{4\lambda} - \frac{ga^4}{4\lambda(a + \lambda t)^3}$$

$$t = \frac{a}{2\lambda} \Rightarrow v = \frac{g\left(a + \lambda \frac{a}{2\lambda}\right)}{4\lambda} - \frac{ga^4}{4\lambda\left(a + \lambda \frac{a}{2\lambda}\right)^3} = \frac{3ag}{8\lambda} - \frac{2ga}{27\lambda} = \frac{65ag}{216\lambda}$$

